
APPENDICES

This page intentionally left blank.

Appendix A
**NEFSC Bycatch Estimation Methodology: Allocation,
Precision, and Accuracy**

This page intentionally left blank.

**NEFSC Bycatch Estimation Methodology:
Allocation, Precision, and Accuracy**

by

Paul J. Rago, Susan E. Wigley, and Michael J. Fogarty

August 2005

Recent Issues in This Series

- 04-07 **Identification and Description of the Common Sponges of Jeffreys Ledge as an Aid in Field Operations.** By K. McCarthy. April 2004.
- 04-08 **Revised Procedures for Calculating Regional Average Water Properties for Northeast Fisheries Science Center Cruises.** By D.G. Mountain, M.H. Taylor, and C. Bascuñán. April 2004.
- 04-09 **Estimation of Striped Bass Discards in the Multispecies Groundfish Fishery during the 2002 Fishing Year (May 2002 - April 2003).** By G.R. Shepherd. June 2004.
- 04-10a **39th Northeast Regional Stock Assessment Workshop (39th SAW) Assessment Summary Report.** [By Northeast Regional Stock Assessment Workshop No. 39.] July 2004.
- 04-10b **39th Northeast Regional Stock Assessment Workshop (39th SAW) Assessment Report.** [By Northeast Regional Stock Assessment Workshop No. 39.] July 2004.
- 04-11 **Bycatch of Sea Turtles in the Mid-Atlantic Sea Scallop (*Placopecten magellanicus*) Dredge Fishery during 2003.** By K.T. Murray. August 2004.
- 04-12 **Description of the 2003 Oceanographic Conditions on the Northeast Continental Shelf.** By C. Bascuñán, M.H. Taylor, and J.P. Manning. September 2004.
- 04-13 **Ninth Flatfish Biology Conference, December 1-2, 2004, Water's Edge Resort, Westbrook, Connecticut.** By R. Mercaldo-Allen (chair), A. Calabrese, D.J. Danila, M.S. Dixon, A. Jearld, D.J. Pacileo, C. Powell, and S.J. Sutherland, steering committee members. November 2004.
- 04-14 **Northeast Fisheries Science Center Publications, Reports, and Abstracts for Calendar Year 2003.** By L. Garner. November 2004.
- 05-01 **Results from the 2004 Cooperative Survey of Atlantic Surfclams.** By J.R. Weinberg, E.N. Powell, C. Pickett, V.A. Nordahl, Jr., and L.D. Jacobson. February 2005.
- 05-02 **Proceedings of a Workshop to Review and Evaluate the Design and Utility of Fish Mark - Recapture Projects in the Northeastern United States; October 19-21, 2004; Nonantum Resort, Kennebunkport, Maine.** By Workshop Organizing Committee (S. Tallack, editor, P. Rago, chairperson, T. Brawn, workshop coordinator, and (alphabetically) S. Cadrin, J. Hoey, and L. Taylor Singer. March 2005.
- 05-03 **Description of the 2004 Oceanographic Conditions on the Northeast Continental Shelf.** By M.H. Taylor, C. Bascuñán, and J.P. Manning. April 2005.
- 05-04 **40th SAW Assessment Report.** By 40th Northeast Regional Stock Assessment Workshop. April 2005.
- 05-05 **Effectiveness of a Square-Mesh Escape Panel in Reducing Finfish Bycatch in a Small-Mesh Bottom Trawl Used in the Longfin Inshore Squid (*Loligo pealeii*) Fishery.** By L. Hendrickson. June 2005.
- 05-06 **Use of the Historic Area Remediation Site by Black Sea Bass and Summer Flounder.** By M.C. Fabrizio, J.P. Pessutti, J.P. Manderson, A.F. Drohan, and B.A. Phelan. June 2005.
- 05-07 **Benthic Macrofauna and Associated Hydrographic Observations Collected in Newark Bay, New Jersey, between June 1993 and March 1994.** By L.L. Stehlik, S.J. Wilk, R.A. Pikanowski, D.G. McMillan, and E.M. MacHaffie. July 2005.
- 05-08 **Mortality and Serious Injury Determinations for Large Whale Stocks along the Eastern Seaboard of the United States, 1999-2003.** By T.V.N. Cole, D.L. Hartley, and R.L. Merrick. July 2005.

NEFSC Bycatch Estimation Methodology: Allocation, Precision, and Accuracy

by

Paul J. Rago^{1,2}, Susan E. Wigley^{1,3}, and Michael J. Fogarty^{1,4}

Postal Address: ¹National Marine Fisheries Serv., 166 Water St., Woods Hole, MA 02543

E-Mail Addresses: ²Paul.Rago@noaa.gov, ³Susan.Wigley@noaa.gov, ⁴Michael.Fogarty@noaa.gov

**U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts**

August 2005

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review but no technical or copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/series/crdlist.htm>. The electronic version will be available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the desired document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

This document's publication history is as follows: manuscript submitted for review -- July 28, 2005; manuscript accepted through technical review -- July 28, 2005; manuscript accepted through policy review -- August 2, 2005; and final copy submitted for publication -- August 11, 2005. This document may be cited as:

Rago, P.J.; Wigley, S.E.; Fogarty, M.J. 2005. NEFSC bycatch estimation methodology: allocation, precision, and accuracy. *U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc.* 05-09; 44 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.

Table of Contents

Executive Summary	v
Introduction	1
Background	1
Definition of Strata – Fishery Identification	2
Data Sources.....	4
Fishing Vessel Trip Report.....	4
Northeast Fisheries Observer Program Data.....	5
Optimization Data Set	6
Imputation Rules for Unobserved Fisheries	6
Sampling Theory and Optimization Models	9
Measures of Overlap	11
Observers Days at Sea Constraints.....	12
Application of the Model	16
Precision, Bias and Sampling Intensity: A Rebuttal to E.A. Babcock et al	18
An Evaluation of Bias in the Northeast Fisheries Observer Program.....	21
Sources of Uncertainty	22
Acknowledgments	24
References	25

List of Tables

Table 1.	The variables, their description, their associated species group, data source, and units of the input data set of the optimization algorithm.	27
Table 2.	Number of trips, by strata, in the Fishing Vessel Trip Reports and Northeast Fisheries Observer Program data sets used in the 2005 sea day optimization.....	29
Table 3.	Summary of fleet sectors, by species group, that are imputed and not imputed; blank cells indicate no fleet activity	30
Table 4.	Summary of contingency table analyses of spatial distribution of VTR and observed trips	31

List of Figures

Figure 1.	An overview of the optimization process used to allocate sea days to fisheries in the Northeast region.....	33
Figure 2.	Number of trips in the 2003/2004 Vessel Trip Report by data subsets for otter trawl, gillnet and longline trips	34
Figure 3.	Number of trips and sea days in the 2003/2004 Northeast Fisheries Observer Program, by data subsets for otter trawl, gillnet and longline trips	35
Figure 4.	The sampling fraction of 2003/2004 Observed trips to Vessel Trip Report trips, by data subset for otter trawl, gillnet and longline trips	36
Figure 5.	Comparison of average kept pounds of groundfish in the Northeast Fisheries Observer Program and Vessel Trip Report data sets for 2003/2004	37

Figure 6. The distribution of differences between the average kept pounds and the standard deviation of average kept pounds of groundfish in the Northeast Fisheries Observer Program and the Vessel Trip Report data for 2003/2004 38

Figure 7. Comparison of average trip duration for trips that caught groundfish in the Northeast Fisheries Observer Program and Vessel Trip Report data sets for 2003/2004 39

Figure 8. The distribution of differences in average trip duration and the standard deviation of average trip duration of trips that caught groundfish in the Northeast Fisheries Observer Program and the Vessel Trip Report data for 2003/2004..... 40

Figure 9. Locations of otter trawl fishing effort in 2003 from vessels using VMS 41

Figure 10. The optimized coefficient of variation of the discard to kept ratio for New England groundfish over a range of sea days; 2,708 sea days are allocated to cover New England groundfish fisheries in 2005 42

Figure 11. The 2003/2004 point estimates of the coefficient of variation of the discard to kept ratio for New England groundfish caught with otter trawl gear, and the expected coefficient of variation of the discard to kept ratio over a range of sample sizes..... 43

Figure 12. The 2003/2004 point estimates of the coefficient of variation of the discard to kept ratio for New England groundfish caught with gillnet gear, and the expected coefficient of variation of the discard to kept ratio over a range of sample sizes..... 44

Executive Summary

This report describes the standardized methodology used to estimate bycatch rates of finfish by commercial fisheries in the Northeast. In this report, bycatch is defined as the observed discarded catch, summed over from eleven different groundfish species. Estimates of unobserved discards are not considered. All retained catches are included whether or not the catches were incidental to the target species. Emphasis is placed on the methods used to define the sampling frame (i.e., the population of commercial fishing trips to be sampled), appropriate stratification, and efficient allocation of sampling effort to these strata. Efficient allocation of sampling effort within a stratified survey design improves the precision of the estimate of overall discard rates. Accuracy of sample estimates is evaluated by comparing various performance measures (e.g., landings, trip duration) between vessels with and without observers present. Although formal statistical distinctions between accuracy and bias of estimators and estimates can be made, in this report we use the terms interchangeably and less formally. A biased estimator is inaccurate; an accurate estimator is unbiased.

This report focuses on bycatch estimates based on discard to kept ratios. Use of this ratio is appropriate for trawl, gillnet and longline fisheries in the Northeast US. A formal assessment of bycatch estimates based on the ratio of discards to fishing effort is not considered in this report. Estimators based on ratios of total discard to fishing effort are more appropriate for fisheries that do not target groundfish, such as the sea scallop and herring fisheries. Evaluations of groundfish bycatch in these fisheries are being conducted by technical committees for their respective fishery management plans.

The Northeast Fisheries Science Center allocates observer sea days to monitor bycatch in commercial fisheries along the Northeast coast. These fisheries are diverse and therefore it is necessary to stratify commercial trips into fleet sectors (strata) with similar characteristics. Data from Northeast Fisheries Observer Program and the Fishing Vessel Trip Report are used together to define the size of the sample and the size of the strata, respectively. We define a total of 227 fisheries for 2005 observer coverage, consisting of three major gear types, four mesh sizes, two levels of trip durations, six port areas, and four seasonal quarters. The total fishing effort for April 2003 to March 2004 in the defined strata comprises 43,703 trips. Our examination of efficacy of observer coverage included results from 1,103 trips and 2,704 sea days. Every effort has been made to make the sampling program synoptic (i.e., cover all the major fisheries that discard commercially important species) and robust to sources of uncertainty. In particular, we utilize discard information at the trip level as opposed to the tow level. Sampling selection relies on observable properties of the strata, rather than desired outcomes (e.g., a targeted “cod” trip). Trips within strata are also assigned a probability of obtaining useful information relative to the species group of interest. The “usefulness” of a trip is conditional on the likelihood that a trip will catch one or more of the species within a predefined group of species.

Our analysis of sea-day allocations and use of optimization methods to improve allocations rest on two primary assumptions. First, the extant data are sufficient to obtain consistent estimates of the underlying variance of the discard ratio per stratum. Consistency is ensured if the samples are representative. Second, the relative size of the strata, i.e., the total number of trips, remains

constant from year to year. This is a more tenuous assumption, as the balance of fishing effort can change in response to changes in resource abundance or regulations. Both of these assumptions are inherent in the use of retrospective data to improve a future sampling program.

The observer sea-day allocation model developed here represents an extension of Neyman optimal allocation (Cochran 1977). Observer trips are allocated to strata as a function of their contribution to the total variance, the expected number of observer days per trip, and the probability that a trip will provide information on one or more of the species groups of interest. The essential features of the sampling design and allocation process are summarized below.

- Strata are defined on the basis of observable properties of the fleet sector
- The sample unit within a stratum is a trip
- The primary response variables are total discards and kept weights of groups of species. Eleven groundfish species constitute one group, monkfish another group, and summer flounder-scup-sea bass, a third group
- The probability of obtaining information on one or more of the species groups from a future trip in a stratum is estimated from analysis of observer data
- An estimate of the probability of not obtaining any information about one of the three species groups is incorporated to allow appropriate increases in sample sizes commensurate with this risk
- Expected average trip durations are defined for each stratum
- Total observer days at sea serve as a constraint on the allocation process
- Additional constraints can be imposed on the minimum and maximum numbers of samples per stratum
- Unsampled strata use imputed (or borrowed) values from adjacent strata to ensure that some information is used for sample selection
- Imputation also identifies gaps in coverage and allows for updates of the population frame as new data are acquired
- Discard ratios and standard errors incorporate the approximate covariance of the ratio
- The precision of the overall discard/kept ratio is the primary performance measure in the allocation process.
- Total variance can be minimized subject to a total observer day constraint, or the number of observer days can be minimized subject to a desired level of precision

Results from the optimization model are used as a tool to improve observer coverage. Some post-processing of the optimized sea days is needed to fine-tune coverage across fleet sectors. Where feasible, the fine-tuning of sea-day allocation capitalizes on the multi-purpose attributes of observer coverage oriented toward assessment of non-fish species (e.g., acquire data in the sea scallop fishery from trips designed to evaluate turtle bycatch rates.)

Presently the model is based on aggregate Discard/Kept (D/K) ratios. These ratios are relevant to most fisheries but, of course, the Discard/Effort (D/E) ratio is important in others. D/E ratio data have been prepared but not yet implemented in the model. D/E ratios are relevant for fisheries such as sea scallops, northern shrimp, and herring. It should be noted that one of the primary difficulties of implementing the D/E methodology is the selection of an appropriate unit of effort.

The “trip” level of effort may be the most useful but additional work will be necessary before extending the methodology to optimally allocate observer coverage to these fisheries.

The optimization methodology addresses the precision of the overall D/K ratio in the context of multiple objectives and limited resources. The issue of accuracy/bias is addressed by comparing various properties of vessels with and without observers onboard. Bias -- the systematic difference between the estimated and true value -- is addressed by first ensuring that the vessel trips are representative, and that a variety of quality assurance/control procedures are employed to accurately monitor vessel performance. Refusals to take an observer and other forms of non-response by industry are possible sources of bias. These sources are addressed via increased use of Enforcement personnel. For these concerns, the NEFSC observer program is consistent with the recommendations of the NMFS National Working Group on Bycatch (NMFS 2004).

Babcock et al. (2003) assert that increases in sampling effort are sufficient to reduce bias. If the presence of observers onboard alters the vessels fishing patterns, then it can be argued that all observed trips yield potentially biased results. If the unobserved vessel fishes with different methods in different areas and so forth, then the increases in sample size can only reduce but not eliminate the scope for bias. A variety of statistical techniques for inferring bias can be applied, but a review of the literature suggests that these techniques have been only moderately successful. Independent measures of vessel behavior may be possible from Vessel Monitoring System data, but such analyses can only detect gross changes from observed trips. Where possible, verification by independent data sources is encouraged, but one should be careful to avoid the problems of incorrectly assuming that a particular methodology is completely unbiased.

Several tests were conducted to address the potential sources of bias by comparing measures of performance for vessels with and without observers present. Bias can arise if the vessels with observers on board consistently catch more or less than other vessels, if the average trip durations change, or if vessels fish in different areas. Each of these hypotheses was tested by comparing observable properties in strata having vessels with and without observers. Average catches (pounds landed) for observed and total trips compare favorably, following an expected linear relationship. The expected difference of the stratum specific means and standard deviations for both kept weight of groundfish and total trip duration was near zero. The frequency distribution of these differences provided no evidence of systematic bias. The mean difference between average catch rates of 238 pounds was not significantly different from zero ($p=0.59$, $df=84$). A paired t-test of the stratum specific standard deviations of pounds kept suggested no significant difference from zero ($p=0.08$). A similar analysis of average trip duration revealed a strong correlation between observed and unobserved trips (Figure 7) and a suggestion that the observed trips were about a half-day longer when the observer was on board ($p = 0.01$). A paired t-test of the difference in stratum specific standard deviations of trip length was not significantly different from zero ($p = 0.60$) (Figure 8B). Some skewing of the differences in mean trip durations was observed, with observed trips being slightly longer.

Two measures of spatial coherence suggest that the spatial distribution of fishing effort for trips having observers closely matches the spatial distribution of all trips. The null hypothesis of

observer proportions equal to the VTR proportions was rejected ($P < 0.05$) in 20 of 65 comparisons. Of these 20 cases, 10 involved ports in Southern New England and the Mid-Atlantic region where landings of New England groundfish are expected to be low. Of the remaining ten cases, five involved the large and extra-large gill net fisheries that mainly target monkfish. Thus, the null hypothesis of equivalent spatial distribution of sampling was rejected in only 5 of 50 fleet sectors, a rejection rate only slightly higher than due to chance alone.

A paper by Murawski et al. (2005 in press) presents information on the spatial distribution of otter trawl fishing effort for vessels with Vessel Monitoring Systems (VMS) with the distribution of tows on observed trips. Qualitatively, the spatial distributions match very well with high concentrations of effort near the boundaries of the existing closed areas on Georges Bank and within the Gulf of Maine. Moreover, the effort concentration profiles deduced from VMS data coincided almost exactly with the profiles derived from observed trips. Overall, these comparisons suggest strong coherency between the two independent measures of fishing locations.

An assessment of the sources of uncertainty in the design and data collected in the Northeast Fisheries Observer program indicates that the level of precision in the discard ratios (d/k) for the New England Groundfish fisheries as a whole is high and there is little evidence of bias. However, at finer temporal and spatial scales, precision of the discard ratios will generally be lower than the aggregate. Precision of the discards estimates will also be lower for individual species, age groups and size classes.

Introduction

Estimation of bycatch in any commercial fishery is a difficult task. At the level of an individual trip, bycatch occurs sporadically over wide geographical ranges. Proper quantification typically requires presence of trained observers. The commercial marine fisheries of the Northeastern US comprise many vessels of widely different sizes, targeting multiple species in a variety of habitats. Overlaying the complexity of the fleet and target species is a complex regulatory environment that constrains fleet behaviors. Since many stocks are in rebuilding phases, the effects of restrictions on landings per trip, and therefore revenue per trip, are difficult to predict. The Northeast Fisheries Observer Program (NEFOP) addresses this complexity by first ensuring that the data obtained from any trip are of the highest quality. This is achieved through a rigorous training program, standardized on-board data collection protocols, and thorough auditing of data. To allow for extrapolation from the sample data to the fleet as a whole, these procedures must be embedded in a statistical sampling design. This report provides a summary of the issues relevant to the design and analysis of the observer sampling program particularly with respect to the allocation of observer days to achieve desired levels of precision.

The NEFOP program incorporates the following important features:

1. Definition of a sampling frame across all relevant fisheries
2. Identification of strata based on observable properties
3. Development of rules for imputing variance estimates in unsampled strata (i.e., “borrowing” estimates from appropriate strata)
4. Use of a trip as the sample unit (rather than individual tow)
5. Definition of discards by species groups, corresponding to the major finfish species within the Northeast US.
6. Use of discard to kept ratios (d/k) for species groups as the primary response variable.
7. Estimation of approximate variances for d/k for groups of species, rather than individual species
8. Allocation of sampling effort based on reduction in total variance of the d/k estimate, subject to total cost constraints.
9. Allowance for observer coverage in remaining fisheries not included in the sampling frame, owing to other priorities (e.g., protected species concerns).
10. Where feasible, capitalize on the multi-purpose attributes of observer coverage oriented toward assessment of non-fish species (e.g., acquire data in sea scallop fishery from trips designed to evaluate turtle bycatch rates.)

In this report we describe the foundations of our standardized approach for bycatch reporting methodologies and the primary sources of uncertainty.

Background

The Northeast Fisheries Science Center (NEFSC) routinely allocates observer coverage to monitor bycatch (fish, invertebrates, and protected species) in the commercial fisheries in the Mid-Atlantic and New England regions. The observer coverage is administered in units of ‘sea

days'. Based on the daily cost of an observer at sea, the available funds determine the number of potential sea days. However, for the New England groundfish fishery, the number of sea days is presently mandated to be 5% coverage of the fishery. The projected fishing activity (in days) for the year is estimated by the available days-at-sea allowed under the Northeast Multispecies Fishery Management Plan. Thus, in a given year, the NEFSC has a mixture of mandated sea days and non-mandated sea days to monitor bycatch in the Northeast region (North Carolina to Maine) for various fisheries.

Allocation of sea days is guided by an optimization algorithm that is based on generalization of the well-known Neyman allocation principle in survey sampling. Precision of the overall estimate of the discard ratio is improved by allocating samples to strata with the greatest contribution to the total variance, subject to an overall constraint on available resources. In this application, "resources" refers to the total number of observer days available. Improvement of the allocation process requires an evaluation of the current sampling design and precision of estimators. The ability to improve the design is contingent on the reliability of the stratum-specific variances and the persistence of these estimates in the future (or at least the next sampling period).

The optimization algorithm can be used to (1) minimize the variance of the discard estimate subject to a given number of sea days, or (2) minimize the number of sea days subject to a desired level of precision. Results from the optimization model are used as a tool to improve the coverage. However, the model does not incorporate information regarding sampling for protected species, nor does it include information for fisheries where the discard ratio may be more appropriately measured by a discard to effort ratio (d/e). Thus the model predictions are conditioned to exploit the multipurpose utility of the protected species sampling, and coverage in important fisheries (like sea scallops) is ensured by reserving some additional days to "level out" sampling that may be required for either protected species or closed area trips.

This report will describe: 1) the fishery identification and data sources used; 2) imputation rules for unobserved fisheries; 3) sampling theory and optimization methods; 4) application of the model to observer coverage; and 5) address accuracy issues discussed by Babcock et al. (2003)

Definition of Strata -- Fishery Identification

Diverse commercial fisheries are prosecuted off the Northeastern coast of the USA. These fisheries vary in size (number of trips) and have varying bycatch rates. To monitor these fisheries with at-sea observers, it is necessary to stratify the trips into fleet sectors with similar characteristics. For this report, fleet sectors are defined as strata within a survey design.

Commercial fishing trips are partitioned into fleet sectors using five classification variables: calendar quarter, gear type, mesh size, geographical region, and trip length. These classification variables are selected because they are generally known *before* a trip occurs. Using these criteria it is possible to generate a list of candidate vessels for each stratum, which simultaneously enables a random selection process and reduces the number of repeat trips on vessels. This is a

critical aspect for both strata definition and sample selection. One cannot base a sampling design on the outcome of a sample observation. In this exercise, it is not possible to select a sampling design that specifically improves the precision of cod discards, since that objective is dependent on the realization of the actual sample. However, it is possible to select samples that will improve the probability of obtaining improved discard estimates by estimating the expected proportion of trips that catch species groups of interest.

Calendar quarter was considered the most feasible temporal unit to capture seasonal variations in fishing activity and bycatch rates over the full range of fisheries. Although some management regulations operate at a finer scale (e.g. weekly), quarterly data can be further subdivided if finer resolution is needed. Otter trawl, gillnet and longline gear were defined as the three major gear types for finfish. Otter trawl and gillnet trips were classified into four mesh size groups: Small (less than 3.99 inch mesh); Medium (between 3.99 and 5.49 inch mesh); Large (between 5.5 and 7.99 inch mesh) and XLarge (8.0 inch mesh or greater). Additionally, trips are classified into six geographical regions based upon the port of departure: ports located within Maine and New Hampshire (ME_NH); Massachusetts (N_MA, excluding Bristol county); Connecticut, RI, and Bristol county, MA (SNE); New Jersey - New York (NJ/NY); Maryland and Delaware (MD/DE); Virginia and North Carolina (VA/NC). Trip length serves as a surrogate for spatial resolution (inshore vs. offshore). Otter trawl trips are further classified into two trip length categories: day trips and multi-day trips. Longline and gillnet gears are not partitioned by trip length.

Due to the mixture of species caught during a trip, it is not sufficient to classify trips with regard to target species because discard of target and non-target species may occur. To account for target and non-target discard, trips in each fleet sector are classified into one or more of three species groups: New England groundfish (NEGF); summer flounder, scup and black sea bass (FSB); and monkfish (MONK). There is often overlap between trips which catch NEGF, FSB and MONK. The estimated number of trips and sea days needed to cover these fleet sectors may be overestimated when the trips are assumed to be independent, therefore the overlapping nature of the fishing fleets are taken into account. Sampling fractions, and how the overlap is accounted for, are described in a later section.

Eleven species constitute the New England groundfish species group: cod, haddock, yellowtail flounder, American plaice, witch flounder, winter flounder, redfish, pollock, white hake, windowpane, and halibut. If a trip catches (retains or discards) at least 1 of the 11 large-mesh regulated species, the trip is categorized as NEGF trip and the haul weights of the 11 species are summed to form an aggregate species total for NEGF. Similarly, if a trip catches (retains or discards) either summer flounder, black sea bass or scup, the trip is categorized as a FSB trip and the haul weights of these species are summed to form an aggregate species total for FSB. If a trip catches (retains or discards) monkfish, then the trip is categorized as a MONK trip. A trip may be categorized to one or more of the three species groups.

Data Sources

Trip characteristics are recorded in both the NEFOP and Fishing Vessel Trip Reports (VTR) data sets. Together, these databases are used to define the size of the sample and the size of the strata, respectively. Data from each source are retrieved and prepared separately before the two sets are combined (Figure 1).

Fishing Vessel Trip Report Data

Beginning in June 1994, the Northeast Region's data collection system was changed from a voluntary to a mandatory reporting system for USA fishermen and dealers who catch and buy/sell groundfish species regulated by the Northeast Multi-species Fishery Management Plan. The mandatory reporting system consists of two components: 1) dealer reporting and 2) vessel trip reporting. Each component contains information needed for fishery management and stock assessment analyses: the dealer reports contain total landings by market category, while the vessel trip reports contain information on area fished, kept and discarded portions of the catch, and fishing effort. The VTR data has been routinely used in management analyses and peer reviewed stock assessments. Details on example applications of the VTR to stock assessments may be found in a large number of reports of the Stock Assessment Review Committee (SARC). Reports prepared since 2000 may be found at <http://www.nefsc.noaa.gov/nefsc/saw/>. Earlier reports are available by contacting saw_reports@noaa.gov.

In this report, the VTR data are used to: 1) define the sampling frame of the commercial fishing trips, and 2) evaluate the accuracy of the observer data with respect to area fished, kept pounds, and trip length. The VTR data are the only synoptic data source for vessel activity, area fished and fishing effort for commercial fisheries. The Vessel Monitoring System data and the Days-At-Sea data systems cover only portions of the fisheries and therefore are limited in use.

The VTR data can be used as a basis for defining the sampling frame, because all federally permitted vessels are required to file a VTR for each fishing trip (see NMFS-NERO http://www.nero.noaa.gov/ro/fso/vtr_inst.pdf). These self-reported data constitute the basis of the fishing activity of the commercial fleets. The VTR trip data are collapsed into fleet sectors and species groups as defined above. For each species group within a fleet sector, the number of trips that caught the species group, the average number of days absent, and the weight of the species in the species group are calculated.

The limitations of self-reported catch data are well known (e.g., Walsh et al. 2002, NMFS 2004). Limitations of the initial data VTR data sets were described by the SARC in 1996 (NMFS 1996). Since then, many of these limitations have been addressed. In particular, subsequent peer-reviews through numerous SARCs and a review by the National Research Council (1998) have identified the strengths, weaknesses, and appropriate uses of the VTR data from the Northeast.

The validity of VTR data as a basis for a sampling frame is supported by comparisons with total landings data from dealer records. All dealers which buy and sell groundfish regulated by federal

FMPs are required to report 100% of the landings. These data are generally thought to constitute a near census of landings of groundfish. The NRC (1998) noted that misreporting of landings is “usually a significant issue only when fisheries are managed by setting a total allowable catch.” On this basis, the magnitude of misreporting by dealers would be low as Northeast groundfish stocks have been managed primarily through effort controls. A comparison of total groundfish landings from VTR and Dealer records for calendar year 2003 reveals close agreement between the two sources:

Species	VTR Landings (mt)	Dealer Landings (mt)	Difference (mt)	Percent Difference
Cod	8240	8692	452	5.2%
Winter flounder	5321	5714	393	6.9%
Witch flounder	2971	3108	137	4.4%
Yellowtail flounder	5208	5530	322	5.8%
American Plaice	2204	2415	211	8.7%
Windowpane flounder	102	60	-42	-70%
Haddock	5778	5874	96	1.6%
White Hake	2268	3305	1037	31.4%
Halibut	11	13	2	15.4%
Redfish	338	360	22	6.1%
Pollock	3839	4188	349	8.3%
Total	36281	39258	2977	7.6%

For the three major species, cod, haddock and yellowtail flounder, the percentage differences range from 1.6% to 5.8%. Only windowpane flounder, white hake and halibut exhibit large percentage differences. Total landings of windowpane flounder and halibut represent small fractions of the total (0.3% of VTR and 0.2% Dealer) landings and these percentage differences are considered negligible. Large percentage differences for white hake may be attributable to confusion between white hake and red hake. White hake can be difficult to distinguish from red hake (sp) and may be identified simply as “hake” by both dealers and fishermen. The overall difference of 7.6% is dominated by large differences in the landings of white hake. Excluding white hake from the comparison reduces the overall percentage difference to 5.4%.

Other measures to ensure the validity of the VTR database include routine auditing procedures, standardized data entry protocols and compliance reviews (pers. comm. Greg Power, Chief, Fisheries Information Section, Northeast Regional Office, NMFS).

Northeast Fisheries Observer Program Data

The NEFOP employs trained, sea-going observers to collect catch data by species and disposition (retained and discarded). Biological samples, gear characteristics data, and economic information are also collected. For the optimization data set, only observed hauls from trips classified as ‘standard sea sampling trips’ are used. Observed trips that were aborted or which

used a ‘limited’ fish sampling protocol (no discard data collected) are excluded. Hail weight can be reported in round or dressed weights; if kept hail weights are reported as ‘dressed’, then the hail weight is converted to round (live) weight using Commercial Fisheries Database System (CFDBS) conversion factors for the species. All discard hail weights are assumed to be round (live) weight.

The NEFOP data are collapsed into strata as defined above. For each stratum, the number of observed trips that caught one or more of the three species groups is calculated. For each fleet sector and species group, the number of observed trips, number of observed hauls, average trip length (in days), kept weight of all species in the species group, discarded weight of all species in species group, and the number of observed days are calculated. A discard ratio and the variance of the ratio are calculated for each stratum (fleet sector and species group).

Optimization Data Set

The VTR and NEFOP data sets are concatenated by fleet sector and species group. A list of variables and their definitions are presented in Table 1. Not all VTR fleet activity may have NEFOP coverage (Table 2). When fleet sectors do not have observer coverage, imputed values are used (Table 3). The imputed values are derived from NEFOP data from similar fleet sectors, thus providing an estimate for the non-observed fleets. Details of the imputation process are provided in the following section.

The optimization tool is flexible and allows the user to select the entire input data set, or a subset. To allocate sea days for an entire year, four calendar quarters of data are used. Using the most recent available data, given the time needed for data entry and auditing, the year consists of calendar quarter 3 and 4 from year -1 and calendar quarter 1 and 2 from the current year.

The three gear types (otter trawl, gillnet, and longline) used in the optimization data set are gear types for which fishing regulations allow finfish to be retained, thus a discard to kept ratio estimator (d/k) is used. Fisheries using other gear types where regulations may prohibit groundfish possession are excluded from the current optimization process because a d/k ratio is not appropriate for these cases.

Imputation rules for unobserved fisheries

Not all of the fishery strata had observed trips between April 2003 and March 2004. To account for the expected variance of the estimates in the missing cells, it was necessary to develop a standardized procedure to handle both missing and minimal levels (e.g., a single trip) of observer coverage. This procedure is referred to hereafter as ‘imputation’ and the estimates derived by the imputation are referred to ‘imputed values’. Imputed values are derived by sequentially relaxing the fleet sector classification. The fleet sectors for each species group (NEGF, FSB, and MONK) are imputed separately. The imputed values fill in missing values for the unobserved strata. Fishery strata are defined with respect to rigid definitions of categorical variables such as region

or quarter. A stratum with missing data must be filled with data from similar strata. To identify suitable candidate strata as “donor” or “parent” cells, it is necessary to “relax” the definitions of the strata. For example, if no trips occur in the Jan.-Mar. quarter, one might relax the definition to include data from the Jan-Jun. half year. The objective process of relaxing strata definitions to impute data is described below.

A fleet sector was not imputed if:

- 1) VTR number of trips = 0 (no imputation needed when there is no fleet activity for the species group);
- 2) VTR number of trips > 0 and standard error was not missing (no imputation needed when there is fleet activity for the species group and there is a standard error of the observer d/k ratio); and
- 3) VTR number of trips > 0 and total observed kept pounds = 0 (no imputation needed when there is fleet activity for the species group and the standard error cannot be calculated); otherwise, the fleet sector was imputed.

The imputation uses three increasing levels of aggregated NEFOP data (using the same data and calculation methods as the original calculations of observed d/k ratio and associated statistics). Three of the five stratification factors are relaxed (region, mesh size and calendar quarter). Gear type and trip length are used, but their stratification is not relaxed. Trip length is not relaxed because the average trip length is used to determine the number of sea days needed to obtain the desired precision level. Gear type is not relaxed because of fundamental differences in catches (retained and discarded) occur using these gear types.

Level 1: Calendar quarter is relaxed to half year and the six geographic regions are relaxed to two regions (NE region = ME/NH, N_MA, SNE; MA region = NY/NJ, DE/MD, NC/VA); gear, mesh size and trip length categories are maintained.

Level 2: Calendar quarter is relaxed to an entire year, the six geographic regions are relaxed to two regions (as in Level 1), and the four mesh groups are relaxed to two mesh groups (SMALL = small and medium mesh groups; LARGE = none, large, and Xlarge mesh groups); gear and trip length categories are maintained.

Level 3: Calendar quarter is relaxed to an entire year (as in Level 2), the six regions are relaxed to one region (all six regions combined), and the four mesh groups are relaxed into one mesh group. This level served as a ‘catch-all’ for all remaining fleets sectors that required imputation.

The VTR-NEFOP data set is merged with Level 1 NEFOP data; if a fleet sector needs imputed values, based on the criteria list above, then the imputed values from the observed trips in Level 1 are transferred to the corresponding VTR-NEFOP fleet sector and species group only if the trips in the Level 1 data set are greater than 1. Data from Level 2 and Level 3 are subsequently merged with the VTR-NEFOP. When imputed values are used in the VTR-NEFOP data set,

the fleet sector and species group is ‘flagged’ with the imputation level used. All fleet sectors that need imputation obtain values at one of the three levels.

Below is a summary of the number of fleet sectors, by imputation level and species group used in the 2005 sea day allocation.

Imputation Level	Species group		
	NEGF	FSB	MONK
Level 0 (no imputation)	150	116	111
Level 1	30	51	44
Level 2	27	41	35
Level 3	20	19	37
Total	227	227	227

To include all fisheries using otter trawl, gillnet and longline gear in the optimization, approximately 33% to 50% of the mean discard rates and variances are imputed or ‘borrowed’.

When a fleet sector and species group is imputed, five variables (number of observed trips, observed d/k ratio, total observed kept pounds, standard error of the d/k ratio, and number of observed days) are estimated with imputed values. Because the aggregated NEFOP data at each level have more observations than the original VTR-NEFOP fleet sector, the imputed values need to be rescaled before they are used. Except for the imputed d/k ratio, the imputed values for the number of observed trips, the total observed kept pounds, the standard error and the number of observed days are re-scaled using a sampling fraction represented by the ratio of the total NEFOP trips for that level, fleet sector and species group to the total VTR trips for that level, fleet sector and species group. Equations used to re-scale imputed values within stratum h are:

$$\begin{aligned}
 T_{vtr} &= \text{total VTR trips of Level}_i \\
 T_{obs} &= \text{total NEFOP trips for Level}_i \\
 T_{imp,h} &= (T_{obs} / T_{vtr}) * \text{Trips}_{vtr,h}; \\
 \text{Kept}_{imp} &= (T_{imp,h} / T_{obs}) * \text{NEFOP kept pounds sum in Level}_i \\
 \text{SE}_{imp} &= (T_{obs} / T_{imp,h})^{1/2} * \text{NEFOP standard error in Level}_i \\
 \text{Days}_{imp} &= (T_{imp,h} / T_{obs}) * \text{total number of NEFOP days in Level}_i \\
 T_{imp,h} &\text{ is rounded to a whole number, if } T_{imp,h} < 1, \text{ then } T_{imp,h} = 1;
 \end{aligned}$$

where Level_i denotes Imputation Level 1, Level 2 or Level 3.

Sampling Theory and Optimization Methods

Fishing trips are considered the primary sample unit in estimating d/k ratios. Fishing trips generally catch multiple species, some of which are not landed owing to various regulations or market conditions. We defined three major groups of species: (1) New England groundfish, (2) summer flounder, scup and sea bass, and (3) monkfish. Fishing trips in a given stratum may catch species from one or more of these groups. The degree of overlap among species groups has important implications for the efficacy of sampling within strata, i.e., the number of samples necessary to achieve a desired level of precision. Because some fraction of trips provide information on more than one species group, estimates of sample size based on the assumption of independence, will overestimate the number of required trips. Developing estimators that explicitly account for the magnitude of overlap can circumvent this potential inefficiency. There are two ways to approach this estimation. One is based on the pattern of overall trips from the vessel trip reports. The second is based on the pattern in observer sampled trips. In theory, if the observed trips are a representative sample, the proportions in the vessel trip reports and observer trips should be the same. In practice, the proportions in the observed trips will deviate from those in the VTRs due to sampling variability and other factors. The selection of observed trips reflects a practical mix of vessel availability, knowledge of vessel operations, familiarity, and safety considerations. These are, of course, important factors for program management, but it must be recognized that these factors introduce bias into estimates.

Both approaches follow the algorithm described below. Let I_{hij} be an indicator variable denoting the presence or absence of species group j within trip i in stratum h . Then $I_{hij}=1$ if species group j is present, else 0. A design matrix can be used to describe each unique trip within a stratum. The design matrix appends to each trip record a set of indicator variables that identify the presence/absence of species groups caught. The following table illustrates a hypothetical case with 7 trips in stratum h .

Example 1

	$I_{h,1}$ j=1	$I_{h,2}$ j=2	$I_{h,3}$ j=3
<u>Trip ID</u>	<u>NEGF</u>	<u>Monk</u>	<u>FSB</u>
1	1	0	0
2	1	1	0
3	1	1	1
4	1	0	1
5	0	1	1
6	0	1	0
7	0	0	1
Sum	4	4	4
$n_h=7$	n_{h1}	n_{h2}	n_{h3}

In this simple example, four of the seven trips caught New England groundfish, four trips caught monkfish, and four caught summer flounder, scup or sea bass. If all of these trips (or trip types) are equally likely, then the probability of obtaining a sample that yields information on NEGF is $4/7$ and so forth. The probability of obtaining information on species j is the sum of the species

group specific trips within the stratum (i.e., n_{hj}) divided by the total number of unique trips within the stratum (n_h). Note that

$$n_h \neq \sum_{j=1}^3 n_{hj}$$

owing to the overlap in coverage for some trips. The probability that a random trip provides information on species group j is defined as

$$\hat{p}_{hj} = \frac{n_{hj}}{n_h} \quad (1)$$

For each stratum, the probabilities can be computed that a random sample will contain information about species group j . The basis for the probability estimator can either be the observed set of trips within a stratum or the total set of trips represented in the VTRs. Applying the same set of indicator variables to the VTR data, one can obtain the population estimates of these quantities as

$$\hat{P}_{hj} = \frac{N_{hj}}{N_h} \quad (2)$$

Eq. 1 establishes the basis for a random sample from the set of observed trips. Eq. 2 establishes the same basis from the VTR. On first principles, Eq. 2 is a better estimator if a representative sample can be taken in a stratum. Eq. 1 is more appropriate if the set of observed trips within a stratum is representative of those trips available for observation.

Using Eq. 1 or 2, it is now possible to examine the effects of altered sample sizes. Let n'_h represent the new total number of trips to be taken in stratum h . For the purpose of evaluating the expected change in variance in the component species groups, the n'_{hj} for each species group need to be redefined. This is accomplished using the equation

$$n'_{hj} = \hat{p}_{hj} n'_h \quad (3)$$

if Eq. 1 is used, or

$$n'_{hj} = \hat{P}_{hj} n'_h \quad (4)$$

if Eq. 2 (based on VTR) is used to estimate the expected probabilities that a trip in stratum h will capture fish from species group j .

Another worked example will reinforce the basic concept of the expected proportions of samples likely to sample species group j . Consider a stratum with 10 observed trips with Eq.1 used to estimate p'_{hj} .

Example 2

<u>Trip ID</u>	<u>I_{h,1}</u> j=1 NEGF	<u>I_{h,2}</u> j=2 Monk	<u>I_{h,3}</u> j=3 FSB
1	1	1	0
2	1	0	0
3	1	0	1
4	1	1	0
5	1	1	1
6	0	0	1
7	0	0	1
8	1	0	1
9	0	1	0
10	0	1	0
Sum	7	4	5
n _h =10	n _{h1}	n _{h2}	n _{h3}
p _{hj}	7/10	4/10	5/10

If the n_h were increased to n'_h=30 then the revised estimates of n'_{hj} would be

$$\hat{n}'_{h1} = \left(\frac{7}{10}\right)30 = 21$$

$$\hat{n}'_{h2} = \left(\frac{4}{10}\right)30 = 12$$

$$\hat{n}'_{h3} = \left(\frac{5}{10}\right)30 = 15$$

Thus, adding 20 trips to stratum h would translate into an expected increase of 14 trips for NEGF (i.e., 21-7), 8 trips for monkfish (i.e., 12-4) and 10 trips for FSB (i.e., 15-5). The increase in the total number of trips for a stratum differs with respect to the pattern of information in the sample. The allowance for non-integer numbers of trips is considered to have a negligible effect. In practice, the actual implementation of a sampling strategy would be based on rounding to the nearest integer, and subject to a lower bound constraint, say n_{hj}= 2.

Example 2 could be repeated for estimates derived from the VTR data. For such an example, the universe of trips would be much larger.

Measures of Overlap

Venn diagrams of the number of trips in the VTR and NEFOP depict the degree of overlap between the three species groups in the two data sets. In the April 2003-March 2004 VTR

database, half of the trips (22,274 trips out of 43,703 trips) are unique to the species groups (Figure 2), while in the NEFOP database, a third of the trips (286 trips out of 1,103 trips) are unique to the species groups (Figure 3). The sampling fractions (NEFOP trips divided by VTR trips) are given in Figure 4. The numbers of trips (and days) in the Venn diagrams are based on whole trips, and therefore slight differences occur in the number of trips between the Venn diagram and d/k ratio analyses (e.g. there are trips in d/k ratio analysis which used two different mesh sizes during a trip).

Observers Days at Sea Constraints

While trips constitute the sampling unit, the total number of sampling units is constrained by the total number of days available during any interval. To consider this component of the sampling design, it is necessary to consider the average trip duration in stratum h . Let t_{hi} be the trip duration (days) for the i -th trip in stratum h . The total number of observed trips in stratum h is n_h and the total number of observed days is $\sum t_{hi}$. The average trip duration is estimated as

$$\bar{t}_h = \frac{\sum_{i=1}^{n_h} t_{hi}}{n_h} \quad (5)$$

The actual number of future observer days that will be required under some new sampling intensity (n'_h) is proportional to n'_h/n_h . Eq. 5 can also be defined in terms of the durations of the trips in the VTR database. The expected total number of days allocated to stratum h is defined as

$$T_h = \bar{t}_h n_h = \sum_{i=1}^{n_h} t_{hi} \quad (6)$$

regardless of whether observer or VTR data are used. The average trip duration in stratum h is not influenced by the number of trips allocated, as long as the trips selected are representative of the basis used to define the species composition of the trips. Recall that either the observer database or the VTR database can be used. Thus the total number of observer days allocated to stratum h under some new allocation is

$$T'_h = \bar{t}_h n'_h \quad (7)$$

The grand total number of days at sea that would be allocated given some new set $\{n'_h\}$ would be

$$T' = \sum_{h=1}^H \bar{t}_h n'_h \quad (8)$$

Some key points in this derivation are:

- It is not possible to derive any real-world sampling program without considering the key uncertainties related to the probability that the trip will be “successful” and that the cost of sea days may vary.
- The number of successful trips, relative to the objective of reducing the variance of the estimate, is a random variable, based on a probability estimate. The expected number of actual trips may not actually result in information necessary to improve the precision of the estimate.
- The “cost” per trip is expressed as the expected duration. Actual duration may also vary within strata, although the stratification is designed reduce the variation in this component.

Optimization is a technique for maximizing (or minimizing) some quantity of interest subject to one or more constraints. Constraints are the key concept. In this application, we consider upper and lower bounds on the size of the sample within a strata, a total constraint on the number of available days, and a constraints related to acceptable levels of precision. For problems that do not explicitly consider dynamic (i.e., time dependent) processes, a variety of optimization methods can be used including linear and nonlinear programming. For this project, the optimization program, Premium Solver Platform (Version 5.5) developed by Frontline Systems, Inc. (2003) was used.

To address the optimization problem, the overall variance of the discard to kept ratio must first be estimated. The discard ratio for species group j in stratum h is the sum of discard weight over all trips divided by sum of kept weights over all trips:

$$\hat{R}_{jh} = \frac{\sum_{i=1}^{n_h} d_{ijh}}{\sum_{i=1}^{n_h} k_{ijh}} \quad (9)$$

where d_{ijh} is the discards for species group j within trip i in stratum h and k_{ijh} is the kept portion of the catch. R_{jh} is the discard rate for species group j in stratum h . The stratum weighted discard to kept ratio for species group j is obtained by weighted sum of discard ratios over all strata:

$$\hat{R}_j = \sum_{h=1}^H \left(\frac{N_h}{\sum_{h=1}^H N_h} \right) \hat{R}_{jh} I_h \quad (10)$$

The variable I_h is a zero/one indicator of whether or not a stratum is included in the computation. The indicator variable can be considered as a composite measure of the suitability of stratum h in the estimator. The indicator variable allows a stratum to be filtered on the basis of one or more metrics. A more complete description of the various types of filtering is described in the next section.

The approximate variance of the estimate of R_{jh} is obtained from a first order Taylor series expansion about the mean:

$$V(\hat{R}_{jh}) = \frac{1}{(n_{jh} - 1)n_{jh}\bar{k}_{jh}^2} \left[\left(\sum_{i=1}^{n_{jh}} d_{ijh} \right)^2 + \hat{R}_{jh}^2 \left(\sum_{i=1}^{n_{jh}} k_{ijh} \right)^2 - 2\hat{R}_{jh} \left(\sum_{i=1}^{n_{jh}} d_{ijh} \right) \left(\sum_{i=1}^{n_{jh}} k_{ijh} \right) \right] \quad (11)$$

where d_{ijh} is the total discard weight of species group j in trip i within stratum h , k_{ijh} is the total kept weight of species group j in trip i within stratum h , n_{jh} is the sample size (number of trips) that caught species group j in stratum h , and \bar{k}_{jh} is the mean kept landing of species group j within stratum h . Note that in this formulation of the variance, the finite population correction factor (fpc), i.e., one minus the sampling fraction within the stratum, has been omitted. This has been done to improve readability. The fpc is included however, in Eq. 11 for the total variance of the d/k ratio.

The variance of the d/k ratio for species group j over the entire set of strata is estimated using standard sampling theory methodology for a stratified random design as

$$V(\hat{R}_j) = \sum_{h=1}^H \left(\frac{N_h - n_{jh}}{N_h} \right) \left(\frac{N_h}{\sum_{h=1}^H N_h} \right)^2 V(\hat{R}_{jh}) I_h \quad (12)$$

The overall coefficient of variation for the discard/kept ratio is defined as

$$CV_j = \frac{\sqrt{V(\hat{R}_j)}}{\hat{R}_j} \quad (13)$$

It is now possible to define an overall estimate of the relative precision of the d/k ratio across all species groups as

$$CV = \sum_{j=1}^3 \lambda_j CV_j \quad (14)$$

where λ_j is an arbitrary weighting factor for species group j . In this formulation, the λ_j can be used as binary factors (0,1) to examine the allocations individually for species groups.

The optimization tool evaluates the potential improvements in the precision of the discard ratio through reallocation of the number of trips to individual strata. Equation 11 illustrates that the variance of the ratio decreases as the number of trips (n_h) increases. Assuming that the data yield representative estimates of the stratum specific variances, then the reduction in total variance can be examined as a function of alternative allocation schemes for each stratum. If n_h^* is defined as the optimal number of trips taken in stratum h , then the variance of the overall ratio is estimated as

$$V(\hat{R}_j^*) = \sum_{h=1}^H \left(\frac{N_h - n_{jh}}{N_h} \right) \left(\frac{N_h}{\sum_{h=1}^H N_h} \right)^2 \left(\frac{n_{jh}}{n_{jh}^*} \right) V(\hat{R}_{jh}) I_h \quad (15)$$

The optimization problem can now be posed as the minimization of the CV of the composite ratio estimate, subject to a total days at sea constraint (T_C) and constraints on the number of trips per stratum.

$$\begin{aligned} & \min \sum_{j=1}^3 \lambda_j CV_j \\ & \text{subject to} \\ & 2 \leq n_{jh}^* \leq N_h \quad , \forall_h \quad (16) \\ & T_C^* \geq \sum_{h=1}^H \bar{t}_h n_h^* \end{aligned}$$

Alternatively, the optimization problem can be defined with the objective of minimizing the total number of days at sea, subject to an acceptable coefficient of variation (CV_{CRIT}). This version of the model can be written as:

$$\begin{aligned}
& \min \sum_{h=1}^H \bar{t}_h n_h^* \\
& \text{subject to} \\
& 2 \leq n_{jh}^* \leq N_h \quad , \forall_h \\
& CV_{CRIT} \geq \sum_{j=1}^3 \lambda_j CV_j
\end{aligned} \tag{17}$$

Another relevant consideration is that a trip may not yield information on any of the target species groups. In some strata, for example, a number of trips fail to capture groundfish, monkfish or the summer flounder, scup and sea bass mixture. To protect against this possibility, it is desirable to inflate the optimal number of trip estimates by the ratio of N_h to N'_h where N_h is the total number of trips in stratum h and N'_h is the number of trips that obtained information on one or more of the species groups.

Application of the Model

Using the optimization algorithm to minimize the variance of the discard estimates subject to a given number of sea days, the allocation of observer sea days for the Mid-Atlantic (M-A) and New England (NE) regions was optimized separately and the resulting allocated sea days combined. Separate analyses were conducted because of differential sea days constraints (mandated sea days for New England groundfish versus non-mandated sea days for the Mid-Atlantic region). Before the optimization began, a portion of the available sea days were set aside to cover fisheries which do not enter the optimization process (e.g. scallop dredge fishery). For these fisheries, sea days are allocated proportional to fishing effort (number of trips or number of days fished).

The Mid-Atlantic optimization used data from the SNE, NJ/NY, DE/MD and VA/NC regions with the species weighting coefficients set to 1 for both FSB and MONK and to 0 for NEGF. The NE optimization used data from the SNE, N_MA, and ME-NH regions, with the species weighting coefficients set to 1 for NEGF and to 0 for both FSB and MONK. Data from the SNE region were included in both optimizations due to the intersection of the NE and M-A regions. Stratum indexes were applied to reduce the data set to contain only the relevant fisheries.

Below is a summary of the indexes and thresholds used in the NE and M-A sea day optimizations.

NE region trip and landings setting and thresholds

<i>Switch</i>	<i>Setting</i>	<i>Threshold (fraction)</i>	<i>Description of Filters that Operate on Entire Strata</i>
I(L_negf%)	1	0.0025	Landings of NEGF<Threshold=>0, else 1
I(L_fsb%)	(All)	0.0001	Landings of FSB<Threshold=>0, else 1
I(L_monk%)	(All)	0.0001	Landings of Monk<Threshold=>0, else 1
sum(I(L_all%))	(All)	NA	If any of Landings indices for NEGF,FSB or Monk=1 then =>1, else 0
I(Nh_negf%)	1	0.0001	Trips of NEGF<Threshold=>0, else 1
I(Nh_fsb%)	(All)	0.0001	Trips of FSB<Threshold=>0, else 1
I(Nh_monk%)	(All)	0.0001	Trips of Monk<Threshold=>0, else 1
I(%TotVTR_3sp)	1	0.00005	Filter on % of total landings of 3 species groups
Filter on All Trips	0	NA	Excludes entire Strata if value=0

M-A region trip and landings settings and thresholds

<i>Switch</i>	<i>Setting</i>	<i>Threshold (fraction)</i>	<i>Description of Filters that Operate on Entire Strata</i>
I(L_negf%)	(All)	0.0025	Landings of NEGF<Threshold=>0, else 1
I(L_fsb%)	1	0.0001	Landings of FSB<Threshold=>0, else 1
I(L_monk%)	1	0.0001	Landings of Monk<Threshold=>0, else 1
sum(I(L_all%))	(All)	NA	If any of Landings indices for NEGF,FSB or Monk=1 then =>1, else 0
I(Nh_negf%)	(All)	0.0001	Trips of NEGF<Threshold=>0, else 1
I(Nh_fsb%)	1	0.0001	Trips of FSB<Threshold=>0, else 1
I(Nh_monk%)	1	0.0001	Trips of Monk<Threshold=>0, else 1
I(%TotVTR_3sp)	1	0.00005	Filter on % of total landings of 3 species groups
Filter on All Trips	0	NA	Excludes entire Strata if value=0

NE and M-A regions d/k ratio thresholds

	<i>Threshold (d/k ratio)</i>	<i>Description of Filters that Operate on Individual Cells (Species within Strata)</i>	<i>Number of Cells Included</i>	<i>Number of Cells Excluded</i>
Max d/k_NEGF	1	Maximum d/k ratio used for NEGF. Values>Threshold excluded	25	11
Max d/k_FSB	2	Maximum d/k ratio used for FSB. Values>Threshold excluded	32	4
Max d/k_Monk	2	Maximum d/k ratio used for Monkfish. Values>Threshold excluded	33	3

Some ‘post-processing’ of the allocation of optimized sea days was necessary. Even though one or more indicator variables (i.e., filters) were applied during optimization, it was necessary to fine-tune the sea day allocations by applying a minimum and maximum amount of coverage, and to maintain coverage of fishing activity throughout the year. The optimized sea days were multiplied by the average trip duration for each stratum to estimate the projected number of observed trips. If the projected number of observed trips was less than 3 trips per strata, then the sea days were redistributed to other strata representing more relevant fisheries. If the number of

potential observed trips in a stratum exceeded 15% of the VTR trips, then the sea days in that stratum were reduced to the number of sea days representing 15% (potential observer trips/VTR trips) coverage. The sea days from strata exceeding the 15% coverage cap were reassigned to other strata.

The number of unique vessels and the vessel selection protocols in a stratum limit the number of trips that can be observed in that stratum. The number of unique vessels varies among strata; in the 2005 sea day optimization, the number of unique vessels in a stratum ranged between 1 and 146 vessels, with 85% of the strata having 50 vessels or less. The vessel selection protocols state a vessel is not to be observed more than twice during a month. As an approximate guide for balancing between the potential number of observed trips and the number of unique vessels in a stratum, a 15% trip coverage cap was selected to prevent assigning more sea days to a stratum than the number of vessels could support. The 15% cap prevented clustering of sampling effort, particularly in instances where the estimate of the variance of d/k might be imprecise. In these instances, the optimization model will tend to allocate large number of trips to such strata to reduce the standard error of the estimate. When the analysis was restricted to the relevant strata for the New England groundfish fisheries, the 15% cap was binding in only 4 of 33 strata for the observer coverage allocation scheme based on 2,708 observer days.

The diagnostics within the optimization tool were used to evaluate the imputation process. The optimization algorithm calculates the d/k ratios and the variance estimates for 'all data' and for 'data without imputed values'. Generally, the d/k ratios and variance estimates were similar between the 'all data' and 'data without imputed values' for each species groups. This indicates that the imputation generally provided consistent values across the three levels of aggregation.

Precision, Bias and Sampling Intensity: A Rebuttal to E.A Babcock et al. (2003)

Understanding the sampling properties of estimates of bycatch derived from observer programs and other sources with respect to accuracy and bias is critical. This section reviews issues related to bycatch estimation in observer programs with an emphasis on potential biases that may exist. The NMFS national bycatch report (NMFS 2004) emphasizes that wherever possible, attempts to detect and guard against bias should be made in observer programs. The report strongly advocates the development of rigorous randomization procedures in sample selection to help ensure representative sampling. All can agree that with unlimited resources, the more observer coverage the better. The real issue however is how to allocate finite resources to meet multiple requirements for stock assessment and protected species evaluation. The cases that Babcock et al. (2003) point to as success stories typically have relative few boats involved compared to many other fisheries. These cases are not representative overall of the issues facing program managers.

Babcock et al. (2003) insufficiently distinguish between two very different types of bias. The first type arises when non-representative sampling occurs. The second type is related to the statistical properties of the consistency of the estimators. These two types of bias are very different and it is important to be clear which type of bias is under consideration. The second type of bias is typically reduced with sufficiently large sample size. However, this may not be

addressed by increases in sample size if fishermen refuse to take observers, if certain classes of boats cannot accommodate observers, etc. Babcock et al. (2003) take as an article of faith that increasing the number of trips will reduce bias. Some of the solutions identified by Babcock et al. (2003) for correcting bias (e.g. the use of bootstrap estimators) apply to correcting bias of the second type. However, no amount of bootstrapping will overcome non-representative sampling.

The mean square error (MSE) of an estimate is composed of two elements, the variance of the estimate and the square of the bias (defined as the difference between the mean of the sample and the true population value). The MSE therefore comprises two additive elements. Cochran (1977) notes that if bias is less than 10% of the standard deviation of the estimate, the effect of this bias on the accuracy of the estimate is negligible. As noted by Babcock et al. (2003), most work on the properties of estimates derived from observer programs have focused on the variance component, with far fewer studies examining bias. For reasons described in detail below, we believe that estimating the bias of the first type is more difficult than intimated by Babcock et al. (2003). It is nonetheless important to try to estimate this quantity. Focusing on the precision part of the MSE in certain analyses does not imply that bias is unimportant, or that it should be dismissed as insolvable as suggested by Babcock et al. (2003)

A critical element of the arguments developed by Babcock et al. (2003) appears to be that increasing the number of trips sampled will, by itself, reduce bias of the first type. This assertion, if true, is important. However, no corroborative evidence is provided. The argument is that fishermen will change behavior if they are subjected to a higher probability of being included in a sample, or of being sampled more frequently by observers. In essence, fishermen will be less likely to fish in a non-typical manner when an observer is on board if the probability of selection is higher. This may not be true if say a particular fishing trip has a 20% chance of being selected vs. a 10% chance and if the fishermen do not know in advance how many trips they may have to accommodate within a specified time period. In any event, we doubt that this can be calculated unless a model of human behavior is part of the estimation procedure.

Babcock et al. (2003) report that Sampson (2002) detected statistically significant differences between a multivariate indicator of landings composition by participants in the Enhanced Data Collection Project (EDCP) of the Oregon Department of Fish and Wildlife and the composition of landings by the entire groundfish trawl fleet. This analysis is used to indicate that biases exist in voluntary programs such as the EDCP and that it is possible to use similar approaches to identify bias in observer programs in general. What Babcock et al. do not report is that Sampson indicated that the multivariate analysis employed (Principal Components Analysis) was only “moderately successful” in capturing the properties of the data. The first three principal components accounted for 15.4, 12.0, and 8.0 % of the variance respectively for trips landing more than 10,000 lbs in which hake comprised less than 50% of the total (designated “Big” trips by Sampson). For trips less than 10,000 lbs in which hake comprised less than 50% of the total (“Small” trips), the first three principal components accounted for 13.7, 10.4, and 9.0% of the variance. Sampson (2002) reported significant differences between the participants in the EDCP and the total fleet in the 1st and 3rd principal components for both Big and Small trips and concluded that the EDCP fleet may not be representative of the entire fleet. However, because the first three PCs captured only a moderate fraction of the variance, these analyses should be viewed with caution. It is worth noting that Sampson provided canonical variable plots of PCA 1

against PCA 2 (Figure 6a and 6b of his report) in which both the information from the EDCP and the whole fleet are superimposed and these show that the data from the EDCP do not appear to be markedly different from the total fleet. A truly important bias should show up clearly in these plots, which take into account more of the variance of the samples than the individual t-tests actually used in the report.

The general issue of testing for bias in observer data using landings data raises some important questions concerning the inferences that can be drawn. In particular, if no significant differences are detected between observer and landings data, this does not guarantee that there is no bias in the estimates of discards.

The other major source of information that could be used to test the representativeness of observer data is to test against self-reported estimates by fishermen. Sampson (2002) made such an analysis for the EDCP data and detected differences. In this case, it was inferred that the self-reported estimates were not accurate. In contrast, Liggins (1997) found no differences between observer data for catch and discards against fleet wide estimates. In general, self-reported estimates are rightly viewed with caution and this is the most commonly available type of discard information against which to compare observer data.

To deal with logistical constraints and their effect on observer programs, Babcock et al. (2003) cite the work of Cotter et al. (2002) using a probability proportional to size (PPS) sampling allocation procedure. However, Cotter et al. (2002) concluded that this approach did not markedly improve the performance of the estimators.

Babcock et al. (2003) refer to the method of collapsing strata as an *ad hoc* procedure when, in fact, it is a very well established method (see Cochran 1977). Bias can occur using this method if an investigator deliberately chooses similar strata to combine. However, methods in which objective rules for combining strata are employed are much less likely to cause bias.

Babcock et al. (2003) assert that Fogarty and Gabriel (2002) assumed that the sampling fraction did not matter. In fact, Fogarty and Gabriel (2002) noted that the sampling fraction does affect the precision of the estimate through the finite population correction factor. The effect indicated by Babcock et al. (2003) is a very well established property of the statistical estimators employed. Fogarty and Gabriel (2002) noted in their analysis that “Ignoring the finite population correction factor results in an overestimate of the standard error...” Fogarty and Gabriel (2002) did not include the FPC in their estimates so as to provide a conservative estimate of the variance (e.g. biased on the high side). This is very different than assuming that the sampling fraction does not matter.

Recommendations made by the NMFS National Working Group on Bycatch (NMFS 2004) largely address the issues of major concern – the importance of obtaining representative sampling, careful consideration of stratification, etc. We recommend that information from observer trips (catch, trip duration, number of hauls/tows, fishing location etc.) also be checked against independent sources of information to see if differences can be detected. The only solution that Babcock et al. (2003) provide when such a bias is detected is to increase the number of trips covered by observers. As noted above, this may or may not be effective. Other solutions

to the problem need to be explored, as well as increasing observer coverage when analyses indicate it is cost-effective to do so given finite resources and competing programmatic needs.

An Evaluation of Bias in the Northeast Fisheries Observer (Sea Sampling) Program

Several tests were conducted to address the potential sources of bias. We compared several measures of performance for vessels with and without observers present. Bias can arise if the observed trips within a stratum are not representative of the other vessels within the stratum. Such bias could arise if the vessels with observers on board consistently catch more or less than other vessels, if the average trip durations change, or if vessels fish in different areas. Each of these hypotheses was tested by comparing observable properties in strata having data from vessels with and without observers.

All vessels are required to report the total trip landings, the number of days absent from port, and the primary statistical area fished. Average catches (pounds landed) for observed and total trips compare favorably (Figure 5), and follow an expected linear relationship. If the observed and unobserved trips within a stratum measure the same underlying process, one would expect no statistical difference in the average catches (and the standard deviations) between the VTR and observer data sets. An examination of the distribution of these differences (Figures 6A and 6B) indicates no evidence of systematic bias. The mean difference of 238 pounds in average catch rates between the two data sets is not significantly different from zero ($p=0.59$, $df=84$). As well, a paired t-test of the stratum specific standard deviations of pounds kept showed no significant difference from zero ($p=0.08$). A strong correlation was detected in trip duration between observed and unobserved trips (Figure 7), with observed trips averaging about a half-day longer ($p = 0.01$) (Figure 8A). However, the difference in stratum specific standard deviations of trip length was not significantly different from zero ($p = 0.60$) (Figure 8B). Some skewing of the differences in mean trip durations is evident, with observed trips being slightly longer.

Two measures of spatial coherence were also examined. Within stratum h the expected number of observer trips by statistical area j as the product of the proportion of VTR trips in Statistical Area j and stratum h (V_{jh}) and the number of observed trips in stratum n_h . Thus, $E_{jh} = V_{jh} * n_h$. These expectations can then be compared to the actual frequencies (O_{jh}) of observed trips by statistical area. Results of these analyses indicate that the spatial distribution of fishing effort for trips with observers on board closely matches the spatial distribution of trips for the stratum as a whole (Table 4). It was possible to compute chi-square statistics for 65 strata. The null hypothesis of observer proportions equal to VTR proportions was rejected ($P<0.05$) in 20 of the 65 comparisons. Of these 20 cases, 11 were from ports in Southern New England and Mid-Atlantic states. Of the remaining nine cases, five involved the large and extra-large gill net fisheries that land both groundfish and monkfish. Thus, the null hypothesis of equivalent spatial distribution of sampling was rejected in only 4 of 50 cases, a rejection rate only slightly higher than expected from chance alone.

As a final measure of the potential spatial bias, a paper by Murawski et al. (2005 in press) is instructive. In this paper, information is presented on the spatial distribution of otter trawl fishing effort for vessels with Vessel Monitoring Systems (VMS) and compared with the

distribution of fishing effort from observed trips (Figure 9). Qualitatively, the spatial distributions match very well with high concentrations of effort near the boundaries of existing closed areas on Georges Bank and within the Gulf of Maine. Moreover, the effort concentration profiles deduced from VMS data coincide almost exactly with the profiles derived from the observed trips. Overall, these comparisons suggest strong coherency between these two independent measures of fishing locations.

Sources of Uncertainty

In the Northeast, every effort is made to ensure representative observer coverage. This is accomplished by stratifying the fleet into homogeneous spatial, temporal and gear groups and by randomly selecting vessels from these strata. Stratification and randomization of sampling units are basic principles of survey design (e. g. Cochran 1977; Thompson 2002) and have been used in previous studies of bycatch to improve both “knowledge of the fleet” (Cotter et al. 2002) and precision of estimates (Allen et al. 2002; Borges et al. 2004). VTR data are used to produce a list of fishing vessels, by quarter and fleet sector. The vessel list contains a randomly ordered list of all vessels that participated in each fleet sector. To obtain a representative sample of the fleet, the NEFOP Area Coordinators use this vessel list, in addition to their local knowledge of fleet activity, to identify vessels on which to place observers. Vessels are required to take an observer if requested to do so. The NEFOP has standard protocols regarding vessel selection. A vessel, using the same gear, is not observed more than twice in the same month— this prevents repeated observations from the same vessel. The NEFOP Area Coordinators have protocols for documenting refusals; a refusal occurs when a vessel owner/captain is asked to take an observer and the owner/captain declines — or agrees but does not follow through (i.e. the vessel leaves the dock without the observer on board). Refusals are forwarded to Law Enforcement. A vessel owner can be prosecuted for failing to take an observer.

An objective process is used for imputation of missing values in unsampled strata. The imputation methodology helps identify gaps in sampling strategy and is an important component for ongoing improvements of the survey design. Stratoudakis et al. (1999) employed a post-stratification technique of “collapsing strata” as a way of dealing with unsampled strata. Our method of imputing means and variances for unsampled strata builds on this approach by utilizing information in comparable strata as a basis for initial sample allocation. Imputation represents a tradeoff between a realistic survey consistent with known fishing patterns and a less realistic pooled survey. Excessive imputation, however, can be indicative of an overly ambitious stratification approach; utilizing the observer data at an unrealistically fine temporal or spatial scale (say daily estimates in a small area) not only leads to an excessive extrapolation, but also violates the premise that observations in the current year are sufficient to predict patterns in the following year.

Persistence of annual patterns is critical to the estimation of an ‘optimal’ scheme. As regulations change and fishing patterns shift, using data based on fleet activity in the preceding year may be problematic. Using the current year’s fishing activity pattern to predict future fishing patterns within strata cannot account for changes induced by variations in resource abundance, revenues, or management regimens. In a study of discards in the North Sea, Statoudakis et al. (1998)

reported immediate increases in discarding rates following increases in minimum size limits, but noted consistent patterns over time and among gears for higher value species such as cod and haddock. Without a predictive model of human behavior, it is not possible to anticipate fine-scale changes in fishing patterns. Rochet et al. (2002) were unable to find reliable predictor variables for prediction of bycatch but it should be noted that their study examined only 26 trips, about two orders of magnitude less than the number of trips considered in this report.

A related source of uncertainty is the ability to make inferences about specific species, stocks or age groups. Our evaluation of the Northeast Observer Program considers discard to kept ratios at the level of species groups. This approach is consistent with recent literature (Allen et al. 2001, Borges et al. 2004). An optimal strategy for New England Groundfish as a group however, will not necessarily be optimal for age 2 haddock on Georges Bank. The precision of discard information required at this level will typically exceed the nominal levels predicted as a result of optimal sampling. Figure 10 illustrates the relationship between the coefficient of variation for the overall New England groundfish discard ratio estimate as a function of total observer days allotted to this fishery. Assuming that 2,708 sea days can be allocated in an optimal manner in 2005, the predicted CV of the d/k ratio is well below 4%. The predicted CV drops to 2.5% at about 4,000 days and drops to about 1% at 20,000 days (about 50% coverage). The continuously decreasing slope of the relationship between CV and observer sea days reflects the reduced effectiveness of additional days as a way of improving overall precision.

Several important points are relevant to the interpretation of Figure 10. First, any non-optimal allocation of sampling effort will tend to increase the overall CV of the d/k ratio. Non-optimal allocations occur when the desired sampling plan cannot be followed, or when the pattern of landings among the strata in the current year differs from the pattern used as a basis for the optimal allocation scheme. Second, the CV of the overall d/k ratio is smaller than the precision of the individual components. Thus, the CV of the d/k ratio for a particular gear type or for a d/k ratio based on a finer temporal or spatial scale will generally be greater than the composite estimate. This property is illustrated in Figures 11 and 12 for quarterly estimates in the New England groundfish otter trawl and gillnet fisheries, respectively. Note that the number of observed otter trawl trips would need to be tripled to reduce the CV of the d/k ratio from 20% to 10%.

The coefficient of variation (CV) of the d/k ratios for New England groundfish are well below the 20% - 30% CV range established by the Atlantic Coastal Cooperative Statistics Program (ACCSP) for high priority commercial fisheries (ACCSP 2001) and by NMFS's National Working Group on Bycatch (NWGB) (NMFS 2004). The NWGB recommends: "For fishery resources, excluding protected species, caught as bycatch in a fishery, the recommended precision goal is a 20-30% CV for estimates of total discards (aggregated over all species) for the fishery; or if total catch cannot be divided into discards and retained catch then the recommended goal for estimates of total catch is a CV of 20-30% (NMFS 2004). Assuming that landings are known without error, the precision of estimated total discard for New England groundfish equals the precision of the d/k ratio for this fishery.

A decrease in precision of the d/k ratio is also expected for any single species analysis. For example, the CV of the d/k ratio for haddock alone will probably be much greater than the CV of

the d/k ratio for the overall groundfish complex. Once again, it is important to remember that the sampling program must be based on observable properties of the strata, not on the outcome of the experiment. Any efforts to improve the precision of the d/k ratio for a single species will come at the expense of reduced precision for other species. Moreover, oversampling of a particular group of vessels may introduce undesirable properties (e.g., repeat trips on a single vessel) that can make the sampling less representative.

An exact definition of an acceptable level of bias and precision depends on the objectives of the analyses and the levels of acceptable risk to the fishery resource and the fishery. The acceptable level of risk must be defined externally by managers but should, at a minimum, consider the risk of stock collapse if management actions are compromised by imprecise information on discards. From the analyses presented in this report, it would appear that the level of precision is high for the groundfish resource as a whole and that there little evidence of bias in the discard rates.

Presently the optimization model uses aggregate d/k ratios, which are appropriate for most fisheries; however, for other fisheries, d/e ratios are more appropriate. The optimization algorithm can handle datasets containing either type of ratio, but not both in the same set (without external weighting). Input data sets with d/e ratios have been developed, but have not yet been incorporated into the overall process. A comparison of the precision of alternative estimators of discard ratios is the subject of ongoing research.

Acknowledgments

We wish to thank Mark Terceiro, Katherine Sosebee, and Ralph Mayo for their insights and assistance in identifying the fishery strata, the bases for imputation, and the iterative process of refining the application. We also thank Fred Serchuk for his constructive comments and review.

References

- ACCSP (Atlantic Coastal Cooperative Statistics Program). 2001. Technical Source Document Series V: Biological Module and Discard, Release and Protected Species Interactions Module. June 28, 2001 draft. 137 p. On-line document: <http://www.accsp.org/tsdocs.htm> .
- Allen, M., D. Kilpatrick, M. Armstrong, R. Briggs, N. Perez, and G. Course. 2001. Evaluation of sampling methods to quantify discarded fish using data collected during discards project EC 95/-94 by Northern Ireland, England, and Spain. *Fish. Res.* 49:241-254.
- Allen, M., D. Kilpatrick, M. Armstrong, R. Briggs, G. Course, and N. Perez. 2002. Multistage cluster sampling design and optimal sampling sizes for estimation of fish discards from commercial trawlers. *Fish. Res.* 55:11-24.
- Babcock, E.A., E. K. Pikitch and C.G. Hudson. 2003. How much observer coverage is enough to adequately estimate bycatch? Report of the Pew Institute for Ocean Science, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL. On-line version: <http://www.oceana.org/uploads/BabcockPikitchGray2003FinalReport.pdf>
- Borges, L., A. F. Zuur, E. Rogan, and R. Officer. 2004. Optimum sampling levels in discard sampling programs. *Can. J. Fish. Aquat. Sci.* 61:1918-1928.
- Cochran, W.L. 1977. *Sampling Techniques*. J. Wiley and Sons. New York.
- Cotter, A.J.R., G. Course, S.T. Buckland and C. Garrod. 2002. A PPS sample survey of English fishing vessels to estimate discarding and retention of North Sea cod, haddock and whiting. *Fisheries Research* 55: 25-35.
- Fogarty, M.J. and W. Gabriel. 2002. Relative precision of discard estimates for the Northeast groundfish complex. Report of National Marine Fisheries Services, Northeast Fisheries Science Center, Woods Hole, MA.
- Frontline Systems. 2003. Premium Solver Platform version 5.5. Incline Village, NV. 222 p.
- Liggins, G.W., M.J. Bradley, S.J. Kennel. 1997. Detection of bias in observer-based estimates of retained and discarded catches from a multispecies trawl fishery. *Fisheries Research Report* 9(3):46-52. University of British Columbia.
- Murawski, S., S. Wigley, M. Fogarty, P. Rago and D. Mountain. (article in press). Effort distribution and catch patterns adjacent to temperate MPAs. *ICES Journal of Marine Science*.
- NMFS (National Marine Fisheries Service). 2004. Evaluating bycatch: a national approach to standardized bycatch monitoring programs. U. S. Dep. Comm., NOAA Tech. Memo. NMFS-F/SPO-66, 108 p. On-line version, http://www.nmfs.noaa.gov/by_catch/SPO_final_rev_12204.pdf

NMFS-NERO (National Marine Fisheries Service) Northeast Regional Office.
http://www.nero.noaa.gov/ro/fso/vtr_inst.pdf

National Research Council (NRC) 1998. Review of Northeast Fishery Stock Assessments. National Academy Press. Washington DC

NEFSC (Northeast Fisheries Science Center). 1996. Analysis of the 1994 fishing vessel logbook data. In: 22nd Northeast Regional Stock Assessment Workshop: Stock Assessment Review Committee consensus summary of assessments. NEFSC Reference Doc. 96-13; 242p.

Rochet, M-J, I. Peronnet, and V. M. Trenkel. 2002. An analysis of discards from the French trawler fleet in the Celtic Sea. ICES J. Mar. Sci. 59:538-552.

Sampson, D. 2002. Final Report to the Oregon Trawl Commission on Analysis of Data from the At-Sea Data Collection Project. Oregon State University. Newport, Oregon. On-line
<http://www.onid.orst.edu/~sampsond/projects/edcp>

Stratoudakis, Y., R. J. Fryer, R. M. Cook. 1998. Discarding practices for commercial gadoids in the North Sea. Can. J. Fish. Aquat. Sci. 55:1632-1644.

Stratoudakis, Y., R. J. Fryer, R. M. Cook, and G. J. Pierce. 1999. Fish discarded from Scottish demersal vessels: Estimators of total discards and annual estimates for targeted gadoids. ICES J. Mar. Sci. 56:592-605.

Thompson, S. K. 2002. Sampling. 2nd ed., J. Wiley and Sons, Inc. New York.

Walsh, W. A., P. Kleiber, and M. McCracken. 2002. Comparison of logbook reports of incidental blue shark catch rates by Hawaii-based longline vessels to fishery observer data by application of a generalized additive model. Fisheries Research 58:79-94.

Table 1. The variables, their description, their associated species group, data source, and units of the input data set of the optimization algorithm.

<i>Variable Name</i>	<i>Definition</i>	<i>Species Group</i>	<i>Data Source</i>	<i>Units</i>
year	Year			categories
negear	gear type			categories
qtr	quarter of year			number
mesh	mesh size			categories
region	state grouping, port of departure			categories
trp	Trip Duration (days)			categories
alltrips	Total number of trips, all species	ALL	VTR	trip
allmnda	Ave number of days absent, all species	ALL	VTR	days
vcount	Total number of VTR trips for 3 sp. Groups	3 Sp Grp	VTR	trip
ocount	Total number of observed trips that caught one or more of the 3 sp groups	3 Sp Grp	VTR	trip
vnegfntrips	Number of VTR trips that caught NEGF	NEGF	VTR	trip
vgfda	Total VTR days absent for trips that caught Groundfish	NEGF	VTR	days
vgftotal	Total VTR pounds(all sp) landed for trips landing groundfish	NEGF	VTR	pounds
vgflb	VTR pounds landed—groundfish	NEGF	VTR	pounds
vgfmnda	VTR average days absent—groundfish	NEGF	VTR	days
onegf	Sum of the "0/1 flags" for observed trips that caught NEGF	NEGF	OBS	trip
ogfntrips	Number of observed trips that caught NEGF	NEGF	OBS	trip
ogfparent	Flag indicating if values of d/k are observed (=1) or imputed (=0)	NEGF	OBS	flag
ogfnewcv	Desired CV closest to 0.30--intermediate value	NEGF	OBS	number
ogfnewntrips	Number of Observed trips necessary to achieve CV=ogfxnewcv	NEGF	OBS	trip
ogfxnewcv	Desired CV=0.30 --exact value	NEGF	OBS	number
ogfavgtriplen	Ave Trip Length in days for observed trips	NEGF	OBS	days
ogfntows	Number of observed Tows	NEGF	OBS	tows
ogfksums	Kept—observed	NEGF	OBS	pounds
ogfdsums	Discarded—observed	NEGF	OBS	pounds
ogfdkratio	d/k ratio	NEGF	OBS	number
ogfse	SE of d/k ratio	NEGF	OBS	number
ogfcv	CV of mean d/k ratio	NEGF	OBS	number
ogfseadays	Number of sea days needed to achieve CV=0.3 (=avg triplen x newntrips)	NEGF	OBS	days
ogfndays	Number of observed days	NEGF	OBS	days
vfsbntrips	Number of VTR Trips that caught FSB	FSB	VTR	trip
vfsbda	Total VTR days absent for trips that caught FSB	FSB	VTR	days
vfsbtotal	Total VTR pounds (all sp) landed for trips landing FSB	FSB	VTR	pounds
vfsblb	VTR pounds landed—FSB	FSB	VTR	pounds
vfsbmnda	VTR average days absent—FSB	FSB	VTR	days
ofsb	Sum of the "0/1 flags" for observed trips that caught FSB	FSB	OBS	trip
ofsbntrips	Number of observed trips that caught FSB	FSB	OBS	trip
ofsbparent	Flag indicating if values of d/k are observed (=1) or imputed (=0)	FSB	OBS	flag
ofsbnewcv	Desired CV closest to 0.30--intermediate value	FSB	OBS	number
ofsbnewntrips	Number of Observed trips necessary to achieve CV=ofsbxnewcv	FSB	OBS	trip
ofsbxnewcv	Desired CV=0.30 --exact value	FSB	OBS	number

ofsbavgtriplen	Ave Trip Length in days for observed trips	FSB	OBS	days
ofsbntows	Number of observed Tows	FSB	OBS	Tows
ofsbksums	Kept—observed	FSB	OBS	pounds
ofsbdsms	Discarded—observed	FSB	OBS	pounds
ofsbdkratio	d/k ratio	FSB	OBS	number
ofsbse	SE of d/k ratio	FSB	OBS	number
ofsbcbv	CV of mean d/k ratio	FSB	OBS	number
ofsbseadays	Number of sea days needed to achieve CV=0.3 (=avg triplen x newntrips)	FSB	OBS	days
ofsbndays	Number of observed days	FSB	OBS	days
vmonkntrips	Number of VTR Trips that caught Monk	Monk	VTR	trip
vmonkda	Total VTR days absent for trips that caught monk	Monk	VTR	days
vmonktotal	Total VTR pounds (all sp) landed for trips landing Monkfish	Monk	VTR	pounds
vmonklb	VTR pounds landed---Monk	Monk	VTR	pounds
vmonkmnda	VTR average days absent—Monk	Monk	VTR	days
omonk	Sum of the "0/1 flags" for observed trips that caught Monkfish	Monk	OBS	trip
omkntrips	Number of observed trips that caught Monk	Monk	OBS	trip
omkparent	Flag indicating if values of d/k are observed (=1) or imputed (=0)	Monk	OBS	flag
omknewcv	Desired CV closest to 0.30--intermediate value	Monk	OBS	number
omknewntrips	Number of Observed trips necessary to achieve CV=omknewcv	Monk	OBS	trip
omkxnewcv	Desired CV=0.30 --exact value	Monk	OBS	number
omkavgtriplen	Ave Trip Length in days for observed trips	Monk	OBS	days
omkntows	Number of observed Tows	Monk	OBS	Tows
omkksums	Kept—observed	Monk	OBS	pounds
omkdsums	Discarded—observed	Monk	OBS	pounds
omkdkratio	d/k ratio	Monk	OBS	number
omkse	SE of d/k ratio	Monk	OBS	number
omkcbv	CV of mean d/k ratio	Monk	OBS	number
omkseadays	Number of sea days needed to achieve CV=0.3 (=avg triplen x newntrips)	Monk	OBS	days
omkndays	Number of observed days	Monk	OBS	days
onegfcpue	Observer Catch(kept) per unit effort (lbs/day) for NEGF	NEGF	OBS	lbs/day
ofsbcpue	Observer Catch (kept) per unit effort (lbs/day) for FSB	FSB	OBS	lbs/day
omkcpue	Observer Catch (kept) per unit effort (lbs/day) for Monk	Monk	OBS	lbs/day
alltotal	Total number of pounds of all species landed in this cell	ALL	VTR	pounds
vnegfcpue	VTR Landings per unit effort (lbs/day) for NEGF	NEGF	VTR	lbs/day
vfsbcpue	VTR Landings per unit effort (lbs/day) for FSB	FSB	VTR	lbs/day
vmkcpue	VTR Landings per unit effort (lbs/day) for Monk	Monk	VTR	lbs/day
L_negf%	Fraction of NEGF landings in stratum h	NEGF	VTR	unitless
L_fsb%	Fraction of FSB landings in stratum h	FSB	VTR	unitless
L_monk%	Fraction of Monk landings in stratum h	Monk	VTR	unitless
Nh_negh%	Fraction of NEGF trips in stratum h	NEGF	VTR	unitless
Nh_fsb%	Fraction of FSB trips in stratum h	FSB	VTR	unitless
Nh_monk%	Fraction of Monk trips in stratum h	Monk	VTR	unitless
I(L_negf%)	Indicator {0,1} for Fraction of NEGF landings in stratum h	NEGF	VTR	switch
I(L_fsb%)	Indicator {0,1} for Fraction of FSB landings in stratum h	FSB	VTR	switch
I(L_monk%)	Indicator {0,1} for Fraction of Monk landings in stratum h	Monk	VTR	switch
sum(I(L_all%))	Indicator {0,1} for composite landings. =0 if all species specific indicators=0,else 1	3 Sp Grp	VTR	switch
I(Nh_negf%)	Indicator {0,1} for Fraction of NEGF trips in stratum h	NEGF	VTR	switch
I(Nh_fsb%)	Indicator {0,1} for Fraction of FSB trips in stratum h	FSB	VTR	switch

<i>I(Nh_monk%)</i>	Indicator {0,1} for Fraction of Monk trips in stratum h	Monk	VTR	switch
<i>sum(I(Nh_all%))</i>	Indicator {0,1} for composite TRIPS. =0 if all species specific indicators=0,else 1	3 Sp Grp	VTR	switch
<i>I(onegfcpue)</i>	Indicator {0,1} for observer CPUE in stratum h for NEGF. 1=> exceeds threshold, else 0	NEGF	OBS	switch
<i>I(ofsbcpue)</i>	Indicator {0,1} for observer CPUE in stratum h for FSB. 1=> exceeds threshold, else 0	FSB	OBS	switch
<i>I(omkcpue)</i>	Indicator {0,1} for observer CPUE in stratum h for Monk. 1=> exceeds threshold, else 0	Monk	OBS	switch
<i>I(vnegfcpue)</i>	Indicator {0,1} for VTR CPUE in stratum h for NEGF. 1=> exceeds threshold, else 0	NEGF	VTR	switch
<i>I(vfsbcpue)</i>	Indicator {0,1} for VTR CPUE in stratum h for FSB. 1=> exceeds threshold, else 0	FSB	VTR	switch
<i>I(vmkcpue)</i>	Indicator {0,1} for VTR CPUE in stratum h for Monk. 1=> exceeds threshold, else 0	Monk	VTR	switch
<i>I(d/k_negf)</i>	Indicator {0,1} for Obsvr d/k ratio in stratum h for NEGF. 1=> exceeds threshold,else 0	NEGF	OBS	switch
<i>I(d/k_fsb)</i>	Indicator {0,1} for Obsvr d/k in stratum h for FSB. 1=> exceeds threshold, else 0	FSB	OBS	switch
<i>I(d/k_monk)</i>	Indicator {0,1} for Obsvr d/k in stratum h for Monk. 1=> exceeds threshold, else 0	Monk	OBS	switch
<i>Total VTR 3spgroup</i>	Sum of landings by strata for each species group	3 Sp Grp	VTR	switch
<i>%Total VTR 3 group</i>	Percent of landings of sum of 3 sp groups in strata	3 Sp Grp	VTR	switch
<i>I(%TotVTR_3sp)</i>	flag for total landings of 3 species groups	3 Sp Grp	VTR	switch
<i>ogfimp_level</i>	Indicator {0,1,2,3} of imputation level	NEGF	OBS	category
<i>ofsbimp_level</i>	Indicator {0,1,2,3} of imputation level	FSB	OBS	category
<i>omonkimp_level</i>	Indicator {0,1,2,3} of imputation level	Monk	OBS	category

Table 2. Number of trips, by strata, in the Fishing Vessel Trip Reports (VTR) and Northeast Fisheries Observer Program (NEFOP) data sets used in the 2005 sea day optimization.

Region	Gear	Mesh	Trip length	QUARTER							
				1		2		3		4	
				VTR	NEFOP	VTR	NEFOP	VTR	NEFOP	VTR	NEFOP
DE/MD	Otter Trawl	Large	day multi-day	17	0	95	0	188	0	52	0
		Medium	day multi-day	8	2	5	0			1	0
		Small	day multi-day	3	0	14	0	3	0	24	0
	Gillnet	Medium		1	0	1	0				
		Small		4	0	1	0	1	0		
		XLarge		12	0	19	0	2	0	8	0
ME_NH	Longline	None		20	0	68	0	6	0	5	0
	Otter Trawl	Large	day multi-day	187	0	102	2	512	6	568	1
		Medium	day multi-day	315	9	279	5	479	9	439	15
		Small	day multi-day			1	0			1	0
		XLarge	day multi-day			3	0	1	0	10	0
	Gillnet	Large		1	0						
		Medium		75	0	242	0	823	10	375	3
		None				1	0	10	0	1	0
		Small						3	0		
		XLarge		19	0	77	0	573	14	247	0
N_MA	Longline	None		407	6	28	1	186	0	243	0
	Otter Trawl	Large	day multi-day	789	20	739	21	2015	54	1232	34
		Medium	day multi-day	501	7	382	13	551	10	613	9
		Small	day multi-day			11	1	1	0		
		XLarge	day multi-day			2	4	3	0	2	1
	Gillnet	Large		13	0	119	2	3	1	15	2
		Medium		12	2	57	2	3	3	15	2
		Small				1	0				
		XLarge						2	0	1	0
		None		1061	81	367	83	1481	94	1024	64
Small			1	0					2	0	
NC/VA	Otter Trawl	Large	day multi-day	2	0	5	0			3	0
		Medium	day multi-day	542	17	117	0			226	3
		Small	day multi-day	4	0	3	0			15	2
		XLarge	day multi-day	35	7	20	0				
	Gillnet	Large		12	4	4	0	2	0	13	0
		Medium		4	0	4	0				
		Small		9	0	46	0	11	0	43	0
		XLarge		19	0	5	0			10	0
NJ/NY	Otter Trawl	Large	day multi-day	2	0	8	0	4	1	15	0
		Medium	day multi-day	38	0	161	0			35	0
		Small	day multi-day	45	0	5	0				
		XLarge	day multi-day	426	4	1878	6	936	0	847	0
	Gillnet	Large		342	4	421	3	580	0	199	1
		Medium		13	1	267	21	464	5	458	4
		Small		170	22	42	5	4	1	64	3
		XLarge		29	0	629	5	894	0	465	0
SNE	Otter Trawl	Large	day multi-day	209	8	99	3	105	1	150	5
		Medium	day multi-day	7	0	4	0	31	0	20	0
		Small	day multi-day			2	0	1	0	2	0
		XLarge	day multi-day			72	0	70	0	29	0
	Gillnet	Large				49	0	81	0	31	0
		Medium				2	0			4	0
		Small		2	0	8	0	49	0	51	0
		XLarge		418	0	699	1	166	0	995	0
SNE	Otter Trawl	Large	day multi-day	273	2	996	20	1399	2	731	2
		Medium	day multi-day	571	37	515	8	621	21	525	25
		Small	day multi-day	25	1	19	3	41	1	158	2
		XLarge	day multi-day	11	0	104	6	304	2	333	10
	Gillnet	Large		503	12	269	8	188	5	373	7
		Medium		3	0	1	0	4	0	11	0
		Small		21	1	124	9	170	3	66	2
		XLarge						1	0		

Table 3. Summary of fleet sectors (strata), by species group, that are imputed (1) and not imputed (0); blank cells indicate no fleet activity.

Region	Gear	Mesh	Trip length	QUARTER														
				1			2			3			4					
				NEGF	FSB	MONK	NEGF	FSB	MONK	NEGF	FSB	MONK	NEGF	FSB	MONK			
DE/MD	Otter Trawl	Large	day				0	1	1	0	1	1	0	1	1	0	1	1
			multi-day	0	1	1	0	1	1	0	1	1	0	1	1	0	1	1
		Medium	day													0	1	0
	multi-day	0	0	1	0	1	1								0	1	0	
	Small	day				0	1	1	0	1	0	1	0	0	0	1	1	
	multi-day	0	1	1	0	1	0								0	1	1	
Gillnet	Medium		0	1	0	0	1	0										
	Small		0	1	0	0	1	0	0	0	0	1						
	XLarge		0	0	1	0	1	1	0	0	1	0	0	1	0	0	1	
ME_NH	Longline	None		1	0	0	1	0	0	1	0	0	1	0	1	0	0	
	Otter Trawl	Large	day	1	0	1	0	1	0	0	1	0	1	0	1	1	1	
			multi-day	0	0	0	0	0	0	0	0	1	0	0	0	1	0	
		Medium	day													0	1	0
		multi-day				1	0	1										
	Small	day									1	0	0	1	0	1		
	multi-day											1	0	1	0	1		
	XLarge	day				1	0	1	1	0	1	0	1	0	0	1		
	multi-day	0	0	1														
	Gillnet	Large		1	0	1	1	1	1	0	1	0	0	1	1	0	1	1
Medium															1	0	1	
None						1	0	1	1	0	1	1	1	0	1	0	1	
Small										1	0	1						
XLarge			1	0	1	1	1	1	0	0	0	1	1	1	1	1		
N_MA	Longline	None		0	0	0	1	0	0	1	0	0	1	0	1	0	0	
	Otter Trawl	Large	day	0	1	0	0	0	0	0	0	1	0	0	0	1	0	
			multi-day	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0
		Medium	day				1	1	1	1	0	1	0	1	1	0	1	
		multi-day				0	0	0	0	1	0	1	1	1	0	1		
	Small	day	1	0	1	0	0	0	0	1	0	0	0	0	0	0		
	multi-day	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0		
	XLarge	day				0	1	0										
	multi-day										1	0	1	1	0	1		
	Gillnet	Large		0	1	0	0	0	0	0	0	0	0	0	0	1	0	
Medium			1	0	0										1	0	1	
None			1	0	1	0	0	1	1	0	1	1	1	0	0	0		
Small			1	0	0	1	0	1	1	0	1	1	0	1	0	1		
XLarge			0	0	0	0	0	0	0	0	0	0	0	0	0	0		
NC/VA	Otter Trawl	Large	day	0	1	0	0	1	1						1	1	1	
			multi-day	0	0	0	0	1	1							1	0	0
		Medium	day	0	1	0	0	1	0									
		multi-day	0	0	0	0	1	1							0	0	1	
	Small	day	0	0	0	0	1	1	0	1	0	1	0	0	1	0		
	multi-day	0	1	1	0	1	1											
Gillnet	Large		0	1	1	0	1	1	0	1	0	0	0	1	1	1		
	Medium		0	1	1	0	1	1						0	1	1		
	Small		0	0	1	0	1	1	0	0	0	0	0	0	1	1		
XLarge		0	1	1	0	1	1						0	1	1			
NJ/NY	Longline	None		1	0	0	1	0	0									
	Otter Trawl	Large	day	0	0	0	0	0	0	1	1	1	1	1	1	1	1	
			multi-day	0	0	0	1	0	0	1	1	1	0	1	1	1		
		Medium	day	1	1	0	0	0	0	0	0	0	0	0	0	0	0	
		multi-day	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	
		Small	day	1	1	1	0	0	0	1	1	1	1	1	1	1	1	
	multi-day	0	0	0	0	0	0	0	1	0	0	1	0	0	1	0		
	XLarge	day				1	1	1	0	1	1	1	0	1	1	1		
	multi-day	0	1	1	0	0	1	0	1	0	1	0	0	1	0	1		
	Gillnet	Large					1	1	1	0	1	1	0	1	0	1	1	
Medium						1	1	1	1	1	1	1	1	1	1	0		
None						0	1	1						0	1	1		
Small			0	0	1	1	1	0	1	1	0	1	0	1	1	1		
XLarge			0	1	1	1	1	1	1	1	1	1	1	1	1	1		
SNE	Otter Trawl	Large	day	0	0	0	0	0	0	1	0	1	0	0	0	0	0	
			multi-day	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		Medium	day				0	0	1	0	1	1	0	0	0	0	0	
		multi-day	0	1	1	1	1	1	0	1	0	1	0	1	1	1		
		Small	day	1	1	1	0	0	0	0	0	0	1	0	0	0	0	
	multi-day	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0		
	XLarge	day				0	1	1							0	1	1	
	multi-day	1	1	1	0	1	1	1	0	1	0	1	0	1	1	1		
	Gillnet	Large		1	0	1	0	0	0	1	0	1	1	1	0	1	1	
		Medium									0	1	0					
None			1	0	1	0	0	1						0	1	0		
Small						0	1	1										
XLarge			0	0	0	0	0	0	0	0	0	0	0	0	0	0		

Table 4. Summary of contingency table analyses of spatial distribution of VTR and observed trips. Expected value of observed trips is based on proportions of VTR trips by Statistical Area. Critical value of Chi-Square statistics is based on alpha level of 0.05. Degrees of freedom are based on number of Statistical Areas reported in VTR database.

Quarter	Gear	Mesh	Region	Trip Duration	Chi Sqr Test Statistic	df	Chi Sqr Crit Value	Signif Level
3	Gill Net	Large	ME_NH	all	41.92	6	12.59	0.000
3	Gill Net	XLarge	ME_NH	all	32.19	4	9.49	0.000
3	Gill Net	Large	N_MA	all	36.92	11	19.68	0.000
3	Gill Net	XLarge	NJ/NY	all	20.30	5	11.07	0.001
4	Gill Net	XLarge	N_MA	all	16.89	4	9.49	0.002
4	Gill Net	Large	ME_NH	all	14.76	4	9.49	0.005
4	Gill Net	XLarge	NJ/NY	all	10.46	2	5.99	0.005
2	Gill Net	XLarge	ME_NH	all	12.06	7	14.07	0.098
2	Gill Net	Large	NC/VA	all	3.06	2	5.99	0.216
1	Gill Net	XLarge	NC/VA	all	2.15	2	5.99	0.341
1	Gill Net	Large	SNE	all	0.40	1	3.84	0.527
4	Gill Net	Large	N_MA	all	2.69	4	9.49	0.611
2	Gill Net	Large	N_MA	all	6.10	8	15.51	0.636
2	Gill Net	XLarge	N_MA	all	1.48	3	7.81	0.687
1	Gill Net	XLarge	N_MA	all	1.23	3	7.81	0.746
3	Gill Net	XLarge	N_MA	all	2.29	5	11.07	0.808
1	Gill Net	Large	N_MA	all	1.29	4	9.49	0.862
2	Longline	None	ME_NH	all	1.15	3	7.81	0.764
1	Longline	None	N_MA	all	1.63	7	14.07	0.977
2	Trawl	Large	N_MA	1day	243.29	6	12.59	0.000
2	Trawl	Medium	SNE	2+day	120.00	3	7.81	0.000
3	Trawl	Large	NJ/NY	1day	80.97	13	22.36	0.000
2	Trawl	Large	NJ/NY	1day	61.00	5	11.07	0.000
4	Trawl	Large	ME_NH	2+day	49.91	9	16.92	0.000
1	Trawl	Small	NJ/NY	1day	32.36	3	7.81	0.000
4	Trawl	Medium	NJ/NY	2+day	28.00	2	5.99	0.000
3	Trawl	Large	N_MA	1day	37.19	9	16.92	0.000
4	Trawl	Small	NJ/NY	1day	15.00	2	5.99	0.001
4	Trawl	Small	N_MA	2+day	14.00	2	5.99	0.001
1	Trawl	Large	NC/VA	2+day	29.65	13	22.36	0.005
2	Trawl	Small	DE/MD	1day	8.67	3	7.81	0.034
1	Trawl	Medium	SNE	2+day	4.00	1	3.84	0.046
2	Trawl	Large	NC/VA	2+day	14.28	8	15.51	0.075
2	Trawl	Large	N_MA	2+day	22.66	15	25.00	0.092
2	Trawl	Small	NJ/NY	1day	13.22	8	15.51	0.105
2	Trawl	Large	DE/MD	2+day	13.03	8	15.51	0.111
4	Trawl	Large	SNE	2+day	2.00	1	3.84	0.157
3	Trawl	Large	ME_NH	1day	14.30	10	18.31	0.160
4	Trawl	Large	NC/VA	2+day	19.92	15	25.00	0.175
2	Trawl	Small	NJ/NY	2+day	7.58	5	11.07	0.181
3	Trawl	Small	NJ/NY	1day	1.00	1	3.84	0.317
1	Trawl	Large	SNE	2+day	3.81	4	9.49	0.432
4	Trawl	Small	N_MA	1day	0.60	1	3.84	0.439
2	Trawl	Medium	N_MA	1day	0.50	1	3.84	0.480
4	Trawl	Large	NC/VA	1day	7.45	8	15.51	0.489
2	Trawl	Large	DE/MD	1day	0.41	1	3.84	0.520
4	Trawl	Small	NJ/NY	2+day	8.01	9	16.92	0.533
4	Trawl	Medium	NC/VA	2+day	0.33	1	3.84	0.564
2	Trawl	Small	SNE	1day	1.00	2	5.99	0.607
4	Trawl	Large	N_MA	1day	5.25	7	14.07	0.630
1	Trawl	Small	N_MA	2+day	1.67	3	7.81	0.644
1	Trawl	Large	NJ/NY	1day	3.08	5	11.07	0.687
4	Trawl	Large	NJ/NY	2+day	0.71	2	5.99	0.700
1	Trawl	Large	N_MA	1day	6.29	10	18.31	0.790
3	Trawl	Large	ME_NH	2+day	3.02	6	12.59	0.807
4	Trawl	Large	N_MA	2+day	5.87	10	18.31	0.826
1	Trawl	Large	N_MA	2+day	1.08	4	9.49	0.897
1	Trawl	Large	ME_NH	1day	3.40	8	15.51	0.907
3	Trawl	Large	N_MA	2+day	2.06	6	12.59	0.914
1	Trawl	Large	NJ/NY	2+day	2.00	6	12.59	0.920
4	Trawl	Large	ME_NH	1day	0.39	3	7.81	0.943
2	Trawl	Large	ME_NH	2+day	4.43	11	19.68	0.956
1	Trawl	Large	ME_NH	2+day	0.85	6	12.59	0.991
3	Trawl	Large	DE/MD	1day	0.81	6	12.59	0.992
2	Trawl	Large	ME_NH	1day	1.67	9	16.92	0.996

Overview of Optimization Process





Figure 1. An overview of the optimization process used to allocate sea days to fisheries in the Northeast region.


**Number of trips in 2003/2004 VTR data subsets
for otter trawl, gillnet and longline trips
(43,703 trips)**



<p>Total Unique Trips: 43,703</p> <p>Total Trips with Overlap: 21,429</p> <p>Sum of Trip Sets: 67,132</p>
--

Figure 2. Number of trips in the 2003/2004 Vessel Trip Report (VTR), by data subsets (New England groundfish -NEGF; Monkfish - MONK; and summer flounder, scup and black sea bass - FSB) for otter trawl, gillnet and longline trips.


**Number of trips and sea days
in the 2003/2004 Observer data subsets
for otter trawl, gillnet and longline trips
(1,103 trips and 2,704 sea days)**



<p>Total Unique Trips: 1,103 Total Trips with Overlap: 817 Sum of Trip Sets: 2,105</p>

Figure 3. Number of trips and sea days in the 2003/2004 Northeast Fisheries Observer Program, by data subsets (New England groundfish - NEFG; Monkfish - MONK; and summer flounder, scup and black sea bass - FSB) for otter trawl, gillnet and longline trips.

**Sampling Fraction: 2003/2004 Observer trips/VTR trips
for otter trawl, gillnet and longline trips
(43,703 unique trips)**



Total Unique Trips: 2.5% (1,103 / 43,703)
Total Trips with Overlap: 3.8% (817/ 21,429)
Sum of Trip Sets: 3.1% (2,105 / 67,132)

Figure 4. The sampling fraction of 2003/2004 Observed trips to Vessel Trip Report trips, by data subset (New England groundfish - NEGF; Monkfish - MONK; and summer flounder, scup and black sea bass - FSB) for otter trawl, gillnet and longline trips.

Comparisons of Ave Kept (lb)





Figure 5. Comparison of average kept pounds of groundfish (natural log scale) in the Northeast Fisheries Observer Program and Vessel Trip Report data sets for 2003/2004. Each point represents the mean of an individual stratum.

VTR vs Obsrvr Ave Kept Comparison



VTR vs Obsrvr SD Kept Comparison




Figure 6. The distribution of differences between the average kept pounds (A) and the standard deviation (SD) of average kept pounds (B) of groundfish in the Northeast Fisheries Observer Program (Obsrvr) and the Vessel Trip Report (VTR) data for 2003/2004. Histograms are non-parametric smooths of the stratum specific differences.

Comparisons of Ave Trip Duration





Figure 7. Comparison of average trip duration (in days) for trips that caught groundfish in the Northeast Fisheries Observer Program and Vessel Trip Report (VTR) data sets for 2003/2004. Each point represents the mean of an individual stratum.

Ave Trip Duration Comparison



SD Trip Duration Comparison




Figure 8. The distribution of differences in average trip duration (in days) (A) and the standard deviation of average trip duration (B) of trips that caught groundfish in the Northeast Fisheries Observer Program (Obsrvr) and the Vessel Trip Report (VTR) data for 2003/2004. Histograms are non-parametric smooths of the stratum specific differences.




Figure 9. Locations of otter trawl fishing effort (color squares) in 2003 from vessels using VMS (vessel monitoring systems). Locations are plotted only for vessels speeds ≤ 3.5 knots and data are aggregated to 1' square. Blue squares represent 1-8 hours, green 9 – 25 hours; yellow 26-63 hours; orange 64 – 145 hours, and red 146 – 309 hours. Observed otter trawl tows (white circles) in 2003. Locations are the starting positions of each tow. Taken from Murawski et al. (article in press).




Figure 10. The optimized coefficient of variation (CV) of the discard to kept ratio (d/k) for New England groundfish over a range of sea days; 2,708 sea days (solid circle) are allocated to cover New England groundfish fisheries in 2005.

New England Groundfish (otter trawl gear)




Figure 11. The 2003/2004 point estimates of the coefficient of variation (CV) of the discard to kept (d/k) ratio for New England groundfish caught with otter trawl gear, and the expected coefficient of variation of the discard to kept ratio over a range of sample sizes (number of trips).

New England Groundfish (gillnet gear)




Figure 12. The 2003/2004 point estimates of the coefficient of variation (CV) of the discard to kept (d/k) ratio for New England groundfish caught with gillnet gear, and the expected coefficient of variation of the discard to kept ratio over a range of sample sizes (number of trips).

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance: All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material lifted from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

Organization: Manuscripts must have an abstract and table of contents, and — if applicable — lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area"/"Experimental Apparatus," "Methods," "Results," "Discussion" and/or "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style: The CRD series is obligated to conform with the style contained in the current edition of the *United States Government Printing Office Style Manual*. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the *CBE Style Manual*. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the International Standardization Organization's guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation: Type a clean/neat, single-spaced version of the document. The document must be paginated continuously from beginning to end and must have a "Table of Contents." Begin the preliminary pages of the document — always the "Table of Contents" — with page "iii." Begin the body of the document — normally the "Introduction" — with page "1," and continuously paginate all pages including tables, figures, appendices, and indices. You can insert blank pages as appropriate throughout the document, but account for them in your pagination (*e.g.*, if your last figure ends on an odd-numbered/right-hand page such as "75," and if your next page is the first page of an appendix, then you would normally insert a blank page after the last figure, and paginate the first page of the appendix as "77" to make it begin on an odd-numbered/right-hand page also). Forward the final version to the Editorial Office as both a paper copy and electronically (*i.e.*, e-mail attachment, 3.5-inch floppy disk, high-density zip disk, or CD). For purposes of publishing the CRD series only, the use of Microsoft Word is preferable to the use of Corel WordPerfect.

Production and Distribution: The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages (pages "i" and "ii") of the document, then combine those covers and preliminary pages with the text that you have supplied. The document will then be issued online.

Paper copies of the four covers and two preliminary pages will be sent to the sole/senior NEFSC author should he/she wish to prepare some paper copies of the overall document as well. The Editorial Office will only produce three paper copies (*i.e.*, two copies for the NEFSC's libraries and one copy for its own archives) of the overall document.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the online version of the document. The sole/senior NEFSC author of the document will receive a list of those so notified.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review, but no technical or copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. There is no scientific review, nor any technical or copy editing, of this report.

OBTAINING A COPY: To obtain a copy of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, or to subscribe to the *Resource Survey Report*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2228) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.

Appendix B
Detailed Tables and Figures from Chapter 5

This page intentionally left blank.




Figure B-1a. Comparison of bluefish discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1b. Comparison of bluefish discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1c. Comparison of bluefish discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1d. Comparison of bluefish discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1e. Comparison of herring discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1f. Comparison of herring discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1g. Comparison of herring discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1h. Comparison of herring discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1i. Comparison of red crab discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1j. Comparison of red crab discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1k. Comparison of red crab discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1l. Comparison of red crab discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1m. Comparison of scallop discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1n. Comparison of scallop discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1o. Comparison of scallop discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1p. Comparison of scallop discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1q. Comparison of squid-butterfish-mackerel discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1r. Comparison of squid-butterfish-mackerel discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1s. Comparison of squid-butterfish-mackerel discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1t. Comparison of squid-butterfish-mackerel discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1u. Comparison of monkfish discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1v. Comparison of monkfish discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1w. Comparison of monkfish discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1x. Comparison of monkfish discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1y. Comparison of Northeast multispecies (large-mesh) discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1z. Comparison of Northeast multispecies (large-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1aa. Comparison of Northeast multispecies (large-mesh) discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1bb. Comparison of Northeast multispecies (large-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1cc. Comparison of Northeast multispecies (small-mesh) discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1dd. Comparison of Northeast multispecies (small-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ee. Comparison of Northeast multispecies (small-mesh) discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ff. Comparison of Northeast multispecies (small-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1gg. Comparison of skates discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1hh. Comparison of skates discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ii. Comparison of skates discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1jj. Comparison of skates discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1kk. Comparison of spiny dogfish discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ll. Comparison of spiny dogfish discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1mm. Comparison of spiny dogfish discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1nn. Comparison of spiny dogfish discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-100. Comparison of fluke-scup-black sea bass discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (; sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1pp. Comparison of fluke-scup-black sea bass discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1qq. Comparison of fluke-scup-black sea bass discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1rr. Comparison of fluke-scup-black sea bass discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ss. Comparison of surfclams/quahogs discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1tt. Comparison of surfclams/quahogs discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1uu. Comparison of tilefish discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm <5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1vv. Comparison of tilefish discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip.




Figure B-1ww. Comparison of tilefish discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-1xx. Comparison of tilefish discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip.




Figure B-2a. Comparison of sea turtles and trip duration (days) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2b. Comparison of sea turtles and kept weight of all species (pounds) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2c. Comparison of seals and trip duration (days) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2d. Comparison of seals and kept weight of all species (pounds) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2e. Comparison of whales and trip duration (days) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2f. Comparison of whales and kept weight of all species (pounds) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2g. Comparison of dolphins/porpoises and trip duration (days) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2h. Comparison of dolphins/porpoises and kept weight of all species (pounds) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2i. Comparison of sea birds and trip duration (days) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-2j. Comparison of sea birds and kept weight of all species (pounds) from 2004 observed longline (010), otter trawl (050), gillnet (100) and scallop dredge (132) trips, by region; each dot represents a trip.




Figure B-3a. Comparison of Northeast multispecies (large-mesh) discards (pounds) and trip duration (days) from 2004 observed otter trawl trips, by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip. Trips with zero discards of Northeast multispecies (large-mesh) are excluded.




Figure B-3b. Comparison of Northeast multispecies (large-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed otter trawl trips by region and mesh size group (sm < 5.5 inches, and lg => 5.5 inches); fourth root transformation used, each dot represents a trip. Trips with zero discards of Northeast multispecies (large-mesh) are excluded




Figure B-3c. Comparison of Northeast multispecies (large-mesh) discards (pounds) and trip duration (days) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip. Trips with zero discards of Northeast multispecies (large-mesh) are excluded.




Figure B-3d. Comparison of Northeast multispecies (large-mesh) discards (pounds) and kept weight of all species (pounds) from 2004 observed gillnet trips by region and mesh size group (lg = 5.5 to 7.99 inches; sm < 5.5 inches, and xlg > 8 inches); fourth root transformation used, each dot represents a trip. Trips with zero discards of Northeast multispecies (large-mesh) are excluded.

010,NE




Figure B-4a. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England longline; each dot represents a species group and mesh size.

050,MA




Figure B-4b. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic otter trawl; each dot represents a species group and mesh size.

050,NE




Figure B-4c. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England otter trawl; each dot represents a species group and mesh size.

132,MA




Figure B-4d. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic scallop dredge; each dot represents a species group and mesh size.

132,NE




Figure B-4e. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England scallop dredge; each dot represents a species group and mesh size.

100,MA




Figure B-4f. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic gillnet; each dot represents a species group and mesh size.

100,NE




Figure B-4g. Comparisons of the total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England gillnet; each dot represents a species group and mesh size.

010,NE




Figure B-5a. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England longline; each dot represents a species group and mesh size.

050,MA




Figure B-5b. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic otter trawl; each dot represents a species group and mesh size.

050,NE




Figure B-5c. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England otter trawl; each dot represents a species group and mesh size.

132,MA




Figure B-5d. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic scallop dredge each dot represents a species group and mesh size

132,NE




Figure B-5e. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England scallop dredge each dot represents a species group and mesh size

100,MA




Figure B-5f. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic gillnet each dot represents a species group and mesh size.

100,NE




Figure B-5g. Comparisons of the standard error (SE) of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England gillnet each dot represents a species group and mesh size.

010,NE




Figure B-5h. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England longline; each dot represents a species group and mesh size.

050,MA




Figure B-5i. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic otter trawl; each dot represents a species group and mesh size.

050,NE




Figure B-5j Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England otter trawl; each dot represents a species group and mesh size.

132,MA




Figure B-5k. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic scallop dredge each dot represents a species group and mesh size

132,NE




Figure B-5l. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England scallop dredge each dot represents a species group and mesh size

100,MA




Figure B-5m. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for Mid-Atlantic gillnet each dot represents a species group and mesh size.

100,NE





Figure B-5n. Comparisons of the CV of total discards derived by the two bycatch ratios (discard-to-days-absent [DDA] and discard-to-kept [DK]) and the three methods (separate ratio [D1], combined ratio [D2] and simple expansion [D3]) for New England gillnet each dot represents a species group and mesh size.

Longline with Region = NE



Longline with Region = NE






Figure B-6a. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for New England longline; each dot represents a species group and mesh size.


Otter Trawl with Region = NE



Otter Trawl Region = MA



Otter Trawl with Region = NE



Otter Trawl Region = MA






Figure B-6b. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for New England otter trawl; each dot represents a species group and mesh size.

Figure B-6c. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for Mid-Atlantic otter trawl; each dot represents a species group and mesh size.


Scallop Dredge with Region = NE



Scallop Dredge with Region = MA



Scallop Dredge with Region = NE



Scallop Dredge with Region = MA






Figure B-6d. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for New England scallop dredge; each dot represents a species group and mesh size.

Figure B-6e. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for Mid-Atlantic otter trawl; each dot represents a species group and mesh size.


Gillnet with Region = NE



Gillnet with Region = MA



Gillnet with Region = NE



Gillnet with Region = MA





Figure B-6f. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for New England gillnet; each dot represents a species group and mesh size.

Figure B-6g. Comparisons of CV of total discards estimated via the combined ratio method (CVD2) and the simple expansion method (CVD3) for discard-to-days-absent (DDA), top panel, and discard-to-kept (DK), bottom panel, for Mid-Atlantic gillnet; each dot represents a species group and mesh size.


Bluefish

Comparisons of Avg Kept (lb)




Fluke-Scup-Black Sea Bass

Comparisons of Avg Kept (lb)



Spiny Dogfish

Comparisons of Avg Kept (lb)



Northeast multispecies (Large-mesh)

Comparisons of Avg Kept (lb)





Figure B-7. Comparisons of average kept pounds (fourth root transformation used), by species group, in the Northeast Fisheries Observer Program and FVTR data sets for 2004. Each dot represents the mean of an individual stratum (fleet).


Northeast multispecies (Small-mesh)

Comparisons of Avg Kept (lb)




Monkfish

Comparisons of Avg Kept (lb)



Herring

Comparisons of Avg Kept (lb)



Red Crab

Comparisons of Avg Kept (lb)





Figure B-7 continued. Comparisons of average kept pounds (fourth root transformation used), by species group, in the Northeast Fisheries Observer Program and FVTR data sets for 2004. Each dot represents the mean of an individual stratum (fleet).


Mackerel-Squid-Butterfish

Comparisons of Avg Kept (lb)




Surfclam – Ocean Quahog

Comparisons of Avg Kept (lb)



Scallops

Comparisons of Avg Kept (lb)



Skate Complex

Comparisons of Avg Kept (lb)





Figure B-7 continued. Comparisons of average kept pounds (fourth root transformation used), by species group, in the Northeast Fisheries Observer Program and FVTR data sets for 2004. Each dot represents the mean of an individual stratum (fleet).

Tilefish

Comparisons of Avg Kept (lb)



All Species

Comparisons of Avg Kept (lb)





Figure B-7 continued. Comparisons of average kept pounds (fourth root transformation used), by species group, in the Northeast Fisheries Observer Program and FVTR data sets for 2004. Each dot represents the mean of an individual stratum (fleet).


Bluefish

VTR vs Obsrvr Ave Kept Comparison




Fluke-Scup-Black Sea Bass

VTR vs Obsrvr Ave Kept Comparison



Spiny Dogfish

VTR vs Obsrvr Ave Kept Comparison



Northeast multispecies (Large-mesh)

VTR vs Obsrvr Ave Kept Comparison





Figure B-8. The distribution of differences in the average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.


Northeast multispecies (Small-mesh)

VTR vs Obsrvr Ave Kept Comparison




Monkfish

VTR vs Obsrvr Ave Kept Comparison



Herring

VTR vs Obsrvr Ave Kept Comparison



Mackerel-Squid-Butterfish

VTR vs Obsrvr Ave Kept Comparison





Figure B-8 continued. The distribution of differences in the average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.


Scallops

VTR vs Obsrvr Ave Kept Comparison




Tilefish

VTR vs Obsrvr Ave Kept Comparison



Skate Complex

VTR vs Obsrvr Ave Kept Comparison



All species

VTR vs Obsrvr Ave Kept Comparison





Figure B-8 continued. The distribution of differences in the average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.


Bluefish

VTR vs Obsrvr SD Kept Comparison




Fluke-Scup-Black Sea Bass

VTR vs Obsrvr SD Kept Comparison



Spiny Dogfish

VTR vs Obsrvr SD Kept Comparison



Northeast multispecies (Large-mesh)

VTR vs Obsrvr SD Kept Comparison





Figure B-9. The distribution of difference between the standard deviation of average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.


Northeast multispecies (small-mesh)

VTR vs Obsrvr SD Kept Comparison




Monkfish

VTR vs Obsrvr SD Kept Comparison



Herring

VTR vs Obsrvr SD Kept Comparison



Mackerel-Squid-butterfish

VTR vs Obsrvr SD Kept Comparison





Figure B-9 continued. The distribution of difference between the standard deviation of average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.


Scallop

VTR vs Obsrvr SD Kept Comparison




Tilefish

VTR vs Obsrvr SD Kept Comparison



Skate Complex

VTR vs Obsrvr SD Kept Comparison



All Species

VTR vs Obsrvr SD Kept Comparison





Figure B-9 continued. The distribution of difference between the standard deviation of average kept pounds of species groups in the Northeast Fisheries Observer Program and the FVTR data for 2004.

ALL TRIPS

Comparisons of Avg Trip Duration



ALL TRIPS

Avg Trip Duration Comparison




Figure B-10. Comparison of average trip duration (days) for all trips in the Northeast Fisheries Observer Program and FVTR data sets for 2004. Each dot represents the mean of an individual stratum (fleet).

SD Trip Duration Comparison




Figure B-11. The distribution of differences between the average trip duration (top), and standard deviation of average trip duration (bottom), for trips in the Northeast Fisheries Observer Program and the FVTR data for 2004

Table B-1. Precision (CV) of total composite discards, by species and fleet, based on 2004 observer data .

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	BLUEFISH	HERRING	SALMON	RED CRAB	SCALLOP	MACK/SQUID-/BUTTERFISH	Mackerel	Illex	Loligo	Butterfish	MONKFISH
Longline	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*
Longline	all	all	MA	all											
Otter Trawl	all	all	NE	small	0.508	0.437	*	0.428	0.710	0.227	0.634	0.320	0.309	0.366	0.405
Otter Trawl	all	all	NE	large	2.474	1.313	*	0.280	0.350	0.572	0.520	1.097	0.610	0.756	0.088
Otter Trawl	all	all	MA	small	0.903	0.784	*	1.394	0.574	0.561	1.044	0.635	0.735	0.571	0.354
Otter Trawl	all	all	MA	large	1.906	0.775	*	*	0.444	0.390	0.489	0.710	0.456	0.502	0.295
Scallop Trawl	open	limited	MA	all	*	*	*	*	0.000	0.000	*	*	0.000	*	0.000
Scallop Trawl	open	general	MA	all	1.141	*	*	0.640	0.224	0.354	*	0.343	0.252	0.976	0.194
Shrimp Trawl	all	all	NE	all	*	0.479	*	*	0.965	0.981	*	*	*	0.981	0.235
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	0.000	0.000	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	large	0.220	0.229	*	0.625	0.969	0.841	0.876	1.067	*	1.520	0.210
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	0.181	0.378	*	0.998	0.421	0.498	0.500	*	*	0.906	0.174
Sink, Anchor, Drift Gillnet	all	all	MA	small	*	*	*	*	*	0.000	*	*	*	0.000	*
Sink, Anchor, Drift Gillnet	all	all	MA	large	1.216	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	0.304	*	*	*	0.587	*	*	*	*	*	0.273
Scallop Dredge	open	limited	NE	all	*	*	*	0.842	0.159	0.689	*	0.490	1.112	1.662	0.319
Scallop Dredge	open	limited	MA	all	*	*	*	1.304	0.200	0.305	1.304	0.514	0.383	0.620	0.174
Scallop Dredge	open	general	NE	all	*	*	*	*	0.094	1.274	*	1.274	*	*	0.560
Scallop Dredge	open	general	MA	all	*	*	*	*	0.359	0.865	*	*	0.865	*	0.202
Scallop Dredge	closed	limited	NE	all	1.077	0.168	*	0.482	0.135	0.421	0.167	0.255	0.468	0.158	0.222
Scallop Dredge	closed	limited	MA	all	1.208	0.660	*	0.357	0.198	0.310	0.648	0.338	0.638	0.303	0.280
Scallop Dredge	closed	general	NE	all											
Scallop Dredge	closed	general	MA	all	*	*	*	*	0.000	*	*	*	*	*	0.000
Mid-water paired & single Trawl	all	all	NE	all	0.770	0.770	*	*	1.464	0.429	0.430	0.872	1.457	1.387	0.724
Mid-water paired & single Trawl	all	all	MA	all	0.539	0.982	*	*	*	0.545	0.539	0.546	0.539	0.539	1.048
Fish Pots/ Traps	all	all	NE	all											
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	0.408
Purse Seine	all	all	NE	all	*	0.981	*	*	*	0.935	*	0.935	*	*	*
Purse Seine	all	all	MA	all											
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	MA	all											
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*
Clam Quahog Dredge	all	all	NE	all											
Clam Quahog Dredge	all	all	MA	all											
Crab Pots	all	all	NE	all											
Crab Pots	all	all	MA	all											
Lobster Pots	all	all	NE	all											
Lobster Pots	all	all	MA	all											

Note: when discard ratio = 0, CV is null (*); Gray-shaded cells indicate unlikely species/gear combinations.

Table B-1 continued. Precision (CV) of total composite discards, by species and fleet, based on 2004 observer data .

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	NE MULTI-SPP (LARGE-MESH)														
					Cod	Haddock	Yellowtail fltd	American plaice	Witch fltd	Winter fltd	Pollock	Redfish	White hake	Windowpane	Hallbut	Ocean pout			
Longline	all	all	NE	all	0.335	0.401	0.389	*	*	*	*	1.191	*	*	*	*	0.569		
Longline	all	all	MA	all															
Otter Trawl	all	all	NE	small	0.233	0.658	0.696	0.409	0.304	0.332	0.430	0.546	0.593	0.459	0.291	0.753	0.321		
Otter Trawl	all	all	NE	large	0.101	0.176	0.265	0.222	0.254	0.145	0.429	0.640	0.248	0.235	0.206	0.424	0.161		
Otter Trawl	all	all	MA	small	0.326	*	*	1.081	1.476	0.489	0.561	*	0.905	0.989	0.399	*	1.506		
Otter Trawl	all	all	MA	large	0.251	3.122	*	0.669	*	0.292	0.413	3.122	0.974	3.133	0.312	*	0.477		
Scallop Trawl	open	limited	MA	all	0.000	*	*	*	*	*	*	*	*	*	0.000	*	*		
Scallop Trawl	open	general	MA	all	0.170	*	*	1.036	*	0.471	0.464	*	*	0.640	0.237	*	*		
Shrimp Trawl	all	all	NE	all	0.224	0.352	0.659	0.552	0.305	0.928	0.269	0.473	0.374	0.232	0.207	*	0.960		
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	NE	large	0.092	0.121	0.186	0.198	0.281	0.406	0.288	0.182	0.261	0.231	0.432	0.449	0.437		
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	0.159	0.175	0.246	0.361	0.337	1.018	0.557	0.317	0.364	0.372	0.815	0.436	0.421		
Sink, Anchor, Drift Gillnet	all	all	MA	small	*	*	*	*	*	*	*	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	MA	large	0.868	*	*	*	*	*	*	*	*	*	0.868	*	*		
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	*	*	*	*	*	*	*	*	*	*	*	*	*		
Scallop Dredge	open	limited	NE	all	0.480	0.850	0.848	0.637	0.848	0.485	1.022	0.848	*	0.525	0.454	*	0.656		
Scallop Dredge	open	limited	MA	all	0.242	*	*	0.705	0.809	0.496	0.581	*	*	0.521	0.323	*	1.091		
Scallop Dredge	open	general	NE	all	0.358	1.226	*	0.494	0.908	0.902	0.213	*	*	*	0.438	*	1.287		
Scallop Dredge	open	general	MA	all	0.311	*	*	0.865	0.857	0.650	0.421	*	*	0.653	0.333	*	*		
Scallop Dredge	closed	limited	NE	all	0.159	0.510	0.423	0.211	0.829	0.188	0.200	*	*	0.478	0.355	0.179	0.427		
Scallop Dredge	closed	limited	MA	all	0.712	*	*	1.256	0.320	0.350	1.269	*	*	0.602	0.886	*	1.239		
Scallop Dredge	closed	general	NE	all															
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*		
Mid-water paired & single Trawl	all	all	NE	all	0.669	1.198	0.951	*	1.155	1.203	1.298	0.967	0.996	1.604	*	*	*		
Mid-water paired & single Trawl	all	all	MA	all	0.708	*	*	*	*	1.146	*	*	*	0.541	*	*	*		
Fish Pots/ Traps	all	all	NE	all															
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*		
Purse Seine	all	all	NE	all	0.973	*	*	*	*	*	*	*	0.973	*	*	*	*		
Purse Seine	all	all	MA	all															
Hand Line	all	all	NE	all	4.030	4.030	*	*	*	*	*	*	*	*	*	*	*		
Hand Line	all	all	MA	all															
Scottish Seine	all	all	NE	all	0.289	0.279	0.279	*	0.279	*	0.543	*	*	0.279	0.354	*	*		
Clam Quahog Dredge	all	all	NE	all															
Clam Quahog Dredge	all	all	MA	all															
Crab Pots	all	all	NE	all															
Crab Pots	all	all	MA	all															
Lobster Pots	all	all	NE	all															
Lobster Pots	all	all	MA	all															

Note: when discard ratio = 0, CV is null (*); Gray-shaded cells indicate unlikely species/gear combinations.

Table B-1 continued. Precision (CV) of total composite discards, by species and fleet, based on 2004 observer data .

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups														
					NE MUL TI-SPP (SMALL-MESH)	Silver hake	Offshore hake	Red hake	SKATE	DOG FISH	FLUKE/SCUPI-BLK SEA BASS	Fluke	Scup	Black sea bass	SURF CLAM/OCEAN QUAHOG	TILE FISH		
Longline	all	all	NE	all	0.910	*	*	0.910	0.614	0.654	*	*	*	*	*	*		
Longline	all	all	MA	all														
Otter Trawl	all	all	NE	small	0.235	0.219	1.511	0.406	0.691	0.322	0.309	0.276	0.551	0.708	1.028	0.304		
Otter Trawl	all	all	NE	large	0.182	0.227	0.322	0.353	0.175	0.245	0.319	0.328	0.918	0.833	1.512	0.529		
Otter Trawl	all	all	MA	small	0.508	0.625	0.683	0.587	0.222	0.367	0.386	0.278	0.560	0.502	0.464	1.155		
Otter Trawl	all	all	MA	large	0.827	0.451	*	1.811	0.209	0.557	0.246	0.266	0.354	0.652	0.609	*		
Scallop Trawl	open	limited	MA	all	*	*	*	*	0.000	*	0.000	0.000	*	0.000	*	*		
Scallop Trawl	open	general	MA	all	0.496	0.508	*	1.141	0.347	0.675	0.505	0.608	0.731	0.638	*	*		
Shrimp Trawl	all	all	NE	all	0.557	0.567	*	0.537	0.799	0.960	*	*	*	*	*	*		
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	0.000	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	NE	large	0.183	0.238	*	0.219	0.228	0.106	0.845	0.898	*	1.602	*	*		
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	0.624	0.207	*	0.864	0.117	0.162	0.233	0.233	0.904	*	*	0.256		
Sink, Anchor, Drift Gillnet	all	all	MA	small	*	*	*	*	*	0.000	0.000	0.000	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	MA	large	*	*	*	*	1.118	1.083	*	*	*	*	*	*		
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	*	*	*	*	0.115	0.129	0.303	0.303	*	*	*	*		
Scallop Dredge	open	limited	NE	all	0.414	0.764	1.173	0.352	0.236	0.515	0.458	0.474	0.322	0.622	0.391	*		
Scallop Dredge	open	limited	MA	all	0.758	0.856	0.738	0.402	0.126	0.230	0.259	0.272	0.704	0.558	0.771	*		
Scallop Dredge	open	general	NE	all	0.104	1.300	*	0.103	0.177	0.318	0.092	0.092	*	*	1.287	*		
Scallop Dredge	open	general	MA	all	0.482	0.467	*	0.857	0.202	0.550	0.461	0.461	*	*	0.830	*		
Scallop Dredge	closed	limited	NE	all	0.396	0.403	0.489	0.448	0.126	0.326	0.291	0.293	0.218	0.161	0.198	*		
Scallop Dredge	closed	limited	MA	all	0.268	0.323	*	0.282	0.142	0.425	0.383	0.385	1.011	0.333	0.321	*		
Scallop Dredge	closed	general	NE	all														
Scallop Dredge	closed	general	MA	all	*	*	*	*	0.000	*	0.000	0.000	*	*	*	*		
Mid-water paired & single Trawl	all	all	NE	all	0.994	1.000	*	0.748	1.177	0.418	0.628	*	0.671	1.626	*	*		
Mid-water paired & single Trawl	all	all	MA	all	0.539	0.539	*	0.539	*	0.246	1.165	1.142	*	1.176	*	*		
Fish Pots/ Traps	all	all	NE	all														
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	0.161	*	0.163	0.161	*	*		
Purse Seine	all	all	NE	all	*	*	*	*	*	0.972	*	*	*	*	*	*		
Purse Seine	all	all	MA	all														
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*		
Hand Line	all	all	MA	all														
Scottish Seine	all	all	NE	all	0.279	0.279	*	0.279	0.319	*	0.253	0.259	0.808	0.808	*	*		
Clam Quahog Dredge	all	all	NE	all														
Clam Quahog Dredge	all	all	MA	all														
Crab Pots	all	all	NE	all														
Crab Pots	all	all	MA	all														
Lobster Pots	all	all	NE	all														
Lobster Pots	all	all	MA	all														

Note: when discard ratio = 0, CV is null (*); Gray-shaded cells indicate unlikely species/gear combinations.

Table B-1 continued. Precision (CV) of total composite discards, by species and fleet, based on 2004 observer data .

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups																
					TURTLES	TURTLE, GREEN	TURTLE, LEATHERBACK	TURTLE, LOGGERHEAD	TURTLE KEMPS RIDLEY	TURTLE, NK	SEALS	SEAL, HARP	SEAL, HOODED	SEAL, HARBOR	SEAL, GRAY	SEAL, NK	WHALES			
Longline	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Otter Trawl	all	all	NE	small	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.931
Otter Trawl	all	all	NE	large	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	1.089
Otter Trawl	all	all	MA	small	0.573	*	*	0.573	*	*	*	*	*	*	*	*	*	*	*	*
Otter Trawl	all	all	MA	large	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Trawl	open	limited	MA	all	0.381	*	*	0.381	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	large	*	*	*	*	*	*	*	0.206	0.293	*	0.273	0.520	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	*	*	*	*	*	*	0.215	0.435	0.751	0.320	0.273	*	*	*	*	
Sink, Anchor, Drift Gillnet	all	all	MA	small	0.626	*	0.787	*	*	1.013	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	MA	large	1.052	1.479	*	1.478	*	*	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	0.495	*	0.730	0.656	*	*	0.692	*	*	1.023	0.924	*	*	*	*	
Scallop Dredge	open	limited	NE	all	0.551	*	*	0.551	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	open	limited	MA	all	0.770	*	*	0.770	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	limited	NE	all	0.165	*	*	0.165	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Mid-water paired & single Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	1.114
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Fish Pots/ Traps	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Clam Quahog Dredge	all	all	NE	all																
Clam Quahog Dredge	all	all	MA	all																
Crab Pots	all	all	NE	all																
Crab Pots	all	all	MA	all																
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Lobster Pots	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*

Note: when discard ratio = 0, CV is null (*); Gray-shaded cells indicate unlikely species/gear combinations.

Table B-1 continued. Precision (CV) of total composite discards, by species and fleet, based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	WHALE, PILOT, LONG-FIN	WHALE, MINKE	WHALE, NK	DOLPHINS/-PORPOISE	DOLPHIN WHITE-SIDED	DOLPHIN-COMMON	DOLPHIN-BOTTLE-NOSE	PORPOISE, HARBOR	PORPOISE/DOLPHIN, NK	SEA BIRDS (ALL)	ALL SPECIES	PILOT coverage
Longline	all	all	NE	all	*	*	*	*	*	*	*	*	*	0.425	0.489	
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	pilot
Otter Trawl	all	all	NE	small	0.931	*	*	0.650	0.936	0.713	*	*	*	0.548	0.193	
Otter Trawl	all	all	NE	large	1.089	*	*	0.389	0.389	*	*	*	*	0.489	0.124	
Otter Trawl	all	all	MA	small	*	*	*	0.557	*	0.557	*	*	*	0.706	0.247	
Otter Trawl	all	all	MA	large	*	*	*	*	*	*	*	*	*	0.672	0.185	
Scallop Trawl	open	limited	MA	all	*	*	*	*	*	*	*	*	*	*	0.000	pilot
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	*	0.243	pilot
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	0.310	
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	0.052	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	*	0.000	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	large	*	*	*	0.359	0.977	*	*	0.384	*	0.342	0.092	
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	*	*	*	0.288	*	*	0.751	0.300	*	0.602	0.085	
Sink, Anchor, Drift Gillnet	all	all	MA	small	*	*	*	*	*	*	*	*	*	0.582	0.000	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	large	*	*	*	*	*	*	*	*	*	0.618	1.078	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	*	*	*	0.924	*	*	*	0.924	*	0.693	0.052	pilot for fish
Scallop Dredge	open	limited	NE	all	*	*	*	*	*	*	*	*	*	0.896	0.197	
Scallop Dredge	open	limited	MA	all	*	*	*	*	*	*	*	*	*	*	0.112	
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	*	0.325	pilot
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	*	0.184	
Scallop Dredge	closed	limited	NE	all	*	*	*	*	*	*	*	*	*	0.163	0.119	
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	*	0.119	
Scallop Dredge	closed	general	NE	all	*	*	*	*	*	*	*	*	*	*		pilot
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	*	0.000	pilot
Mid-water paired & single Trawl	all	all	NE	all	1.114	*	*	0.786	0.786	*	*	*	*	0.554	0.317	
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	0.408	
Fish Pots/ Traps	all	all	NE	all	*	*	*	*	*	*	*	*	*	*		pilot
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	0.137	pilot
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	0.715	
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	*		pilot
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	4.030	pilot
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	*		pilot
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*	0.423	pilot
Clam Quahog Dredge	all	all	NE	all												pilot
Clam Quahog Dredge	all	all	MA	all												pilot
Crab Pots	all	all	NE	all												pilot
Crab Pots	all	all	MA	all												pilot
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	*		pilot
Lobster Pots	all	all	MA	all												pilot

Note: when discard ratio = 0, CV is null (*); Gray-shaded cells indicate unlikely species/gear combinations.

Table B-2. Ranking of total discards within fleet (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	BLUEFISH	HERRING	SALMON	RED CRAB	SCALLOP	Mackerel	Illex	Loligo	Butterfish	MONKFISH
Longline	all	all	NE	all	8	8	*	8	8	8	8	8	8	8
Longline	all	all	MA	all										
Otter Trawl	all	all	NE	small	16	12	*	25	29	6	5	8	4	7
Otter Trawl	all	all	NE	large	22	23	*	12	20	29	27	31	30	3
Otter Trawl	all	all	MA	small	14	22	*	26	15	5	7	11	4	13
Otter Trawl	all	all	MA	large	16	24	*	26	6	21	20	12	15	8
Scallop Trawl	open	limited	MA	all	8	8	*	8	1	8	8	6	8	4
Scallop Trawl	open	general	MA	all	15	20	*	12	2	20	16	9	11	4
Shrimp Trawl	all	all	NE	all	20	1	*	20	17	20	20	20	19	14
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	3	3	*	3	3	2	3	3	3	3
Sink, Anchor, Drift Gillnet	all	all	NE	large	9	15	*	22	24	12	23	27	25	8
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	6	19	*	21	18	8	27	27	23	3
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	4	*	4	4	4	4	4	2	4
Sink, Anchor, Drift Gillnet	all	all	MA	large	2	5	*	5	5	5	5	5	5	5
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	4	7	*	7	6	7	7	7	7	3
Scallop Dredge	open	limited	NE	all	26	26	*	24	1	26	17	16	18	3
Scallop Dredge	open	limited	MA	all	24	24	*	22	1	23	14	12	18	3
Scallop Dredge	open	general	NE	all	17	17	*	17	3	17	14	17	17	1
Scallop Dredge	open	general	MA	all	16	16	*	16	2	16	16	13	16	3
Scallop Dredge	closed	limited	NE	all	19	28	*	25	1	27	20	13	24	3
Scallop Dredge	closed	limited	MA	all	20	19	*	24	1	15	14	12	23	3
Scallop Dredge	closed	general	NE	all										
Scallop Dredge	closed	general	MA	all	5	5	*	5	1	5	5	5	5	3
Mid-water paired & single Trawl	all	all	NE	all	9	3	*	23	21	1	10	15	7	12
Mid-water paired & single Trawl	all	all	MA	all	11	10	*	15	15	14	2	7	9	3
Fish Pots/ Traps	all	all	NE	all										
Fish Pots/ Traps	all	all	MA	all	4	4	*	4	4	4	4	4	4	3
Purse Seine	all	all	NE	all	5	2	*	5	5	5	4	5	5	5
Purse Seine	all	all	MA	all										
Hand Line	all	all	NE	all	2	2	*	2	2	2	2	2	2	2
Hand Line	all	all	MA	all										
Scottish Seine	all	all	NE	all	13	13	*	13	13	13	13	13	13	13
Clam Quahog Dredge	all	all	NE	all										
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all										
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-2 *continued*. Ranking of total discards within fleet (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Cod	Had-dock	Yellowtail fltd	American plaice	Whitch fltd	Winter fltd	Pollock	Redfish	White hake	Window-pane	Hailbut	Ocean pout	
Longline	all	all	NE	all	2	4	8	8	8	8	6	8	8	8	8	5	
Longline	all	all	MA	all													
Otter Trawl	all	all	NE	small	18	13	22	20	14	15	28	24	17	26	30	21	
Otter Trawl	all	all	NE	large	8	10	7	9	5	14	16	13	18	4	21	11	
Otter Trawl	all	all	MA	small	28	28	24	25	17	16	28	21	23	12	28	19	
Otter Trawl	all	all	MA	large	22	26	17	26	9	10	19	25	23	7	26	11	
Scallop Trawl	open	limited	MA	all	8	8	8	8	8	8	8	8	8	3	8	8	
Scallop Trawl	open	general	MA	all	20	20	18	20	17	8	20	20	13	5	20	20	
Shrimp Trawl	all	all	NE	all	8	15	7	3	13	5	6	12	11	9	20	16	
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	
Sink, Anchor, Drift Gillnet	all	all	NE	small	3	3	3	3	3	3	3	3	3	3	3	3	
Sink, Anchor, Drift Gillnet	all	all	NE	large	2	10	5	13	21	7	4	11	6	20	18	17	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	4	13	10	20	26	15	7	22	9	25	17	12	
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	4	4	4	4	4	4	4	4	4	4	4	
Sink, Anchor, Drift Gillnet	all	all	MA	large	5	5	5	5	5	5	5	5	5	4	5	5	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	7	7	7	7	7	7	7	7	7	7	7	7	
Scallop Dredge	open	limited	NE	all	19	22	11	23	7	6	25	26	20	10	26	15	
Scallop Dredge	open	limited	MA	all	24	24	16	21	8	15	24	24	10	6	24	17	
Scallop Dredge	open	general	NE	all	13	17	4	12	10	5	17	17	17	8	17	14	
Scallop Dredge	open	general	MA	all	16	16	13	13	11	6	16	16	10	4	16	16	
Scallop Dredge	closed	limited	NE	all	15	14	4	10	11	9	29	29	16	6	26	21	
Scallop Dredge	closed	limited	MA	all	25	25	6	18	7	11	25	25	16	8	25	22	
Scallop Dredge	closed	general	NE	all													
Scallop Dredge	closed	general	MA	all	5	5	5	5	5	5	5	5	5	5	5	5	
Mid-water paired & single Trawl	all	all	NE	all	16	6	23	13	18	19	8	5	14	23	23	23	
Mid-water paired & single Trawl	all	all	MA	all	15	15	15	15	13	15	15	15	12	15	15	15	
Fish Pots/ Traps	all	all	NE	all													
Fish Pots/ Traps	all	all	MA	all	4	4	4	4	4	4	4	4	4	4	4	4	
Purse Seine	all	all	NE	all	5	5	5	5	5	5	5	3	5	5	5	5	
Purse Seine	all	all	MA	all													
Hand Line	all	all	NE	all	1	2	2	2	2	2	2	2	2	2	2	2	
Hand Line	all	all	MA	all													
Scottish Seine	all	all	NE	all	7	11	13	8	13	9	13	13	11	2	13	13	
Clam Quahog Dredge	all	all	NE	all													
Clam Quahog Dredge	all	all	MA	all													
Crab Pots	all	all	NE	all													
Crab Pots	all	all	MA	all													
Lobster Pots	all	all	NE	all													
Lobster Pots	all	all	MA	all													

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-2 continued. Ranking of total discards within fleet (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Silver hake	Offshore hake	Red hake	SKATE	DOG FISH	Fluke	Scup	Black sea bass	SURF CLAM- OCEAN QUAHOG	TILEFISH
Longline	all	all	NE	all	8	8	7	3	1	8	8	8	8	
Longline	all	all	MA	all										
Otter Trawl	all	all	NE	small	2	23	9	1	3	10	11	19	31	27
Otter Trawl	all	all	NE	large	15	26	19	1	2	6	17	24	28	25
Otter Trawl	all	all	MA	small	6	20	8	1	2	10	3	9	18	27
Otter Trawl	all	all	MA	large	14	26	18	1	2	3	4	5	13	26
Scallop Trawl	open	limited	MA	all	8	8	8	2	8	5	8	7	8	8
Scallop Trawl	open	general	MA	all	7	20	14	1	3	6	10	19	20	20
Shrimp Trawl	all	all	NE	all	2	20	10	4	18	20	20	20	20	20
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	3	3	3	3	1	3	3	3	3	3
Sink, Anchor, Drift Gillnet	all	all	NE	large	14	27	16	3	1	19	27	26	27	27
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	16	27	11	2	1	5	23	27	27	14
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	4	4	4	1	3	4	4	4	4
Sink, Anchor, Drift Gillnet	all	all	MA	large	5	5	5	3	1	5	5	5	5	5
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	7	7	7	2	1	5	7	7	7	7
Scallop Dredge	open	limited	NE	all	13	14	8	2	9	4	21	12	5	26
Scallop Dredge	open	limited	MA	all	9	20	13	2	5	4	19	11	7	24
Scallop Dredge	open	general	NE	all	16	17	9	2	6	7	17	17	11	17
Scallop Dredge	open	general	MA	all	9	16	12	1	8	5	16	16	7	16
Scallop Dredge	closed	limited	NE	all	12	22	7	2	8	5	23	17	18	29
Scallop Dredge	closed	limited	MA	all	10	25	13	2	5	4	17	9	21	25
Scallop Dredge	closed	general	NE	all										
Scallop Dredge	closed	general	MA	all	5	5	5	2	5	4	5	5	5	5
Mid-water paired & single Trawl	all	all	NE	all	4	23	17	11	2	23	20	22	23	23
Mid-water paired & single Trawl	all	all	MA	all	6	15	5	15	1	8	15	4	15	15
Fish Pots/ Traps	all	all	NE	all										
Fish Pots/ Traps	all	all	MA	all	4	4	4	4	4	4	2	1	4	4
Purse Seine	all	all	NE	all	5	5	5	5	1	5	5	5	5	5
Purse Seine	all	all	MA	all										
Hand Line	all	all	NE	all	2	2	2	2	2	2	2	2	2	2
Hand Line	all	all	MA	all										
Scottish Seine	all	all	NE	all	5	13	3	4	13	1	10	6	13	13
Clam Quahog Dredge	all	all	NE	all										
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all										
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-2 continued. Ranking of total discards within fleet (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	TURTLE, GREEN	TURTLE, LEATHERBACK	TURTLE, LOGGERHEAD	TURTLE, KEMP'S RIDLEY	TURTLE, NK	SEAL, HARP	SEAL, HOODED	SEAL, HARBOR	SEAL, GRAY	SEAL, NK
Longline	all	all	NE	all	2	2	2	*	2	2	2	2	2	*
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Otter Trawl	all	all	NE	small	5	5	5	*	5	5	5	5	5	*
Otter Trawl	all	all	NE	large	4	4	4	*	4	4	4	4	4	*
Otter Trawl	all	all	MA	small	4	4	2	*	4	4	4	4	4	*
Otter Trawl	all	all	MA	large	2	2	2	*	2	2	2	2	2	*
Scallop Trawl	open	limited	MA	all	2	2	1	*	2	2	2	2	2	*
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	large	7	7	7	*	7	2	7	3	4	*
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	8	8	8	*	8	4	6	1	3	*
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	2	4	*	3	4	4	4	4	*
Sink, Anchor, Drift Gillnet	all	all	MA	large	2	4	2	*	4	4	4	4	4	*
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	7	4	1	*	7	7	7	6	5	*
Scallop Dredge	open	limited	NE	all	3	3	1	*	3	3	3	3	3	*
Scallop Dredge	open	limited	MA	all	2	2	1	*	2	2	2	2	2	*
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	limited	NE	all	3	3	2	*	3	3	3	3	3	*
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	NE	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	*
Mid-water paired & single Trawl	all	all	NE	all	4	4	4	*	4	4	4	4	4	*
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Fish Pots/ Traps	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Clam Quahog Dredge	all	all	NE	all										
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-2 continued. Ranking of total discards within fleet (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	WHALE: PILOT, LONG-FIN	WHALE: MINKE	WHALE: NK	DOLPHIN WHITE-SIDED	DOLPHIN-COMMON	DOLPHIN-BOTTLE-NOSE	PORPOISE-HARBOR	PORPOISE/DOLP HIN, NK	SEA BIRDS (ALL)	PILOT coverage
Longline	all	all	NE	all	2	*	*	2	2	2	2	*	1	
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Otter Trawl	all	all	NE	small	3	*	*	3	1	5	5	*	2	
Otter Trawl	all	all	NE	large	3	*	*	1	4	4	4	*	2	
Otter Trawl	all	all	MA	small	4	*	*	4	1	4	4	*	3	
Otter Trawl	all	all	MA	large	2	*	*	2	2	2	2	*	1	
Scallop Trawl	open	limited	MA	all	2	*	*	2	2	2	2	*	2	pilot
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	pilot
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	large	7	*	*	6	7	7	5	*	1	
Sink, Anchor, Drift Gillnet	all	all	NE	xlge	8	*	*	8	8	6	2	*	5	
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	*	*	4	4	4	4	*	1	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	large	4	*	*	4	4	4	4	*	1	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	xlge	7	*	*	7	7	7	1	*	3	pilot for fish
Scallop Dredge	open	limited	NE	all	3	*	*	3	3	3	3	*	2	
Scallop Dredge	open	limited	MA	all	2	*	*	2	2	2	2	*	2	
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	pilot
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	
Scallop Dredge	closed	limited	NE	all	3	*	*	3	3	3	3	*	1	
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	
Scallop Dredge	closed	general	NE	all										pilot
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	pilot
Mid-water paired & single Trawl	all	all	NE	all	3	*	*	2	4	4	4	*	1	
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	
Fish Pots/ Traps	all	all	NE	all										pilot
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Clam Quahog Dredge	all	all	NE	all										pilot
Clam Quahog Dredge	all	all	MA	all										pilot
Crab Pots	all	all	NE	all										pilot
Crab Pots	all	all	MA	all										pilot
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Lobster Pots	all	all	MA	all										pilot

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-3. Ranking of total discards within species group (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	BLUEFISH	HERRING	SALMON	RED CRAB	SCALLOP	Mackerel	Illex	Loligo	Butterfish	MONKFISH
Longline	all	all	NE	all	14	13	*	11	19	13	15	14	16	21
Longline	all	all	MA	all										
Otter Trawl	all	all	NE	small	2	2	*	2	13	2	1	1	1	4
Otter Trawl	all	all	NE	large	4	5	*	1	11	7	4	8	7	3
Otter Trawl	all	all	MA	small	3	7	*	6	10	3	2	2	2	11
Otter Trawl	all	all	MA	large	8	9	*	11	7	8	8	3	5	10
Scallop Trawl	open	limited	MA	all	14	13	*	11	3	13	15	6	16	13
Scallop Trawl	open	general	MA	all	11	13	*	3	8	13	13	9	8	14
Shrimp Trawl	all	all	NE	all	14	3	*	11	16	13	15	14	12	19
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	14	13	*	11	19	6	15	14	16	21
Sink, Anchor, Drift Gillnet	all	all	NE	large	7	6	*	4	17	5	10	14	11	15
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	5	8	*	5	15	4	15	14	13	5
Sink, Anchor, Drift Gillnet	all	all	MA	small	14	13	*	11	19	13	15	14	3	21
Sink, Anchor, Drift Gillnet	all	all	MA	large	1	13	*	11	19	13	15	14	16	21
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	6	13	*	11	14	13	15	14	16	12
Scallop Dredge	open	limited	NE	all	14	13	*	7	2	13	5	5	6	1
Scallop Dredge	open	limited	MA	all	14	13	*	8	1	10	7	4	9	2
Scallop Dredge	open	general	NE	all	14	13	*	11	9	13	12	14	16	6
Scallop Dredge	open	general	MA	all	14	13	*	11	6	13	15	13	16	9
Scallop Dredge	closed	limited	NE	all	10	12	*	9	4	12	14	7	14	7
Scallop Dredge	closed	limited	MA	all	13	11	*	10	5	9	11	11	15	8
Scallop Dredge	closed	general	NE	all										
Scallop Dredge	closed	general	MA	all	14	13	*	11	12	13	15	14	16	16
Mid-water paired & single Trawl	all	all	NE	all	9	1	*	11	18	1	6	10	4	17
Mid-water paired & single Trawl	all	all	MA	all	12	10	*	11	19	11	3	12	10	18
Fish Pots/ Traps	all	all	NE	all										
Fish Pots/ Traps	all	all	MA	all	14	13	*	11	19	13	15	14	16	20
Purse Seine	all	all	NE	all	14	4	*	11	19	13	9	14	16	21
Purse Seine	all	all	MA	all										
Hand Line	all	all	NE	all	14	13	*	11	19	13	15	14	16	21
Hand Line	all	all	MA	all										
Scottish Seine	all	all	NE	all	14	13	*	11	19	13	15	14	16	21
Clam Quahog Dredge	all	all	NE	all										
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all										
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-3 continued. Ranking of total discards within species group (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Cod	Had-dock	Yellowtail fld	American plaice	Witch fld	Winter fld	Pollock	Redfish	White hake	Window-pane	Hallbut	Ocean pout	
Longline	all	all	NE	all	3	3	15	15	17	17	8	10	17	18	6	5	
Longline	all	all	MA	all													
Otter Trawl	all	all	NE	small	5	1	3	2	2	1	4	2	1	10	3	2	
Otter Trawl	all	all	NE	large	1	2	1	1	1	2	1	1	2	1	1	1	
Otter Trawl	all	all	MA	small	14	11	12	10	6	4	10	5	7	4	6	6	
Otter Trawl	all	all	MA	large	10	11	9	15	4	5	7	9	13	3	6	3	
Scallop Trawl	open	limited	MA	all	14	11	15	15	17	17	10	10	17	2	6	14	
Scallop Trawl	open	general	MA	all	14	11	14	15	14	13	10	10	12	11	6	14	
Shrimp Trawl	all	all	NE	all	7	8	7	3	10	7	5	6	6	14	6	10	
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*	*	*	
Sink, Anchor, Drift Gillnet	all	all	NE	small	14	11	15	15	17	17	10	10	17	18	6	14	
Sink, Anchor, Drift Gillnet	all	all	NE	large	2	5	4	5	12	10	2	4	4	16	4	8	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	4	6	8	8	16	12	3	8	5	17	2	7	
Sink, Anchor, Drift Gillnet	all	all	MA	small	14	11	15	15	17	17	10	10	17	18	6	14	
Sink, Anchor, Drift Gillnet	all	all	MA	large	14	11	15	15	17	17	10	10	17	6	6	14	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	14	11	15	15	17	17	10	10	17	18	6	14	
Scallop Dredge	open	limited	NE	all	8	9	5	13	3	3	9	10	9	9	6	4	
Scallop Dredge	open	limited	MA	all	14	11	11	12	5	11	10	10	3	5	6	9	
Scallop Dredge	open	general	NE	all	13	11	6	7	9	8	10	10	17	12	6	11	
Scallop Dredge	open	general	MA	all	14	11	13	11	11	9	10	10	10	8	6	14	
Scallop Dredge	closed	limited	NE	all	9	7	2	4	7	6	10	10	11	7	5	12	
Scallop Dredge	closed	limited	MA	all	14	11	10	14	8	14	10	10	14	15	6	13	
Scallop Dredge	closed	general	NE	all													
Scallop Dredge	closed	general	MA	all	14	11	15	15	17	17	10	10	17	18	6	14	
Mid-water paired & single Trawl	all	all	NE	all	11	4	15	6	13	16	6	3	8	18	6	14	
Mid-water paired & single Trawl	all	all	MA	all	14	11	15	15	15	17	10	10	15	18	6	14	
Fish Pots/ Traps	all	all	NE	all													
Fish Pots/ Traps	all	all	MA	all	14	11	15	15	17	17	10	10	17	18	6	14	
Purse Seine	all	all	NE	all	14	11	15	15	17	17	10	7	17	18	6	14	
Purse Seine	all	all	MA	all													
Hand Line	all	all	NE	all	6	11	15	15	17	17	10	10	17	18	6	14	
Hand Line	all	all	MA	all													
Scottish Seine	all	all	NE	all	12	10	15	9	17	15	10	10	16	13	6	14	
Clam Quahog Dredge	all	all	NE	all													
Clam Quahog Dredge	all	all	MA	all													
Crab Pots	all	all	NE	all													
Crab Pots	all	all	MA	all													
Lobster Pots	all	all	NE	all													
Lobster Pots	all	all	MA	all													

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-3 continued. Ranking of total discards within species group (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Silver hake	Offshore hake	Red hake	SKATE	DOG FISH	Fluke	Scup	Black sea bass	SURF CLAM/OCEAN QUAHOG	TILEFISH
Longline	all	all	NE	all	18	7	13	17	10	20	14	16	11	5
Longline	all	all	MA	all										
Otter Trawl	all	all	NE	small	1	1	1	3	4	1	2	4	8	1
Otter Trawl	all	all	NE	large	3	3	3	1	3	2	4	7	6	2
Otter Trawl	all	all	MA	small	2	4	2	6	7	6	1	2	5	4
Otter Trawl	all	all	MA	large	8	7	8	5	5	3	3	3	3	5
Scallop Trawl	open	limited	MA	all	18	7	19	8	23	11	14	10	11	5
Scallop Trawl	open	general	MA	all	11	7	16	10	13	16	5	14	11	5
Shrimp Trawl	all	all	NE	all	4	7	10	18	22	20	14	16	11	5
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	18	7	19	22	20	20	14	16	11	5
Sink, Anchor, Drift Gillnet	all	all	NE	large	9	7	11	15	2	17	14	13	11	5
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	13	7	9	11	8	9	13	16	11	3
Sink, Anchor, Drift Gillnet	all	all	MA	small	18	7	19	22	6	13	14	16	11	5
Sink, Anchor, Drift Gillnet	all	all	MA	large	18	7	19	16	1	20	14	16	11	5
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	18	7	19	14	12	15	14	16	11	5
Scallop Dredge	open	limited	NE	all	7	2	4	2	16	4	7	5	1	5
Scallop Dredge	open	limited	MA	all	6	5	6	4	14	5	8	6	2	5
Scallop Dredge	open	general	NE	all	17	7	7	13	19	14	14	16	7	5
Scallop Dredge	open	general	MA	all	12	7	15	7	21	10	14	16	4	5
Scallop Dredge	closed	limited	NE	all	10	6	5	9	17	7	12	12	9	5
Scallop Dredge	closed	limited	MA	all	15	7	18	12	18	8	11	8	10	5
Scallop Dredge	closed	general	NE	all										
Scallop Dredge	closed	general	MA	all	18	7	19	19	23	18	14	16	11	5
Mid-water paired & single Trawl	all	all	NE	all	5	7	14	20	9	20	10	15	11	5
Mid-water paired & single Trawl	all	all	MA	all	16	7	17	22	15	19	14	11	11	5
Fish Pots/ Traps	all	all	NE	all										
Fish Pots/ Traps	all	all	MA	all	18	7	19	22	23	20	6	1	11	5
Purse Seine	all	all	NE	all	18	7	19	22	11	20	14	16	11	5
Purse Seine	all	all	MA	all										
Hand Line	all	all	NE	all	18	7	19	22	23	20	14	16	11	5
Hand Line	all	all	MA	all										
Scottish Seine	all	all	NE	all	14	7	12	21	23	12	9	9	11	5
Clam Quahog Dredge	all	all	NE	all										
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all										
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-3 continued. Ranking of total discards within species group (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	TURTLE, GREEN	TURTLE, LEATHERBACK	TURTLE, LOGGERHEAD	TURTLE, KEMP'S RIDLEY	TURTLE, NK	SEAL, HARP	SEAL, HOODED	SEAL, HARBOR	SEAL, GRAY	SEAL, NK
Longline	all	all	NE	all	2	3	8	*	2	3	2	4	4	*
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Otter Trawl	all	all	NE	small	2	3	8	*	2	3	2	4	4	*
Otter Trawl	all	all	NE	large	2	3	8	*	2	3	2	4	4	*
Otter Trawl	all	all	MA	small	2	3	4	*	2	3	2	4	4	*
Otter Trawl	all	all	MA	large	2	3	8	*	2	3	2	4	4	*
Scallop Trawl	open	limited	MA	all	2	3	1	*	2	3	2	4	4	*
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	*
Sink, Anchor, Drift Gillnet	all	all	NE	large	2	3	8	*	2	1	2	2	2	*
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	2	3	8	*	2	2	1	1	1	*
Sink, Anchor, Drift Gillnet	all	all	MA	small	2	2	8	*	1	3	2	4	4	*
Sink, Anchor, Drift Gillnet	all	all	MA	large	1	3	6	*	2	3	2	4	4	*
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	2	1	5	*	2	3	2	3	3	*
Scallop Dredge	open	limited	NE	all	2	3	2	*	2	3	2	4	4	*
Scallop Dredge	open	limited	MA	all	2	3	3	*	2	3	2	4	4	*
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	limited	NE	all	2	3	7	*	2	3	2	4	4	*
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	NE	all	*	*	*	*	*	*	*	*	*	*
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	*
Mid-water paired & single Trawl	all	all	NE	all	2	3	8	*	2	3	2	4	4	*
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Fish Pots/ Traps	all	all	NE	all										
Fish Pots/ Traps	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	*
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Clam Quahog Dredge	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Clam Quahog Dredge	all	all	MA	all										
Crab Pots	all	all	NE	all										
Crab Pots	all	all	MA	all										
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	*
Lobster Pots	all	all	MA	all										

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-3 continued. Ranking of total discards within species group (fish and protected species ranked separately) based on 2004 observer data.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	WHALE, PILOT, LONG-FIN	WHALE, MINKE	WHALE, NK	DOLPHIN WHITE-SIDED	DOLPHIN-COMMON	DOLPHIN-BOTTLE-NOSE	PORPOISE-HARBOR	PORPOISE/DOLP-HIN, NK	SEA BIRDS (ALL)	PILOT coverage
Longline	all	all	NE	all	4	*	*	5	3	2	4	*	12	
Longline	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Otter Trawl	all	all	NE	small	1	*	*	3	1	2	4	*	6	
Otter Trawl	all	all	NE	large	2	*	*	1	3	2	4	*	5	
Otter Trawl	all	all	MA	small	4	*	*	5	2	2	4	*	11	
Otter Trawl	all	all	MA	large	4	*	*	5	3	2	4	*	3	
Scallop Trawl	open	limited	MA	all	4	*	*	5	3	2	4	*	14	pilot
Scallop Trawl	open	general	MA	all	*	*	*	*	*	*	*	*	*	pilot
Shrimp Trawl	all	all	NE	all	*	*	*	*	*	*	*	*	*	
Shrimp Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	small	*	*	*	*	*	*	*	*	*	pilot
Sink, Anchor, Drift Gillnet	all	all	NE	large	4	*	*	4	3	2	2	*	1	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	4	*	*	5	3	1	1	*	9	
Sink, Anchor, Drift Gillnet	all	all	MA	small	4	*	*	5	3	2	4	*	8	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	large	4	*	*	5	3	2	4	*	7	pilot for fish
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	4	*	*	5	3	2	3	*	10	pilot for fish
Scallop Dredge	open	limited	NE	all	4	*	*	5	3	2	4	*	4	
Scallop Dredge	open	limited	MA	all	4	*	*	5	3	2	4	*	14	
Scallop Dredge	open	general	NE	all	*	*	*	*	*	*	*	*	*	pilot
Scallop Dredge	open	general	MA	all	*	*	*	*	*	*	*	*	*	
Scallop Dredge	closed	limited	NE	all	4	*	*	5	3	2	4	*	13	
Scallop Dredge	closed	limited	MA	all	*	*	*	*	*	*	*	*	*	
Scallop Dredge	closed	general	NE	all	*	*	*	*	*	*	*	*	*	pilot
Scallop Dredge	closed	general	MA	all	*	*	*	*	*	*	*	*	*	pilot
Mid-water paired & single Trawl	all	all	NE	all	3	*	*	2	3	2	4	*	2	
Mid-water paired & single Trawl	all	all	MA	all	*	*	*	*	*	*	*	*	*	
Fish Pots/ Traps	all	all	NE	all										pilot
Fish Pots/ Traps	all	all	MA	all										pilot
Purse Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	
Purse Seine	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Hand Line	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Hand Line	all	all	MA	all	*	*	*	*	*	*	*	*	*	pilot
Scottish Seine	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Clam Quahog Dredge	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Clam Quahog Dredge	all	all	MA	all										pilot
Crab Pots	all	all	NE	all										pilot
Crab Pots	all	all	MA	all										pilot
Lobster Pots	all	all	NE	all	*	*	*	*	*	*	*	*	*	pilot
Lobster Pots	all	all	MA	all										pilot

Gray-shaded cells indicate unlikely combinations of species/gear; * indicate no discards of these species.

Table B-4. Number of sea days needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	BLUEFISH	HERRING	SALMON	RED CRAB	SCALLOP	MACK/SQUID/BUTTERFISH	Mackereel	Illex	Loligo	Butterfish	MONKFISH
Longline	all	all	NE	all	35	35	35	35	35	35	35	35	35	35	35
Longline	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76
Otter Trawl	all	all	NE	small	1103	882	211	848	1998	249	1748	487	454	633	757
Otter Trawl	all	all	NE	large	26644	12864	730	798	1233	3159	2582	9820	3561	5259	81
Otter Trawl	all	all	MA	small	2231	1869	196	5417	1162	1125	2841	1362	1697	1160	497
Otter Trawl	all	all	MA	large	3625	883	342	342	311	242	374	753	327	394	140
Scallop Trawl	open	limited	MA	all	95	95	95	95	95	95	95	95	95	95	95
Scallop Trawl	open	general	MA	all	155	51	51	399	119	181	51	277	200	270	115
Shrimp Trawl	all	all	NE	all	42	92	42	42	353	364	42	42	42	364	22
Shrimp Trawl	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12
Sink, Anchor, Drift Gillnet	all	all	NE	large	443	486	141	2592	4357	3758	3929	5405	141	6119	408
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	267	1004	144	3266	1255	1701	1708	144	144	4216	238
Sink, Anchor, Drift Gillnet	all	all	MA	small	62	62	62	62	62	62	62	62	62	62	62
Sink, Anchor, Drift Gillnet	all	all	MA	large	105	29	29	29	29	29	29	29	29	29	29
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	131	68	68	68	301	68	68	68	68	68	104
Scallop Dredge	open	limited	NE	all	269	269	269	1596	80	1380	269	709	3260	6097	320
Scallop Dredge	open	limited	MA	all	329	329	329	8713	280	641	8713	1662	986	2528	213
Scallop Dredge	open	general	NE	all	92	92	92	92	204	176	92	176	92	92	117
Scallop Dredge	open	general	MA	all	96	96	96	96	54	293	96	96	293	96	17
Scallop Dredge	closed	limited	NE	all	3861	344	139	1473	167	1301	327	564	1531	273	429
Scallop Dredge	closed	limited	MA	all	1777	772	108	341	157	337	764	370	989	324	283
Scallop Dredge	closed	general	NE	all	24	24	24	24	24	24	24	24	24	24	24
Scallop Dredge	closed	general	MA	all	21	21	21	21	21	21	21	21	21	21	21
Mid-water paired & single Trawl	all	all	NE	all	699	747	56	56	1793	346	347	829	1786	1712	718
Mid-water paired & single Trawl	all	all	MA	all	182	453	35	35	35	167	181	165	182	182	492
Fish Pots/ Traps	all	all	NE	all	20	20	20	20	20	20	20	20	20	20	20
Fish Pots/ Traps	all	all	MA	all	40	40	40	40	40	40	40	40	40	40	103
Purse Seine	all	all	NE	all	19	219	19	19	19	206	19	206	19	19	19
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9
Hand Line	all	all	NE	all	72	72	72	72	72	72	72	72	72	72	72
Hand Line	all	all	MA	all	133	133	133	133	133	133	133	133	133	133	133
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12
Clam Quahog Dredge	all	all	NE	all	50	50	50	50	50	50	50	50	50	50	50
Clam Quahog Dredge	all	all	MA	all	84	84	84	84	84	84	84	84	84	84	84
Crab Pots	all	all	NE	all	101	101	101	101	101	101	101	101	101	101	101
Crab Pots	all	all	MA	all	28	28	28	28	28	28	28	28	28	28	28
Lobster Pots	all	all	NE	all	439	439	439	439	439	439	439	439	439	439	439
Lobster Pots	all	all	MA	all	89	89	89	89	89	89	89	89	89	89	89
Total Sea Days					43,547	23,025	4,573	27,698	15,384	17,200	25,658	24,643	17,279	31,311	6,541
Total Sea Days excluding shaded cells					35,867	19,828	0	5,547	6,049	15,522	24,114	17,398	15,185	19,567	5,528

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-4 continued. Number of sea days needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	NE MUL FISPP (LARGE-MESH)													
					Cod	Haddock	Yellowtail fltd	American plaice	Witch fltd	Winter fltd	Pollock	Redfish	White haike	Windowpane	Halibut	Ocean pout		
Longline	all	all	NE	all	27	39	36	35	35	35	35	310	35	35	35	35	76	
Longline	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	76	
Otter Trawl	all	all	NE	small	266	1861	1892	774	442	523	852	1351	1507	969	405	2129	490	
Otter Trawl	all	all	NE	large	107	321	719	505	663	219	1837	3732	630	568	440	1793	268	
Otter Trawl	all	all	MA	small	429	196	196	2763	3849	904	1172	196	2272	2580	633	196	3911	
Otter Trawl	all	all	MA	large	101	5866	342	677	342	137	272	5866	1315	5882	156	342	358	
Scallop Trawl	open	limited	MA	all	95	95	95	95	95	95	95	95	95	95	95	95	95	
Scallop Trawl	open	general	MA	all	85	51	51	505	51	255	423	51	51	399	170	51	51	
Shrimp Trawl	all	all	NE	all	20	50	171	121	38	328	29	89	56	22	17	42	349	
Shrimp Trawl	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	76	
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12	12	
Sink, Anchor, Drift Gillnet	all	all	NE	large	83	142	320	360	693	1325	724	315	607	488	1500	1532	1504	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	206	249	473	963	833	3315	1829	750	948	979	2892	1281	1245	
Sink, Anchor, Drift Gillnet	all	all	MA	small	62	62	62	62	62	62	62	62	62	62	62	62	62	
Sink, Anchor, Drift Gillnet	all	all	MA	large	19	29	29	29	29	29	29	29	29	29	19	29	29	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	68	68	68	68	68	68	68	68	68	68	68	68	68	
Scallop Dredge	open	limited	NE	all	708	1855	1611	1077	1611	728	2851	1611	269	848	633	269	1298	
Scallop Dredge	open	limited	MA	all	411	329	329	2708	3159	1660	2108	329	329	1687	715	329	6549	
Scallop Dredge	open	general	NE	all	82	120	92	85	110	146	92	92	92	92	90	92	190	
Scallop Dredge	open	general	MA	all	40	96	96	293	288	171	73	96	96	173	46	96	96	
Scallop Dredge	closed	limited	NE	all	227	1735	1311	390	3105	315	355	139	139	1576	988	439	1332	
Scallop Dredge	closed	limited	MA	all	1136	108	108	1819	333	404	1829	108	108	800	1409	108	1803	
Scallop Dredge	closed	general	NE	all	24	24	24	24	24	24	24	24	24	24	24	24	24	
Scallop Dredge	closed	general	MA	all	21	21	21	21	21	21	21	21	21	21	21	21	21	
Mid-water paired & single Trawl	all	all	NE	all	688	1451	855	56	1437	1037	1616	1217	1219	1128	56	56	56	
Mid-water paired & single Trawl	all	all	MA	all	281	35	35	35	35	547	35	35	35	176	35	35	35	
Fish Pots/ Traps	all	all	NE	all	20	20	20	20	20	20	20	20	20	20	20	20	20	
Fish Pots/ Traps	all	all	MA	all	40	40	40	40	40	40	40	40	40	40	40	40	40	
Purse Seine	all	all	NE	all	217	19	19	19	19	19	19	19	217	19	19	19	19	
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9	9	
Hand Line	all	all	NE	all	137	137	72	72	72	72	72	72	72	72	72	72	72	
Hand Line	all	all	MA	all	133	133	133	133	133	133	133	133	133	133	133	133	133	
Scottish Seine	all	all	NE	all	14	12	12	12	12	12	18	12	12	12	15	12	12	
Clam Quahog Dredge	all	all	NE	all	50	50	50	50	50	50	50	50	50	50	50	50	50	
Clam Quahog Dredge	all	all	MA	all	84	84	84	84	84	84	84	84	84	84	84	84	84	
Crab Pots	all	all	NE	all	101	101	101	101	101	101	101	101	101	101	101	101	101	
Crab Pots	all	all	MA	all	28	28	28	28	28	28	28	28	28	28	28	28	28	
Lobster Pots	all	all	NE	all	439	439	439	439	439	439	439	439	439	439	439	439	439	
Lobster Pots	all	all	MA	all	89	89	89	89	89	89	89	89	89	89	89	89	89	
Total Sea Days					6,712	16,129	10,194	14,724	18,580	13,608	17,695	17,846	11,464	19,958	11,769	10,382	21,169	
Total Sea Days excluding shaded cells					6,450	15,330	5,143	13,481	9,033	9,941	16,768	7,934	5,380	15,652	10,603	7,758	20,321	

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-4 continued. Number of sea days needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Species												
					NE MULTISPP (SMALL-MESH)	Silver hake	Offshore hake	Red hake	SKATE	DOG FISH	FLUKE/SCUPEL-BLK SEA BASS	Fluke	Scup	Black sea bass	SURF CLAM-OCEAN QUAHOG	TILEFISH	
Longline	all	all	NE	all	185	35	35	185	89	99	35	35	35	35	35	35	
Longline	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	
Otter Trawl	all	all	NE	small	269	233	5131	762	2024	492	455	365	1344	2115	3822	441	
Otter Trawl	all	all	NE	large	341	530	1053	1255	316	614	1034	1091	7275	6224	15593	2692	
Otter Trawl	all	all	MA	small	944	1324	1690	1198	202	532	584	314	1120	952	836	3057	
Otter Trawl	all	all	MA	large	998	321	342	3401	70	481	98	114	201	643	584	342	
Scallop Trawl	open	limited	MA	all	95	95	95	95	95	95	95	95	95	95	95	95	
Scallop Trawl	open	general	MA	all	292	297	51	155	80	443	408	75	102	396	51	51	
Shrimp Trawl	all	all	NE	all	123	128	42	115	247	349	42	42	42	42	42	42	
Shrimp Trawl	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12	
Sink, Anchor, Drift Gillnet	all	all	NE	large	313	514	141	439	482	109	3767	4020	141	6294	141	141	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	2059	345	144	2902	109	214	417	418	4204	144	144	502	
Sink, Anchor, Drift Gillnet	all	all	MA	small	62	62	62	62	62	62	62	62	62	62	62	62	
Sink, Anchor, Drift Gillnet	all	all	MA	large	29	29	29	29	99	96	29	29	29	29	29	29	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	68	68	68	68	55	58	120	120	68	68	68	68	
Scallop Dredge	open	limited	NE	all	534	1706	3605	390	177	807	649	693	325	1172	478	269	
Scallop Dredge	open	limited	MA	all	3080	3657	3443	1081	114	371	465	512	2642	2077	2958	329	
Scallop Dredge	open	general	NE	all	135	204	92	148	120	120	92	92	92	92	190	92	
Scallop Dredge	open	general	MA	all	96	90	96	288	17	124	88	88	96	96	271	96	
Scallop Dredge	closed	limited	NE	all	1180	1214	1618	1434	145	857	703	712	452	252	375	139	
Scallop Dredge	closed	limited	MA	all	287	339	108	298	88	567	481	485	1568	378	334	108	
Scallop Dredge	closed	general	NE	all	24	24	24	24	24	24	24	24	24	24	24	24	
Scallop Dredge	closed	general	MA	all	21	21	21	21	21	21	21	21	21	21	21	21	
Mid-water paired & single Trawl	all	all	NE	all	1218	1226	56	826	1034	316	697	56	797	1134	56	56	
Mid-water paired & single Trawl	all	all	MA	all	182	182	35	182	35	43	557	544	35	563	35	35	
Fish Pots/ Traps	all	all	NE	all	20	20	20	20	20	20	20	20	20	20	20	20	
Fish Pots/ Traps	all	all	MA	all	40	40	40	40	40	40	40	40	73	40	40	40	
Purse Seine	all	all	NE	all	19	19	19	19	19	217	19	19	19	19	19	19	
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9	
Hand Line	all	all	NE	all	72	72	72	72	72	72	72	72	72	72	72	72	
Hand Line	all	all	MA	all	133	133	133	133	133	133	133	133	133	133	133	133	
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	30	30	30	30	12	12	
Clam Quahog Dredge	all	all	NE	all	50	50	50	50	50	50	50	50	50	50	50	50	
Clam Quahog Dredge	all	all	MA	all	84	84	84	84	84	84	84	84	84	84	84	84	
Crab Pots	all	all	NE	all	101	101	101	101	101	101	101	101	101	101	101	101	
Crab Pots	all	all	MA	all	28	28	28	28	28	28	28	28	28	28	28	28	
Lobster Pots	all	all	NE	all	439	439	439	439	439	439	439	439	439	439	439	439	
Lobster Pots	all	all	MA	all	89	89	89	89	89	89	89	89	89	89	89	89	
Total Sea Days					13,792	13,902	19,241	16,616	6,965	8,351	12,200	11,286	22,079	24,187	27,502	9,984	
Total Sea Days excluding shaded cells					12,562	12,637	6,272	15,136	4,901	6,943	9,850	9,591	10,416	10,673	133	6,703	

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-4 continued. Number of sea days needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Species													
					TURTLES	TURTLE, GREEN	TURTLE, LEATHERBACK	TURTLE, LOGGERHEAD	TURTLE, KEMPS RIDLEY	TURTLE, NK	SEALS	SEAL, HARP	SEAL, HOODED	SEAL, HARBOR	SEAL, GRAY	SEAL, NK		
Longline	all	all	NE	all	35	35	35	35	35	35	35	35	35	35	35	35	35	
Longline	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	76	
Otter Trawl	all	all	NE	small	211	211	211	211	211	211	211	211	211	211	211	211	211	
Otter Trawl	all	all	NE	large	730	730	730	730	730	730	730	730	730	730	730	730	730	
Otter Trawl	all	all	MA	small	1229	196	196	1229	196	196	196	196	196	196	196	196	196	
Otter Trawl	all	all	MA	large	342	342	342	342	342	342	342	342	342	342	342	342	342	
Scallop Trawl	open	limited	MA	all	95	95	95	95	95	95	95	95	95	95	95	95	95	
Scallop Trawl	open	general	MA	all	51	51	51	51	51	51	51	51	51	51	51	51	51	
Shrimp Trawl	all	all	NE	all	42	42	42	42	42	42	42	42	42	42	42	42	42	
Shrimp Trawl	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76	76	
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12	12	
Sink, Anchor, Drift Gillnet	all	all	NE	large	141	141	141	141	141	141	531	1007	141	889	2518	141		
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	144	144	144	144	144	144	470	1694	3812	973	731	144		
Sink, Anchor, Drift Gillnet	all	all	MA	small	1259	62	1841	62	62	1757	62	62	62	62	62	62	62	
Sink, Anchor, Drift Gillnet	all	all	MA	large	653	913	29	913	29	29	29	29	29	29	29	29	29	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	468	68	799	764	68	68	804	68	68	1175	1272	68		
Scallop Dredge	open	limited	NE	all	1261	269	269	1261	269	269	269	269	269	269	269	269	269	
Scallop Dredge	open	limited	MA	all	3956	329	329	3956	329	329	329	329	329	329	329	329	329	
Scallop Dredge	open	general	NE	all	92	92	92	92	92	92	92	92	92	92	92	92	92	
Scallop Dredge	open	general	MA	all	96	96	96	96	96	96	96	96	96	96	96	96	96	
Scallop Dredge	closed	limited	NE	all	414	139	139	414	139	139	139	139	139	139	139	139	139	
Scallop Dredge	closed	limited	MA	all	108	108	108	108	108	108	108	108	108	108	108	108	108	
Scallop Dredge	closed	general	NE	all	24	24	24	24	24	24	24	24	24	24	24	24	24	
Scallop Dredge	closed	general	MA	all	21	21	21	21	21	21	21	21	21	21	21	21	21	
Mid-water paired & single Trawl	all	all	NE	all	56	56	56	56	56	56	56	56	56	56	56	56	56	
Mid-water paired & single Trawl	all	all	MA	all	35	35	35	35	35	35	35	35	35	35	35	35	35	
Fish Pots/ Traps	all	all	NE	all	20	20	20	20	20	20	20	20	20	20	20	20	20	
Fish Pots/ Traps	all	all	MA	all	40	40	40	40	40	40	40	40	40	40	40	40	40	
Purse Seine	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	19	19	
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9	9	
Hand Line	all	all	NE	all	72	72	72	72	72	72	72	72	72	72	72	72	72	
Hand Line	all	all	MA	all	133	133	133	133	133	133	133	133	133	133	133	133	133	
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12	
Clam Quahog Dredge	all	all	NE	all	50	50	50	50	50	50	50	50	50	50	50	50	50	
Clam Quahog Dredge	all	all	MA	all	84	84	84	84	84	84	84	84	84	84	84	84	84	
Crab Pots	all	all	NE	all	101	101	101	101	101	101	101	101	101	101	101	101	101	
Crab Pots	all	all	MA	all	28	28	28	28	28	28	28	28	28	28	28	28	28	
Lobster Pots	all	all	NE	all	439	439	439	439	439	439	439	439	439	439	439	439	439	
Lobster Pots	all	all	MA	all	89	89	89	89	89	89	89	89	89	89	89	89	89	
Total Sea Days					12,721	5,457	7,083	12,079	4,573	6,269	6,025	6,990	8,241	7,257	8,741	4,573		
Total Sea Days excluding shaded cells					12,721	3,082	6,142	12,079	3,632	6,269	4,742	5,706	5,319	5,436	6,920	3,290		

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-4 continued. Number of sea days needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	WHALES	WHALE, PILOT, LONG-FIN	WHALE, MINKE	WHALE, NK	DOLPHINS- PORPOISE	DOLPHIN WHITE-SIDED	DOLPHIN COMMON	DOLPHIN BOTTLE-NOSE	PORPOISE HARBOR	PORPOISE/ DOLPHIN, NK	SEA BIRDS (ALL)	ALL SPECIES
Longline	all	all	NE	all	35	35	35	35	35	35	35	35	35	35	267	57
Longline	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	76
Otter Trawl	all	all	NE	small	3082	3082	211	211	2265	3099	2594	211	211	211	1870	183
Otter Trawl	all	all	NE	large	10526	10526	730	730	2111	2111	730	730	730	730	3237	159
Otter Trawl	all	all	MA	small	196	196	196	196	1164	196	1164	196	196	196	1880	250
Otter Trawl	all	all	MA	large	342	342	342	342	342	342	342	342	342	342	727	55
Scallop Trawl	open	limited	MA	all	95	95	95	95	95	95	95	95	95	95	95	95
Scallop Trawl	open	general	MA	all	51	51	51	51	51	51	51	51	51	51	51	38
Shrimp Trawl	all	all	NE	all	42	42	42	42	42	42	42	42	42	42	42	39
Shrimp Trawl	all	all	MA	all	76	76	76	76	76	76	76	76	76	76	76	55
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12
Sink, Anchor, Drift Gillnet	all	all	NE	large	141	141	141	141	1398	3642	141	141	1557	141	1306	82
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	144	144	144	144	806	144	144	3816	871	144	2661	59
Sink, Anchor, Drift Gillnet	all	all	MA	small	62	62	62	62	62	62	62	62	62	62	880	62
Sink, Anchor, Drift Gillnet	all	all	MA	large	29	29	29	29	29	29	29	29	29	29	311	95
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	68	68	68	68	1272	68	68	68	1272	68	806	51
Scallop Dredge	open	limited	NE	all	269	269	269	269	269	269	269	269	269	269	3194	123
Scallop Dredge	open	limited	MA	all	329	329	329	329	329	329	329	329	329	329	329	89
Scallop Dredge	open	general	NE	all	92	92	92	92	92	92	92	92	92	92	92	88
Scallop Dredge	open	general	MA	all	96	96	96	96	96	96	96	96	96	96	96	14
Scallop Dredge	closed	limited	NE	all	139	139	139	139	139	139	139	139	139	139	407	130
Scallop Dredge	closed	limited	MA	all	108	108	108	108	108	108	108	108	108	108	108	61
Scallop Dredge	closed	general	NE	all	24	24	24	24	24	24	24	24	24	24	24	24
Scallop Dredge	closed	general	MA	all	21	21	21	21	21	21	21	21	21	21	21	21
Mid-water paired & single Trawl	all	all	NE	all	1606	1606	56	56	1464	1464	56	56	56	56	808	193
Mid-water paired & single Trawl	all	all	MA	all	35	35	35	35	35	35	35	35	35	35	35	111
Fish Pots/ Traps	all	all	NE	all	20	20	20	20	20	20	20	20	20	20	20	20
Fish Pots/ Traps	all	all	MA	all	40	40	40	40	40	40	40	40	40	40	40	37
Purse Seine	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	143
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9
Hand Line	all	all	NE	all	72	72	72	72	72	72	72	72	72	72	72	137
Hand Line	all	all	MA	all	133	133	133	133	133	133	133	133	133	133	133	133
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	20
Clam Quahog Dredge	all	all	NE	all	50	50	50	50	50	50	50	50	50	50	50	50
Clam Quahog Dredge	all	all	MA	all	84	84	84	84	84	84	84	84	84	84	84	84
Crab Pots	all	all	NE	all	101	101	101	101	101	101	101	101	101	101	101	101
Crab Pots	all	all	MA	all	28	28	28	28	28	28	28	28	28	28	28	28
Lobster Pots	all	all	NE	all	439	439	439	439	439	439	439	439	439	439	439	439
Lobster Pots	all	all	MA	all	89	89	89	89	89	89	89	89	89	89	89	89
Total Sea Days					18,791	18,791	4,573	4,573	13,507	13,752	7,923	8,245	7,920	4,573	20,503	3,513
Total Sea Days excluding shaded cells					17,714	17,714	2,593	3,497	13,507	13,752	7,923	7,708	7,829	4,573	20,503	3,513

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-5. Number of trips needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	BLUEFISH	HERRING	SALMON	RED CRAB	SCALLOP	MACK-/SQUID-/ BUTTERFISH	Mackerel	Illex	Loligo	Butterfish	MONKFISH
Longline	all	all	NE	all	26	26	26	26	26	26	26	26	26	26	26
Longline	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12
Otter Trawl	all	all	NE	small	364	291	70	280	659	82	576	161	150	209	250
Otter Trawl	all	all	NE	large	11227	5420	304	336	520	1331	1088	4138	1501	2216	34
Otter Trawl	all	all	MA	small	1189	995	104	2885	619	599	1513	725	904	618	265
Otter Trawl	all	all	MA	large	1879	458	177	177	161	125	194	390	170	204	72
Scallop Trawl	open	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12
Scallop Trawl	open	general	MA	all	72	25	25	196	56	85	25	132	94	129	54
Shrimp Trawl	all	all	NE	all	42	91	42	42	350	361	42	42	42	361	22
Shrimp Trawl	all	all	MA	all	13	13	13	13	13	13	13	13	13	13	13
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12
Sink, Anchor, Drift Gillnet	all	all	NE	large	327	359	104	1913	3216	2774	2900	3990	104	4517	301
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	175	657	94	2139	822	1114	1119	94	94	2760	156
Sink, Anchor, Drift Gillnet	all	all	MA	small	58	58	58	58	58	58	58	58	58	58	58
Sink, Anchor, Drift Gillnet	all	all	MA	large	100	27	27	27	27	27	27	27	27	27	27
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	94	51	51	51	211	51	51	51	51	51	76
Scallop Dredge	open	limited	NE	all	25	25	25	146	7	126	25	65	298	558	29
Scallop Dredge	open	limited	MA	all	36	36	36	966	31	71	966	184	109	280	24
Scallop Dredge	open	general	NE	all	71	71	71	71	149	130	71	130	71	71	89
Scallop Dredge	open	general	MA	all	69	69	69	69	39	210	69	69	210	69	12
Scallop Dredge	closed	limited	NE	all	449	40	15	171	19	151	38	66	178	32	50
Scallop Dredge	closed	limited	MA	all	194	84	12	37	17	37	83	40	108	35	31
Scallop Dredge	closed	general	NE	all	12	12	12	12	12	12	12	12	12	12	12
Scallop Dredge	closed	general	MA	all	15	15	15	15	15	15	15	15	15	15	15
Mid-water paired & single Trawl	all	all	NE	all	266	285	21	21	683	132	132	316	681	652	274
Mid-water paired & single Trawl	all	all	MA	all	52	130	12	12	12	48	52	47	52	52	141
Fish Pots/ Traps	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19
Fish Pots/ Traps	all	all	MA	all	37	37	37	37	37	37	37	37	37	37	97
Purse Seine	all	all	NE	all	10	108	10	10	10	102	10	102	10	10	10
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9
Hand Line	all	all	NE	all	68	68	68	68	68	68	68	68	68	68	68
Hand Line	all	all	MA	all	126	126	126	126	126	126	126	126	126	126	126
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12
Clam Quahog Dredge	all	all	NE	all	69	69	69	69	69	69	69	69	69	69	69
Clam Quahog Dredge	all	all	MA	all	69	69	69	69	69	69	69	69	69	69	69
Crab Pots	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12
Crab Pots	all	all	MA	all	27	27	27	27	27	27	27	27	27	27	27
Lobster Pots	all	all	NE	all	353	353	353	353	353	353	353	353	353	353	353
Lobster Pots	all	all	MA	all	75	75	75	75	75	75	75	75	75	75	75
Total Trips					17,678	10,260	2,306	10,588	8,647	8,594	10,019	11,808	5,891	13,889	3,015
Total Trips excluding shaded cells					15,925	9,034	0	1,539	2,468	7,333	8,922	6,413	4,300	5,539	2,219

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-5 continued. Number of trips needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	NE MUL FISPP (LARGE-MESH)													
					Cod	Haddock	Yellowtail fltd	American plaice	Witch fltd	Winter fltd	Pollock	Redfish	White hake	Windowpane	Halibut	Ocean pout		
Longline	all	all	NE	all	21	30	28	26	26	26	26	242	26	26	26	26	59	
Longline	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12	
Otter Trawl	all	all	NE	small	88	614	624	255	146	173	281	446	497	319	133	702	161	
Otter Trawl	all	all	NE	large	45	135	303	213	279	92	774	1572	266	239	185	755	113	
Otter Trawl	all	all	MA	small	229	104	104	1472	2050	481	624	104	1210	1374	337	104	2083	
Otter Trawl	all	all	MA	large	52	3040	177	351	177	71	141	3040	681	3048	81	177	185	
Scallop Trawl	open	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12	
Scallop Trawl	open	general	MA	all	41	25	25	244	25	125	207	25	25	196	81	25	25	
Shrimp Trawl	all	all	NE	all	20	49	169	120	37	325	29	89	56	22	17	42	346	
Shrimp Trawl	all	all	MA	all	13	13	13	13	13	13	13	13	13	13	13	13	13	
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12	12	
Sink, Anchor, Drift Gillnet	all	all	NE	large	61	105	236	266	512	978	534	232	448	360	1107	1131	1110	
Sink, Anchor, Drift Gillnet	all	all	NE	xlg	135	163	310	631	545	2171	1197	491	621	641	1893	839	815	
Sink, Anchor, Drift Gillnet	all	all	MA	small	58	58	58	58	58	58	58	58	58	58	58	58	58	
Sink, Anchor, Drift Gillnet	all	all	MA	large	17	27	27	27	27	27	27	27	27	27	17	27	27	
Sink, Anchor, Drift Gillnet	all	all	MA	xlg	51	51	51	51	51	51	51	51	51	51	51	51	51	
Scallop Dredge	open	limited	NE	all	65	170	147	99	147	67	261	147	25	78	58	25	119	
Scallop Dredge	open	limited	MA	all	46	36	36	300	350	184	234	36	36	187	79	36	726	
Scallop Dredge	open	general	NE	all	64	91	71	66	84	109	71	71	71	71	70	71	140	
Scallop Dredge	open	general	MA	all	29	69	69	210	206	123	53	69	69	124	33	69	69	
Scallop Dredge	closed	limited	NE	all	26	202	152	45	361	37	41	15	15	183	115	51	155	
Scallop Dredge	closed	limited	MA	all	124	12	12	199	36	44	200	12	12	87	154	12	197	
Scallop Dredge	closed	general	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12	
Scallop Dredge	closed	general	MA	all	15	15	15	15	15	15	15	15	15	15	15	15	15	
Mid-water paired & single Trawl	all	all	NE	all	262	553	326	21	548	395	616	464	465	430	21	21	21	
Mid-water paired & single Trawl	all	all	MA	all	81	12	12	12	12	157	12	12	12	50	12	12	12	
Fish Pots/ Traps	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	19	19	
Fish Pots/ Traps	all	all	MA	all	37	37	37	37	37	37	37	37	37	37	37	37	37	
Purse Seine	all	all	NE	all	107	10	10	10	10	10	10	10	107	10	10	10	10	
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9	9	
Hand Line	all	all	NE	all	129	129	68	68	68	68	68	68	68	68	68	68	68	
Hand Line	all	all	MA	all	126	126	126	126	126	126	126	126	126	126	126	126	126	
Scottish Seine	all	all	NE	all	14	12	12	12	12	12	18	12	12	12	15	12	12	
Clam Quahog Dredge	all	all	NE	all	69	69	69	69	69	69	69	69	69	69	69	69	69	
Clam Quahog Dredge	all	all	MA	all	69	69	69	69	69	69	69	69	69	69	69	69	69	
Crab Pots	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12	
Crab Pots	all	all	MA	all	27	27	27	27	27	27	27	27	27	27	27	27	27	
Lobster Pots	all	all	NE	all	353	353	353	353	353	353	353	353	353	353	353	353	353	
Lobster Pots	all	all	MA	all	75	75	75	75	75	75	75	75	75	75	75	75	75	
Total Trips					2,641	6,572	3,902	5,630	6,641	6,657	6,407	8,167	5,730	8,536	5,495	5,198	7,437	
Total Trips excluding shaded cells					2,464	6,143	2,403	4,676	2,717	5,239	5,685	3,669	2,618	7,001	4,602	3,710	6,881	

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-5 continued. Number of trips needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups														
					NE MULTI-SPP (SMALL-MESH)	Silver hake	Offshore hake	Red hake	SKATE	DOG FISH	FLUKE/SCUP/-BLK SEA BASS	Fluke	Scup	Black sea bass	SURF CLAM/OCEAN QUAHOG	TILEFISH		
Longline	all	all	NE	all	144	26	26	144	69	78	26	26	26	26	26	26		
Longline	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	12		
Otter Trawl	all	all	NE	small	89	77	1692	251	668	162	150	120	443	698	1260	146		
Otter Trawl	all	all	NE	large	143	223	444	529	133	259	436	460	3065	2623	6570	1134		
Otter Trawl	all	all	MA	small	503	705	900	638	108	283	311	167	596	507	445	1628		
Otter Trawl	all	all	MA	large	517	166	177	1762	36	249	51	59	104	333	303	177		
Scallop Trawl	open	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	12		
Scallop Trawl	open	general	MA	all	142	144	25	72	39	216	200	36	48	195	25	25		
Shrimp Trawl	all	all	NE	all	122	127	42	114	245	346	42	42	42	42	42	42		
Shrimp Trawl	all	all	MA	all	13	13	13	13	13	13	13	13	13	13	13	13		
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12		
Sink, Anchor, Drift Gillnet	all	all	NE	large	231	380	104	324	356	81	2780	2967	104	4646	104	104		
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	1348	226	94	1900	71	140	273	273	2753	94	94	329		
Sink, Anchor, Drift Gillnet	all	all	MA	small	58	58	58	58	58	58	58	58	58	58	58	58		
Sink, Anchor, Drift Gillnet	all	all	MA	large	27	27	27	27	95	91	27	27	27	27	27	27		
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	51	51	51	51	42	44	87	87	51	51	51	51		
Scallop Dredge	open	limited	NE	all	49	156	330	36	16	74	59	63	30	107	44	25		
Scallop Dredge	open	limited	MA	all	342	406	382	120	13	41	52	57	293	230	328	36		
Scallop Dredge	open	general	NE	all	102	149	71	111	91	91	71	71	71	71	140	71		
Scallop Dredge	open	general	MA	all	69	65	69	206	12	89	63	63	69	69	194	69		
Scallop Dredge	closed	limited	NE	all	137	141	188	167	17	100	82	83	52	29	44	15		
Scallop Dredge	closed	limited	MA	all	31	37	12	33	10	62	53	53	171	41	37	12		
Scallop Dredge	closed	general	NE	all	12	12	12	12	12	12	12	12	12	12	12	12		
Scallop Dredge	closed	general	MA	all	15	15	15	15	15	15	15	15	15	15	15	15		
Mid-water paired & single Trawl	all	all	NE	all	464	467	21	315	394	121	266	21	304	432	21	21		
Mid-water paired & single Trawl	all	all	MA	all	52	52	12	52	12	12	160	156	12	161	12	12		
Fish Pots/ Traps	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	19		
Fish Pots/ Traps	all	all	MA	all	37	37	37	37	37	37	37	37	69	37	37	37		
Purse Seine	all	all	NE	all	10	10	10	10	10	107	10	10	10	10	10	10		
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9		
Hand Line	all	all	NE	all	68	68	68	68	68	68	68	68	68	68	68	68		
Hand Line	all	all	MA	all	126	126	126	126	126	126	126	126	126	126	126	126		
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	30	30	30	30	12	12		
Clam Quahog Dredge	all	all	NE	all	69	69	69	69	69	69	69	69	69	69	69	69		
Clam Quahog Dredge	all	all	MA	all	69	69	69	69	69	69	69	69	69	69	69	69		
Crab Pots	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12		
Crab Pots	all	all	MA	all	27	27	27	27	27	27	27	27	27	27	27	27		
Lobster Pots	all	all	NE	all	353	353	353	353	353	353	353	353	353	353	353	353		
Lobster Pots	all	all	MA	all	75	75	75	75	75	75	75	75	75	75	75	75		
Total Trips					5,584	4,647	5,688	7,873	3,447	3,658	6,227	5,872	9,332	11,423	10,788	4,971		
Total Trips excluding shaded cells					4,637	3,673	2,211	6,842	2,243	2,511	4,981	4,874	4,476	4,543	139	3,180		

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-5 continued. Number of trips needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	Species												
					TURTLES	TURTLE, GREEN	TURTLE, LEATHERBACK	TURTLE, LOGGERHEAD	TURTLE, KEMPS RIDLEY	TURTLE, NK	SEALS	SEAL, HARP	SEAL, HOODED	SEAL, HARBOR	SEAL, GRAY	SEAL, NK	
Longline	all	all	NE	all	26	26	26	26	26	26	26	26	26	26	26	26	26
Longline	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Otter Trawl	all	all	NE	small	70	70	70	70	70	70	70	70	70	70	70	70	70
Otter Trawl	all	all	NE	large	304	304	304	304	304	304	304	304	304	304	304	304	304
Otter Trawl	all	all	MA	small	654	104	104	654	104	104	104	104	104	104	104	104	104
Otter Trawl	all	all	MA	large	177	177	177	177	177	177	177	177	177	177	177	177	177
Scallop Trawl	open	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Scallop Trawl	open	general	MA	all	25	25	25	25	25	25	25	25	25	25	25	25	25
Shrimp Trawl	all	all	NE	all	42	42	42	42	42	42	42	42	42	42	42	42	42
Shrimp Trawl	all	all	MA	all	13	13	13	13	13	13	13	13	13	13	13	13	13
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12	12
Sink, Anchor, Drift Gillnet	all	all	NE	large	104	104	104	104	104	104	392	743	104	656	1859	104	104
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	94	94	94	94	94	94	308	1109	2496	637	479	94	94
Sink, Anchor, Drift Gillnet	all	all	MA	small	1195	58	1747	58	58	1668	58	58	58	58	58	58	58
Sink, Anchor, Drift Gillnet	all	all	MA	large	604	845	27	845	27	27	27	27	27	27	27	27	27
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	351	51	600	573	51	51	604	51	51	882	955	51	51
Scallop Dredge	open	limited	NE	all	115	25	25	115	25	25	25	25	25	25	25	25	25
Scallop Dredge	open	limited	MA	all	439	36	36	439	36	36	36	36	36	36	36	36	36
Scallop Dredge	open	general	NE	all	71	71	71	71	71	71	71	71	71	71	71	71	71
Scallop Dredge	open	general	MA	all	69	69	69	69	69	69	69	69	69	69	69	69	69
Scallop Dredge	closed	limited	NE	all	48	15	15	48	15	15	15	15	15	15	15	15	15
Scallop Dredge	closed	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Scallop Dredge	closed	general	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Scallop Dredge	closed	general	MA	all	15	15	15	15	15	15	15	15	15	15	15	15	15
Mid-water paired & single Trawl	all	all	NE	all	21	21	21	21	21	21	21	21	21	21	21	21	21
Mid-water paired & single Trawl	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Fish Pots/ Traps	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	19	19
Fish Pots/ Traps	all	all	MA	all	37	37	37	37	37	37	37	37	37	37	37	37	37
Purse Seine	all	all	NE	all	10	10	10	10	10	10	10	10	10	10	10	10	10
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9	9
Hand Line	all	all	NE	all	68	68	68	68	68	68	68	68	68	68	68	68	68
Hand Line	all	all	MA	all	126	126	126	126	126	126	126	126	126	126	126	126	126
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Clam Quahog Dredge	all	all	NE	all	69	69	69	69	69	69	69	69	69	69	69	69	69
Clam Quahog Dredge	all	all	MA	all	69	69	69	69	69	69	69	69	69	69	69	69	69
Crab Pots	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12	12
Crab Pots	all	all	MA	all	27	27	27	27	27	27	27	27	27	27	27	27	27
Lobster Pots	all	all	NE	all	353	353	353	353	353	353	353	353	353	353	353	353	353
Lobster Pots	all	all	MA	all	75	75	75	75	75	75	75	75	75	75	75	75	75
Total Trips					5,397	3,124	4,543	4,722	2,306	3,916	3,360	3,961	4,708	4,232	5,349	2,306	2,306
Total Trips excluding shaded cells					5,397	2,096	4,169	4,722	1,933	3,916	3,023	3,623	3,579	3,613	4,730	1,969	1,969

Gray-shaded cells indicate unlikely combinations of species/gear.

Table B-5 continued. Number of trips needed to achieve a CV of 30 percent based on the total composite discards.

Gear Type	Access Area (Open-Closed)	Trip Category (General/Limited)	Region	mesh groups	WHALES	WHALE, PILOT, LONG-FIN	WHALE, MINKE	WHALE, NK	DOLPHINS- PORPOISE	DOLPHIN WHITE- SIDED	DOLPHIN- COMMON	DOLPHIN- BOTTLE- NOSE	PORPOISE, HARBOR	PORPOISE/ DOLPHIN, NK	SEA BIRDS (ALL)	ALL SPECIES
Longline	all	all	NE	all	26	26	26	26	26	26	26	26	26	26	208	44
Longline	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	12
Otter Trawl	all	all	NE	small	1016	1016	70	70	747	1022	855	70	70	70	617	60
Otter Trawl	all	all	NE	large	4435	4435	304	304	890	890	304	304	304	304	1364	67
Otter Trawl	all	all	MA	small	104	104	104	104	620	104	620	104	104	104	1001	133
Otter Trawl	all	all	MA	large	177	177	177	177	177	177	177	177	177	177	377	29
Scallop Trawl	open	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	12
Scallop Trawl	open	general	MA	all	25	25	25	25	25	25	25	25	25	25	25	18
Shrimp Trawl	all	all	NE	all	42	42	42	42	42	42	42	42	42	42	42	38
Shrimp Trawl	all	all	MA	all	13	13	13	13	13	13	13	13	13	13	13	9
Sink, Anchor, Drift Gillnet	all	all	NE	small	12	12	12	12	12	12	12	12	12	12	12	12
Sink, Anchor, Drift Gillnet	all	all	NE	large	104	104	104	104	1032	2688	104	104	1149	104	964	61
Sink, Anchor, Drift Gillnet	all	all	NE	xlq	94	94	94	94	528	94	94	2499	570	94	1742	38
Sink, Anchor, Drift Gillnet	all	all	MA	small	58	58	58	58	58	58	58	58	58	58	835	58
Sink, Anchor, Drift Gillnet	all	all	MA	large	27	27	27	27	27	27	27	27	27	27	288	91
Sink, Anchor, Drift Gillnet	all	all	MA	xlq	51	51	51	51	955	51	51	51	955	51	605	39
Scallop Dredge	open	limited	NE	all	25	25	25	25	25	25	25	25	25	25	292	11
Scallop Dredge	open	limited	MA	all	36	36	36	36	36	36	36	36	36	36	36	10
Scallop Dredge	open	general	NE	all	71	71	71	71	71	71	71	71	71	71	71	69
Scallop Dredge	open	general	MA	all	69	69	69	69	69	69	69	69	69	69	69	10
Scallop Dredge	closed	limited	NE	all	15	15	15	15	15	15	15	15	15	15	47	15
Scallop Dredge	closed	limited	MA	all	12	12	12	12	12	12	12	12	12	12	12	7
Scallop Dredge	closed	general	NE	all	12	12	12	12	12	12	12	12	12	12	12	12
Scallop Dredge	closed	general	MA	all	15	15	15	15	15	15	15	15	15	15	15	15
Mid-water paired & single Trawl	all	all	NE	all	612	612	21	21	558	558	21	21	21	21	308	73
Mid-water paired & single Trawl	all	all	MA	all	12	12	12	12	12	12	12	12	12	12	12	32
Fish Pots/ Traps	all	all	NE	all	19	19	19	19	19	19	19	19	19	19	19	19
Fish Pots/ Traps	all	all	MA	all	37	37	37	37	37	37	37	37	37	37	37	34
Purse Seine	all	all	NE	all	10	10	10	10	10	10	10	10	10	10	10	71
Purse Seine	all	all	MA	all	9	9	9	9	9	9	9	9	9	9	9	9
Hand Line	all	all	NE	all	68	68	68	68	68	68	68	68	68	68	68	129
Hand Line	all	all	MA	all	126	126	126	126	126	126	126	126	126	126	126	126
Scottish Seine	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	20
Clam Quahog Dredge	all	all	NE	all	69	69	69	69	69	69	69	69	69	69	69	69
Clam Quahog Dredge	all	all	MA	all	69	69	69	69	69	69	69	69	69	69	69	69
Crab Pots	all	all	NE	all	12	12	12	12	12	12	12	12	12	12	12	12
Crab Pots	all	all	MA	all	27	27	27	27	27	27	27	27	27	27	27	27
Lobster Pots	all	all	NE	all	353	353	353	353	353	353	353	353	353	353	353	353
Lobster Pots	all	all	MA	all	75	75	75	75	75	75	75	75	75	75	75	75

Total Trips 7,975 7,975 2,306 2,306 6,887 6,965 3,607 4,711 4,731 2,306 9,877 1,992
 Total Trips excluding shaded cells 7,720 7,720 1,550 2,051 6,887 6,965 3,607 4,429 4,698 2,306 9,877 1,992

Gray-shaded cells indicate unlikely combinations of species/gear.