

Northeast Multispecies Fishery

Final Year-End Results for Fishing Year 2011

- Tables 1 through 9: Year-end total groundfish caught, landed, and discard estimates
- Tables 25 through 28: U.S./Canada total groundfish caught
- Table 29: FY 2010 & 2011 GOM cod and haddock recreational catch estimates

October 1, 2012

Table 1: FY 2011 Northeast Multispecies Catch (mt)

Final

Stock	Components with ACLs and sub-ACLs; (with accountability measures (AMs))							sub-components: No AMs	
	Total Groundfish	Groundfish Fishery	Sector	Common Pool	Recreational*	Midwater Trawl Herring Fishery**	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
GB cod	3,406.0	3,276.7	3,215.3	61.5				38.9	90.3
GOM cod	6,347.3	6,101.8	4,368.0	93.4	1,640.3			216.4	29.0
GB Haddock	4,185.2	3,840.5	3,828.8	11.7		101.8		3.9	239.0
GOM Haddock	737.5	724.1	483.7	1.9	238.5	0.2		4.9	8.3
GB Yellowtail Flounder	1,105.9	990.0	988.0	2.0			83.9	0.0	32.0
SNE/MA Yellowtail Flounder	503.6	376.2	364.0	12.2			110.9	1.1	15.4
CC/GOM Yellowtail Flounder	853.1	806.5	795.1	11.4				38.5	8.1
Plaice	1,662.1	1,636.1	1,631.6	4.5				12.1	14.0
Witch Flounder	1,177.4	997.1	992.9	4.2				22.5	157.7
GB Winter Flounder	1,980.5	1,925.4	1,924.2	1.1				0.0	55.2
GOM Winter Flounder	287.3	160.8	158.2	2.6				113.3	13.2
SNE/MA Winter Flounder	295.7	93.9	86.9	7.0				40.0	161.9
Redfish	2,722.6	2,706.7	2,703.2	3.6				3.6	12.2
White Hake	3,038.2	3,028.5	3,014.4	14.1				2.6	7.1
Pollock	9,064.5	7,612.4	7,543.1	69.2				694.0	758.1
Northern Windowpane	190.6	156.5	156.2	0.3				0.0	34.2
Southern Windowpane	528.6	111.5	83.0	28.5				16.6	400.5
Ocean Pout	88.5	60.7	56.3	4.4				0.0	27.8
Halibut	52.0	42.6	41.4	1.2				7.1	2.4
Wolffish	33.0	32.9	32.2	0.7				0.0	0.1

*Recreational estimates based on Marine Recreational Information Program (MRIP) data.

**Landings extrapolated from observer data.

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Source: NMFS Northeast Regional Office

October 1, 2012, run date of June 28, 2012

Any value for a non-allocated species may include landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories. These are SNE winter flounder, northern windowpane, southern windowpane, ocean pout, halibut, and wolffish.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 2: FY 2011 Northeast Multispecies Landings (mt)

Final

Stock	Components with ACLs and sub-ACLs; (with accountability measures (AMs))							sub-components: No AMs	
	Total Groundfish	Groundfish Fishery	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery**	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
GB cod	3,250.5	3,129.1	3,071.0	58.1				38.9	82.5
GOM cod	6,158.2	5,932.7	4,222.6	69.8	1,640.3			216.4	9.1
GB Haddock	3,883.1	3,758.5	3,746.8	11.7		101.8		3.9	19.0
GOM Haddock	721.9	716.6	476.2	1.9	238.5	0.2		4.9	0.3
GB Yellowtail Flounder	951.0	940.9	939.1	1.8			10.2	0.0	0.0
SNE/MA Yellowtail Flounder	365.6	356.7	345.3	11.5			1.0	1.1	6.8
CC/GOM Yellowtail Flounder	761.3	720.1	711.5	8.6				38.5	2.7
Plaice	1,453.8	1,439.8	1,435.9	3.9				12.1	1.9
Witch Flounder	958.9	934.8	930.9	3.9				22.5	1.6
GB Winter Flounder	1,913.2	1,912.1	1,911.0	1.1				0.0	1.0
GOM Winter Flounder	241.4	155.7	153.1	2.6				76.9	8.8
SNE/MA Winter Flounder	61.3	3.6	3.3	0.3				32.1	25.5
Redfish	2,526.0	2,522.2	2,518.8	3.4				3.6	0.2
White Hake	2,999.0	2,994.9	2,981.8	13.1				2.6	1.5
Pollock	8,947.6	7,499.2	7,433.7	65.5				694.0	754.3
Northern Windowpane	0.4	0.0	0.0	0.0				0.0	0.4
Southern Windowpane	41.8	2.3	0.2	2.2				16.6	22.9
Ocean Pout	0.1	0.0	0.0	0.0				0.0	0.1
Halibut	19.0	11.4	10.3	1.0				7.1	0.6
Wolffish	0.1	0.0	0.0	0.0				0.0	0.1

**Landings extrapolated from observer data.

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Source: NMFS Northeast Regional Office

October 1, 2012, run date of June 28, 2012

Any value for a non-allocated species may include landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories. These are SNE winter flounder, northern windowpane, southern windowpane, ocean pout, halibut, and wolffish.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 3: FY 2011 Northeast Multispecies Estimated Discards (mt)

Final

Stock	Components with ACLs and sub-ACLs; (with accountability measures (AMs))							sub-components: No AMs	
	Total Groundfish	Groundfish Fishery	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	State Water ¹	Other
	A to G	A+B+C	A	B	C	D	E	F	G
GB cod	155.4	147.7	144.3	3.4					7.8
GOM cod	189.1	169.1	145.5	23.7	NA*				19.9
GB Haddock	302.0	82.0	82.0	0.0		0.0			220.0
GOM Haddock	15.6	7.5	7.4	0.1	NA*	0.0			8.1
GB Yellowtail Flounder	154.9	49.1	48.9	0.2			73.7	0.0	32.0
SNE/MA Yellowtail Flounder	138.0	19.5	18.7	0.8			109.9		8.6
CC/GOM Yellowtail Flounder	91.9	86.4	83.7	2.7					5.5
Plaice	208.3	196.3	195.7	0.5					12.1
Witch Flounder	218.5	62.4	62.0	0.3					156.1
GB Winter Flounder	67.4	13.2	13.2	0.1				0.0	54.1
GOM Winter Flounder	45.9	5.1	5.1	0.1				36.3	4.4
SNE/MA Winter Flounder	234.5	90.2	83.5	6.7				7.9	136.3
Redfish	196.6	184.6	184.4	0.2					12.0
White Hake	39.2	33.6	32.6	1.1					5.6
Pollock	117.0	113.2	109.4	3.8					3.8
Northern Windowpane	190.2	156.5	156.2	0.3					33.8
Southern Windowpane	486.8	109.2	82.8	26.3					377.6
Ocean Pout	88.5	60.7	56.3	4.4					27.8
Halibut	33.0	31.2	31.1	0.1					1.8
Wolffish	32.9	32.9	32.2	0.7					0.0

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the most recent assessments for these stocks

¹Commercial discard estimate not available

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Source: NMFS Northeast Regional Office

October 1, 2012, run date of June 28, 2012

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 4: FY 2011 Northeast Multispecies Annual Catch Limits (mt)

Stock	Components with ACLs and sub-ACLs; (with accountability measures (AMs))							sub-components: No AMs	
	Total ACL	Groundfish sub-ACL ¹	Sector sub-ACL	Common Pool sub-ACL	Recreational sub-ACL	Midwater Trawl Herring Fishery sub-ACL	Scallop Fishery sub-ACL	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
GB cod	4,540	4,301	4,208	93				48	191
GOM cod	8,545	7,649	4,721	104	2,824			597	299
GB Haddock	32,611	30,580	30,393	187		318		342	1,370
GOM Haddock	1,141	1,086	770	8	308	11		9	35
GB Yellowtail Flounder	1,416	1,142	1,122	20			200.8	0	73
SNE Yellowtail Flounder	641	524	404	120			82	7	27
CC/GOM Yellowtail Flounder	992	940	913	27				10	42
Plaice	3,280	3,108	3,038	70				34	138
Witch Flounder	1,304	1,236	1,211	25				14	55
GB Winter Flounder	2,118	2,007	1,993	14				0	111
GOM Winter Flounder	524	329	313	16				163	32
SNE/MA Winter Flounder	842	726	NA	726				72	45
Redfish	7,959	7,541	7,505	36				84	334
White Hake	3,138	2,974	2,946	28				33	132
Pollock	16,166	13,952	13,848	104				769	1,445
Northern Windowpane	161	110	NA	110				2	49
Southern Windowpane	225	154	NA	154				2	69
Ocean Pout	253	239	NA	239				3	11
Halibut	76	33	NA	33				39	4
Wolffish	77	73	NA	73				1	3

¹ Except for GOM cod and GOM haddock, the groundfish sub-ACL is equal to the commercial (sector + common pool) groundfish sub-ACL. For GOM cod and GOM haddock, the groundfish sub-ACL includes the commercial sub-ACL and the recreational sub-ACL. For inseason monitoring, the commercial groundfish sub-ACL for GOM cod is 4,825 mt and 778 mt for GOM haddock.

Values in metric tons of live weight

Source: NMFS Northeast Regional Office
October 1, 2012

Table 5: FY 2011 Northeast Multispecies Percents of Catch Limits Taken (%)

Final

Stock	Components with ACLs and sub-ACLs; (with accountability measures (AMs))							sub-components: No AMs	
	Total Groundfish	Groundfish Fishery	Sector	Common Pool	Recreational*	Midwater Trawl Herring Fishery	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
GB cod	75.0	76.2	76.4	66.1				81.1	47.3
GOM cod	74.3	79.8	92.5	89.9	58.1			36.3	9.7
GB Haddock	12.8	12.6	12.6	6.3		32.0		1.1	17.4
GOM Haddock	64.6	66.7	62.8	24.3	77.4	1.7		54.6	23.8
GB Yellowtail Flounder	78.1	86.7	88.1	10.1			41.8	NA	43.9
SNE Yellowtail Flounder	78.6	71.8	90.1	10.2			135.2	15.6	57.1
CC/GOM Yellowtail Flounder	86.0	85.8	87.1	42.1				384.8	19.3
Plaice	50.7	52.6	53.7	6.4				35.5	10.1
Witch Flounder	90.3	80.7	82.0	16.8				161.0	286.8
GB Winter Flounder	93.5	95.9	96.5	8.2				NA	49.7
GOM Winter Flounder	54.8	48.9	50.5	16.5				69.5	41.3
SNE/MA Winter Flounder	35.1	12.9	NA	NA				55.6	359.7
Redfish	34.2	35.9	36.0	9.9				4.3	3.7
White Hake	96.8	101.8	102.3	50.4				7.9	5.3
Pollock	56.1	54.6	54.5	66.6				90.3	52.5
Northern Windowpane	118.4	142.2	NA	NA				0.5	69.7
Southern Windowpane	234.9	72.4	NA	NA				829.1	580.4
Ocean Pout	35.0	25.4	NA	NA				0.0	253.0
Halibut	68.5	129.1	NA	NA				18.1	59.2
Wolffish	42.8	45.1	NA	NA				0.0	3.0

* To evaluate whether recreational catches exceeded any of the recreational sub-ACLs, and determine whether a recreational AM was triggered, the 2-year average of FY 2010 and FY 2011 was used. Please see Table 29.

Values in percent of live weight (%)

Sector and common pool include estimate of missing dealer reports

Source: NMFS Northeast Regional Office

October 1, 2012, run date of June 28, 2012

Any value for a non-allocated species may include landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories. These are SNE winter flounder, northern windowpane, southern windowpane, ocean pout, halibut, and wolffish.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 6: FY 2011 Northeast Multispecies Estimated State Water Sub-Component (mt)

Stock	Final								
	Total			Commercial			Recreational		
	Catch	Landings	Discard	Catch	Landings ¹	Discard ²	Catch	Landings	Discard
A+B+C+D	A+C	B+D	A+B	A	B	C+D	C	D	
GB cod	38.9	38.9		38.9	38.9		0.0	0.0	NA*
GOM cod	216.4	216.4		216.4	216.4		-.**	-.**	NA*
GB Haddock	3.9	3.9		3.9	3.9		NA*	NA*	NA*
GOM Haddock	4.9	4.9		4.9	4.9		-.**	-.**	NA*
GB Yellowtail Flounder	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
SNE/MA Yellowtail Flounder	1.1	1.1		1.1	1.1				
CC/GOM Yellowtail Flounder	38.5	38.5		38.5	38.5				
Plaice	12.1	12.1		12.1	12.1				
Witch Flounder	22.5	22.5		22.5	22.5				
GB Winter Flounder	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
GOM Winter Flounder	113.3	76.9	36.3	22.4	22.4		90.8	54.5	36.3
SNE/MA Winter Flounder	40.0	32.1	7.9	24.9	24.9		15.1	7.2	7.9
Redfish	3.6	3.6		3.6	3.6				
White Hake	2.6	2.6		2.6	2.6				
Pollock	694.0	694.0		36.8	36.8		657.3	657.3	NA*
Northern Windowpane	0.0	0.0		0.0	0.0				
Southern Windowpane	16.6	16.6		16.6	16.6				
Ocean Pout	0.0	0.0		0.0	0.0				
Halibut	7.1	7.1		7.1	7.1				
Wolffish	0.0	0.0		0.0	0.0				

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the most recent assessments for these stocks

**Some or all landings attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

¹CY 2012 commercial landings are estimated

FY 2011 GOM cod state recreational landings are estimated at 536.1 metric tons.

²Commercial discard estimate not available

FY 2011 GOM haddock state recreational landings are estimated at 12.3 metric tons.

Values in metric tons of live weight

Source: NMFS Northeast Regional Office

October 1, 2012, run date of June 28, 2012

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 7: FY 2011 Northeast Multispecies Other Sub-Component Catch Detail (mt)

Final

Stock	Total	SCALLOP ¹	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	REDCRAB	RESEARCH
GB cod	90.3	5.7	0.7	0.0	0.3	0.7	0.1	0.1	0.0	12.3
GOM cod	29.0	-	0.6	0.0	3.0	0.1	0.0	0.0	-	8.7
GB Haddock	239.0	2.4	0.2	0.0	11.9**	2.0	0.1	0.0	0.0	18.1
GOM Haddock	8.3	-	0.0	0.0	2.6**	0.1	-	-	-	0.2
GB Yellowtail Flounder	32.0	**	0.1	0.0	0.8	0.0	-	0.0	0.0	-
SNE Yellowtail Flounder	15.4	**	3.9	-	0.1	0.0	0.0	0.0	0.0	3.4
CC/GOM Yellowtail Flounder	8.1	2.9	0.1	0.0	0.5	0.0	0.0	0.0	-	2.5
Plaice	14.0	0.0	1.6	0.0	1.4	0.5	0.3	0.0	0.0	1.5
Witch Flounder	157.7	18.0	11.0	0.0	7.8	1.5	0.4	0.1	0.0	1.1
GB Winter Flounder	55.2	38.4	0.2	-	0.3	0.0	-	-	-	-
GOM Winter Flounder	13.2	2.0	0.0	0.0	0.2	0.0	-	-	-	0.2
SNE Winter Flounder	161.9	60.3	18.6	0.0	2.1	0.5	0.0	0.2	0.0	3.5
Redfish	12.2	0.0	4.0	0.0	0.2	0.1	0.0	0.0	0.0	0.1
White Hake	7.1	2.0	0.3	0.0	0.2	0.1	0.0	0.6	0.0	0.0
Pollock	758.1	-	1.0	0.0	0.5	0.2	0.1	0.0	0.0	0.6
Northern Windowpane	34.2	33.0	0.0	0.0	0.1	0.0	-	0.0	0.0	0.0
Southern Windowpane	400.5	135.3	86.5	-	1.4	0.5	0.1	0.6	0.0	0.0
Ocean Pout	27.8	6.4	2.5	0.0	0.9	0.1	0.0	0.0	0.0	0.0
Halibut	2.4	0.8	0.1	-	0.0	0.4	-	0.0	-	0.0
Wolffish	0.1	-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

**Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 7: FY 2011 Northeast Multispecies Other Sub-Component Catch Detail (mt)

Final

Stock	Total	SCUP	SHRIMP	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH ¹	WHITING	UNKNOWN	RECREATIONAL
GB cod	90.3	0.3	0.0	0.2	0.1	0.0	0.0	0.0	0.0	15.2	54.6
GOM cod	29.0	2.5	0.7	0.4	3.1	0.0	-	0.0	2.6	7.3	**
GB Haddock	239.0	0.1	0.1	81.0	41.9	0.0	0.0	0.0	0.7	80.4	NA*
GOM Haddock	8.3	-	0.5	0.0	0.8	-	-	0.0	1.9	2.4	**
GB Yellowtail Flounder	32.0	0.2	0.0	0.1	30.2	-	-	0.0	-	0.7	
SNE Yellowtail Flounder	15.4	1.6	0.0	1.0	0.6	0.0	0.0	0.0	0.0	4.7	
CC/GOM Yellowtail Flounder	8.1	0.3	0.1	0.0	0.4	0.0	-	0.0	0.3	0.9	
Plaice	14.0	1.0	0.0	2.3	1.5	0.0	0.0	0.0	0.1	3.7	
Witch Flounder	157.7	7.3	0.2	39.9	22.1	0.0	0.0	0.0	1.0	47.4	
GB Winter Flounder	55.2	0.9	0.0	0.2	13.3	-	-	-	0.0	1.7	
GOM Winter Flounder	13.2	-	0.0	0.0	0.1	-	-	0.0	0.2	0.2	10.3
SNE Winter Flounder	161.9	9.7	0.0	16.1	6.1	0.0	0.0	0.0	0.1	33.0	11.7
Redfish	12.2	2.7	0.0	0.9	0.9	0.0	0.0	0.0	0.0	3.4	
White Hake	7.1	0.3	0.0	1.1	0.7	0.0	0.0	0.0	0.0	1.6	
Pollock	758.1	0.7	0.0	0.1	0.2	0.0	0.0	0.0	0.0	6.2	748.5
Northern Windowpane	34.2	0.0	0.0	0.0	0.9	0.0	-	0.0	0.0	0.1	
Southern Windowpane	400.5	55.8	0.0	17.3	16.1	0.0	0.0	0.0	0.1	86.8	
Ocean Pout	27.8	1.7	0.0	5.6	3.2	0.0	0.0	0.0	0.1	7.2	
Halibut	2.4	0.1	0.0	0.3	0.1	-	-	-	0.0	0.5	
Wolffish	0.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

**Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 8: FY 2011 Northeast Multispecies Other Sub-Component Landings Detail (mt)

Final

Stock	Total	SCALLOP ¹	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	REDCRAB	RESEARCH
GB cod	82.5	0.0	0.3	-	-	0.5	-	0.1	-	12.3
GOM cod	9.1	-	-	-	-	0.0	-	-	-	8.7
GB Haddock	19.0	0.0	-	-	**	-	-	-	-	18.1
GOM Haddock	0.3	-	-	-	**	-	-	-	-	0.2
GB Yellowtail Flounder	0.0	**	-	-	-	-	-	-	-	-
SNE Yellowtail Flounder	6.8	**	1.3	-	-	-	-	0.0	-	3.4
CC/GOM Yellowtail Flounder	2.7	0.1	-	-	-	-	-	-	-	2.5
Plaice	1.9	0.0	0.1	-	-	-	-	-	-	1.5
Witch Flounder	1.6	0.2	0.1	-	-	0.0	-	-	-	1.1
GB Winter Flounder	1.0	0.9	-	-	-	-	-	-	-	-
GOM Winter Flounder	8.8	-	0.0	-	-	-	-	-	-	0.2
SNE Winter Flounder	25.5	1.2	3.6	-	-	0.0	-	0.1	-	3.5
Redfish	0.2	-	0.0	-	-	0.0	-	-	-	0.1
White Hake	1.5	0.0	0.2	-	-	0.0	-	0.6	-	0.0
Pollock	754.3	-	-	-	-	-	-	-	-	0.6
Northern Windowpane	0.4	-	-	-	-	-	-	-	-	-
Southern Windowpane	22.9	-	3.3	-	-	-	-	0.0	-	-
Ocean Pout	0.1	-	0.0	-	-	-	-	-	-	-
Halibut	0.6	-	-	-	-	0.4	-	0.0	-	-
Wolffish	0.1	-	-	-	-	-	-	-	-	-

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

**Some or all landings attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 8: FY 2011 Northeast Multispecies Other Sub-Component Landings Detail (mt)

Final

Stock	Total	SCUP	SHRIMP	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH ¹	WHITING	UNKNOWN	RECREATIONAL
GB cod	82.5	0.0	-	0.0	0.0	-	-	-	-	14.6	54.6
GOM cod	9.1	-	-	-	-	-	-	-	-	0.4	**
GB Haddock	19.0	-	-	-	-	-	-	-	-	0.8	NA*
GOM Haddock	0.3	-	-	-	-	-	-	-	-	0.0	**
GB Yellowtail Flounder	0.0	-	-	-	-	-	-	-	-	0.0	
SNE Yellowtail Flounder	6.8	-	-	-	-	-	-	-	-	2.1	
CC/GOM Yellowtail Flounder	2.7	-	-	-	-	-	-	-	-	0.1	
Plaice	1.9	-	-	0.0	0.1	-	-	-	-	0.1	
Witch Flounder	1.6	0.0	-	-	0.0	-	-	-	-	0.2	
GB Winter Flounder	1.0	0.0	-	0.0	-	-	-	-	-	0.1	
GOM Winter Flounder	8.8	-	-	-	-	-	-	-	-	0.0	8.5
SNE Winter Flounder	25.5	0.3	-	0.2	0.0	-	-	-	-	8.2	8.4
Redfish	0.2	-	-	0.1	0.0	-	-	-	-	0.0	
White Hake	1.5	0.2	-	-	0.1	-	-	-	-	0.4	
Pollock	754.3	-	-	-	0.0	-	-	-	-	5.3	748.5
Northern Windowpane	0.4	-	-	-	0.4	-	-	-	-	0.0	
Southern Windowpane	22.9	0.1	-	1.1	0.8	-	-	-	-	17.5	
Ocean Pout	0.1	-	-	-	-	-	-	-	-	0.1	
Halibut	0.6	-	-	-	-	-	-	-	-	0.2	
Wolfish	0.1	-	-	-	-	-	-	-	-	0.1	

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

**Some or all landings attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 9: FY 2011 Northeast Multispecies Other Sub-Component Estimated Discards Detail (mt)

Final

Stock	Total	SCALLOP ¹	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	REDCRAB	RESEARCH
GB cod	7.8	5.6	0.4	0.0	0.3	0.2	0.1	0.0	0.0	0.0
GOM cod	19.9	-	0.6	0.0	3.0	0.1	0.0	0.0	-	0.0
GB Haddock	220.0	2.4	0.2	0.0	11.9**	2.0	0.1	0.0	0.0	0.0
GOM Haddock	8.1	-	0.0	0.0	2.6**	0.1	-	-	-	0.0
GB Yellowtail Flounder	32.0	**	0.1	0.0	0.8	0.0	-	0.0	0.0	-
SNE Yellowtail Flounder	8.6	**	2.6	-	0.1	0.0	0.0	0.0	0.0	0.0
CC/GOM Yellowtail Flounder	5.5	2.8	0.1	0.0	0.5	0.0	0.0	0.0	-	0.0
Plaice	12.1	-	1.5	0.0	1.4	0.5	0.3	0.0	0.0	0.0
Witch Flounder	156.1	17.8	10.9	0.0	7.8	1.5	0.4	0.1	0.0	0.0
GB Winter Flounder	54.1	37.5	0.2	-	0.3	0.0	-	-	-	-
GOM Winter Flounder	4.4	2.0	0.0	0.0	0.2	0.0	-	-	-	0.0
SNE Winter Flounder	136.3	59.0	15.0	0.0	2.1	0.5	0.0	0.1	0.0	0.0
Redfish	12.0	0.0	4.0	0.0	0.2	0.1	0.0	0.0	0.0	0.0
White Hake	5.6	2.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0
Pollock	3.8	-	1.0	0.0	0.5	0.2	0.1	0.0	0.0	0.0
Northern Windowpane	33.8	33.0	0.0	0.0	0.1	0.0	-	0.0	0.0	0.0
Southern Windowpane	377.6	135.3	83.2	-	1.4	0.5	0.1	0.6	0.0	0.0
Ocean Pout	27.8	6.4	2.5	0.0	0.9	0.1	0.0	0.0	0.0	0.0
Halibut	1.8	0.8	0.1	-	0.0	0.0	-	0.0	-	0.0
Wolffish	0.0	-	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

**Some or all discards attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 9: FY 2011 Northeast Multispecies Other Sub-Component Estimated Discards Detail (mt)

Final

Stock	Total	SCUP	SHRIMP	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH	WHITING	UNKNOWN	RECREATIONAL
GB cod	7.8	0.2	0.0	0.2	0.1	0.0	0.0	0.0	0.0	0.6	NA*
GOM cod	19.9	2.5	0.7	0.4	3.1	0.0	-	0.0	2.6	6.9	NA*
GB Haddock	220.0	0.1	0.1	81.0	41.9	0.0	0.0	0.0	0.7	79.6	NA*
GOM Haddock	8.1	-	0.5	0.0	0.8	-	-	0.0	1.9	2.3	NA*
GB Yellowtail Flounder	32.0	0.2	0.0	0.1	30.2	-	-	0.0	-	0.7	
SNE Yellowtail Flounder	8.6	1.6	0.0	1.0	0.6	0.0	0.0	0.0	0.0	2.6	
CC/GOM Yellowtail Flounder	5.5	0.3	0.1	0.0	0.4	0.0	-	0.0	0.3	0.9	
Plaice	12.1	1.0	0.0	2.3	1.4	0.0	0.0	0.0	0.1	3.6	
Witch Flounder	156.1	7.3	0.2	39.9	22.1	0.0	0.0	0.0	1.0	47.2	
GB Winter Flounder	54.1	0.9	0.0	0.2	13.3	-	-	-	0.0	1.6	
GOM Winter Flounder	4.4	-	0.0	0.0	0.1	-	-	0.0	0.2	0.2	1.8
SNE Winter Flounder	136.3	9.5	0.0	15.9	6.1	0.0	0.0	0.0	0.1	24.7	3.3
Redfish	12.0	2.7	0.0	0.8	0.9	0.0	0.0	0.0	0.0	3.4	
White Hake	5.6	0.1	0.0	1.1	0.6	0.0	0.0	0.0	0.0	1.2	
Pollock	3.8	0.7	0.0	0.1	0.2	0.0	0.0	0.0	0.0	0.9	NA*
Northern Windowpane	33.8	0.0	0.0	0.0	0.5	0.0	-	0.0	0.0	0.1	
Southern Windowpane	377.6	55.6	0.0	16.2	15.3	0.0	0.0	0.0	0.1	69.3	
Ocean Pout	27.8	1.7	0.0	5.6	3.2	0.0	0.0	0.0	0.1	7.2	
Halibut	1.8	0.1	0.0	0.3	0.1	-	-	-	0.0	0.3	
Wolffish	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	

Values in metric tons of live weight

¹Based on scallop fishing year March, 2011 through February, 2012

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

Source: NMFS Northeast Regional Office
October 1, 2012, run date of June 28, 2012

**Some or all discards attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not counted above.

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 25: FY 2011 End of Year Accounting of U.S./Canada Multispecies - U.S. Catch (mt)

Stock	U.S. Catch	U.S. Catch by Fishery Component							
		Groundfish	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery*	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
Eastern U.S./Canada Cod	164.7	164.2	162.4	1.8				0.0	0.5
Eastern U.S./Canada Haddock	1,077.6	1,060.2	1,060.2	0.0		10.8		0.0	6.6
U.S./Canada Yellowtail Flounder	1,105.9	990.0	988.0	2.0			83.9	0.0	32.0

*Landings extrapolated from observer data.
 Values in live weight
 Includes estimate of missing dealer reports
 August 21, 2012, run date of June 28, 2012

Table 26: FY 2011 End of Year Northeast Multispecies Trips, Vessels, Observers, and A DAS Used in the U.S./Canada Area

Area ¹	Number of Vessels		Number of Trips		A DAS Used		Number of Observed Trips	
	Sector	Common Pool	Sector	Common Pool	Sector	Common Pool	Sector	Common Pool
Eastern U.S./Canada Area	64	1	416	1	1,996	8	116	0
Western U.S./Canada Area	127	4	1,105	13	5,036	93	314	1
Total	127	4	1,175	14	5,199	101	336	1

¹Area based on area fished. Totals don't sum due to multi-area trips

Source: NMFS Northeast Regional Office
 August 21, 2012, run date of June 28, 2012

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 25A: FY 2011 End of Year Accounting of U.S./Canada Multispecies - U.S. Landings (mt)

Stock	U.S. Landings	U.S. Catch by Fishery Component							
		Groundfish	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery*	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
Eastern U.S./Canada Cod	131.0	130.9	129.2	1.8				0.0	0.0
Eastern U.S./Canada Haddock	1,028.5	1,017.7	1,017.7	0.0		10.8		0.0	0.0
U.S./Canada Yellowtail Flounder	951.0	940.9	939.1	1.8			10.2	0.0	0.0

*Landings extrapolated from observer data.
 Values in live weight
 Includes estimate of missing dealer reports
 Source: NMFS Northeast Regional Office
 August 21, 2012, run date of June 28, 2012

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 25B: FY 2011 End of Year Accounting of U.S./Canada Multispecies - U.S. Discards (mt)

Stock	U.S. Discards	U.S. Catch by Fishery Component							
		Groundfish	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
Eastern U.S./Canada Cod	33.7	33.2	33.2	0.0				0.0	0.5
Eastern U.S./Canada Haddock	49.1	42.5	42.5	0.0		0.0		0.0	6.6
U.S./Canada Yellowtail Flounder	154.9	49.1	48.9	0.2			73.7	0.0	32.0

Values in live weight

Includes estimate of missing dealer reports

Source: NMFS Northeast Regional Office

August 21, 2012, run date of June 28, 2012

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 27: FY 2011 End of Year Accounting of U.S./Canada Multispecies - U.S. Catch TACs (mt)

Stock	U.S. TAC ^{1,2}	Fishery Component U.S. TAC							
		Groundfish	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
Eastern U.S./Canada Cod	200	200	196	4				0	0
Eastern U.S./Canada Haddock	9,460	9,640	9,581	59		NA		0	0
U.S./Canada Yellowtail Flounder	1,458.0	1,142.0	1,122.0	20.0			200.8	0	73.0

¹The 2011 U.S. TAC for Eastern Georges Bank haddock was transposed, so 9,640 mt actually got allocated to sectors and the common pool.

²The 2011 U.S. TAC for Georges Bank yellowtail flounder is different than the total ACL for this stock.

Values in live weight

Includes estimate of missing dealer reports

Source: NMFS Northeast Regional Office

August 21, 2012

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 28: FY 2011 End of Year Accounting of U.S./Canada Multispecies - Percent of U.S. TAC Caught (%)

Stock	% of U.S. TAC	Percent of Each Fishery Component U.S. TAC Caught							
		Groundfish	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	State Water	Other
	A to G	A+B+C	A	B	C	D	E	F	G
Eastern U.S./Canada Cod	82.4	82.1	82.8	44.9				NA	NA
Eastern U.S./Canada Haddock	11.4	11.0	11.1	0.0		NA		NA	NA
U.S./Canada Yellowtail Flounder	75.9	86.7	88.1	10.1			41.8	NA	43.9

Values in percent live weight (%)
 Includes estimate of missing dealer reports
 Source: NMFS Northeast Regional Office
 August 21, 2012, run date of June 28, 2012

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

Table 29: FY 2010 & 2011 GOM Cod and Haddock Recreational Catch (mt)

Final

Stock	Fishing Year	Recreational Catch				
		Catch	Landings	Discard	Recreational sub-ACL	Percent of Catch Limit Taken
		C+D	C	D		
GOM cod	2010	1,506.9	1,506.9	NA*	2,673.0	56.4
	2011	1,640.3	1,640.3	NA*	2,824.0	58.1
	Average	1,573.6	1,573.6	NA*	2,748.5	57.3
GOM Haddock	2010	297.4	297.4	NA*	324.0	91.8
	2011	238.5	238.5	NA*	308.0	77.4
	Average	267.9	267.9	NA*	316.0	84.8

*Estimates not applicable. Recreational amounts are not attributed to the ACL consistent with the assessments for these stocks used to set FY 2011 quotas.

To evaluate whether recreational catches exceeded any of the recreational sub-ACLs, the average recreational catch for FY 2010 and FY 2011 was compared to the average recreational sub-ACLs for the stock for FY 2010 and FY 2011.

FY 2010 & 2011 recreational estimates based on Marine Recreational Information Program (MRIP) data.

Values in metric tons of live weight

Source: NMFS Northeast Regional Office

October 1, 2012

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.