

Table 8 Abundance and Biomass from NEFSC Spring Surveys for Barndoor Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1968-2000.

| | weight/tow | | | number/tow | | | ind wt | length | | | | nonzero | | | |
|------|------------|--------|-------|------------|--------|-------|--------|--------|----|----------|---------|---------|---------|----|----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% mean | 95% max | tows | no fish | | |
| 1968 | 0.374 | 0.075 | 0.673 | 0.138 | 0.026 | 0.249 | 2.716 | 41 | 46 | 61 | 71.7 | 115 | 118 | 10 | 21 |
| 1969 | 0.658 | -0.364 | 1.681 | 0.145 | -0.011 | 0.301 | 4.539 | 33 | 42 | 70 | 83.1 | 119 | 120 | 8 | 22 |
| 1970 | 0.111 | 0.033 | 0.188 | 0.047 | 0.017 | 0.078 | 2.350 | 45 | 44 | 62 | 68.2 | 104 | 105 | 9 | 10 |
| 1971 | 0.116 | 0.018 | 0.214 | 0.102 | 0.021 | 0.183 | 1.134 | 26 | 31 | 59 | 57.1 | 69 | 80 | 8 | 20 |
| 1972 | 0.222 | 0.028 | 0.416 | 0.023 | 0.005 | 0.041 | 9.617 | 63 | 62 | 119 | 104.7 | 123 | 124 | 6 | 6 |
| 1973 | 0.010 | -0.001 | 0.022 | 0.017 | 0.000 | 0.034 | 0.621 | 51 | 51 | 51 | 54.1 | 59 | 60 | 3 | 3 |
| 1974 | 0.020 | -0.005 | 0.045 | 0.017 | -0.002 | 0.037 | 1.146 | 43 | 43 | 58 | 53.3 | 59 | 60 | 3 | 3 |
| 1975 | 0.001 | -0.001 | 0.003 | 0.001 | -0.001 | 0.003 | 0.900 | 60 | 60 | 60 | 60.0 | 60 | 60 | 1 | 1 |
| 1976 | 0.010 | -0.010 | 0.030 | 0.006 | -0.005 | 0.017 | 1.800 | 61 | 61 | 61 | 61.0 | 61 | 61 | 1 | 1 |
| 1977 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1978 | 0.015 | -0.009 | 0.040 | 0.016 | -0.006 | 0.039 | 0.933 | 51 | 50 | 55 | 56.3 | 61 | 62 | 2 | 3 |
| 1979 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1980 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1981 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1982 | 0.002 | -0.001 | 0.005 | 0.002 | -0.002 | 0.005 | 1.000 | 54 | 54 | 54 | 54.0 | 54 | 54 | 1 | 1 |
| 1983 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1984 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1985 | 0.001 | 0.000 | 0.002 | 0.007 | -0.004 | 0.017 | 0.076 | 20 | 20 | 20 | 24.6 | 37 | 38 | 2 | 2 |
| 1986 | 0.003 | -0.001 | 0.007 | 0.011 | -0.004 | 0.026 | 0.250 | 33 | 33 | 41 | 37.5 | 41 | 42 | 2 | 2 |
| 1987 | 0.002 | -0.002 | 0.006 | 0.007 | -0.006 | 0.020 | 0.300 | 37 | 37 | 37 | 37.0 | 37 | 37 | 1 | 1 |
| 1988 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1989 | 0.007 | -0.007 | 0.021 | 0.006 | -0.006 | 0.019 | 1.100 | 60 | 60 | 60 | 60.0 | 60 | 60 | 1 | 1 |
| 1990 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1991 | 0.002 | -0.002 | 0.006 | 0.007 | -0.006 | 0.020 | 0.300 | 38 | 38 | 38 | 38.0 | 38 | 38 | 1 | 1 |
| 1992 | 0.136 | -0.117 | 0.389 | 0.013 | -0.006 | 0.032 | 10.397 | 41 | 41 | 117 | 98.2 | 124 | 125 | 2 | 4 |
| 1993 | 0.032 | 0.024 | 0.039 | 0.028 | 0.005 | 0.051 | 1.147 | 31 | 31 | 37 | 45.3 | 89 | 90 | 5 | 5 |
| 1994 | 0.084 | -0.023 | 0.191 | 0.029 | -0.001 | 0.059 | 2.926 | 46 | 46 | 65 | 70.1 | 120 | 121 | 4 | 6 |
| 1995 | 0.015 | -0.007 | 0.037 | 0.012 | -0.005 | 0.029 | 1.254 | 55 | 55 | 63 | 59.6 | 63 | 64 | 2 | 2 |
| 1996 | 0.062 | -0.039 | 0.162 | 0.025 | -0.003 | 0.054 | 2.465 | 23 | 23 | 66 | 63.2 | 111 | 112 | 4 | 6 |
| 1997 | 0.077 | 0.006 | 0.148 | 0.035 | 0.007 | 0.063 | 2.216 | 39 | 39 | 67 | 68.7 | 89 | 90 | 6 | 7 |
| 1998 | 0.169 | -0.024 | 0.363 | 0.061 | 0.015 | 0.106 | 2.799 | 26 | 26 | 60 | 64.4 | 122 | 123 | 8 | 15 |
| 1999 | 0.279 | -0.102 | 0.660 | 0.052 | 0.011 | 0.094 | 5.343 | 28 | 28 | 74 | 80.9 | 125 | 126 | 8 | 11 |
| 2000 | 0.473 | 0.246 | 0.699 | 0.138 | 0.076 | 0.200 | 3.419 | 19 | 20 | 68 | 71.4 | 125 | 127 | 14 | 29 |

Table 9 Abundance and Biomass from NEFSC Autumn Surveys for Barndoor Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1963-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|--------|-------|------------|--------|-------|--------|--------|----|----------|---------|------|---------|----|-----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% mean | 95% max | tows | no fish | | |
| 1963 | 2.633 | 1.604 | 3.663 | 0.762 | 0.468 | 1.056 | 3.458 | 28 | 44 | 69 | 74.6 | 121 | 136 | 47 | 120 |
| 1964 | 1.212 | 0.489 | 1.934 | 0.400 | 0.229 | 0.570 | 3.030 | 40 | 41 | 69 | 72.7 | 112 | 122 | 32 | 63 |
| 1965 | 1.822 | 1.115 | 2.528 | 0.695 | 0.441 | 0.949 | 2.622 | 27 | 42 | 67 | 69.9 | 111 | 134 | 36 | 95 |
| 1966 | 0.811 | 0.394 | 1.229 | 0.459 | 0.243 | 0.675 | 1.767 | 23 | 38 | 60 | 63.0 | 88 | 115 | 26 | 62 |
| 1967 | 0.438 | -0.025 | 0.901 | 0.064 | 0.017 | 0.111 | 6.844 | 45 | 52 | 65 | 81.0 | 119 | 120 | 10 | 14 |
| 1968 | 0.285 | 0.123 | 0.447 | 0.132 | 0.067 | 0.198 | 2.150 | 42 | 42 | 67 | 69.1 | 96 | 132 | 18 | 29 |
| 1969 | 0.054 | -0.003 | 0.111 | 0.035 | -0.006 | 0.076 | 1.551 | 51 | 51 | 62 | 62.0 | 73 | 74 | 5 | 8 |
| 1970 | 0.066 | -0.046 | 0.178 | 0.011 | -0.005 | 0.027 | 5.868 | 66 | 66 | 65 | 89.1 | 128 | 129 | 2 | 2 |
| 1971 | 0.170 | -0.051 | 0.392 | 0.117 | -0.077 | 0.311 | 1.455 | 35 | 35 | 53 | 54.6 | 63 | 120 | 6 | 19 |
| 1972 | 0.096 | -0.073 | 0.265 | 0.012 | -0.001 | 0.026 | 7.751 | 59 | 59 | 70 | 90.3 | 132 | 133 | 3 | 3 |
| 1973 | 0.004 | -0.001 | 0.009 | 0.008 | -0.003 | 0.019 | 0.474 | 41 | 41 | 47 | 48.7 | 52 | 53 | 2 | 3 |
| 1974 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1975 | 0.017 | -0.016 | 0.049 | 0.010 | -0.010 | 0.031 | 1.600 | 70 | 70 | 70 | 70.0 | 70 | 70 | 1 | 2 |
| 1976 | 0.047 | 0.002 | 0.091 | 0.058 | -0.003 | 0.119 | 0.810 | 50 | 50 | 51 | 54.6 | 61 | 62 | 7 | 10 |
| 1977 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1978 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1979 | 0.009 | -0.008 | 0.026 | 0.003 | -0.003 | 0.009 | 3.000 | 78 | 78 | 78 | 78.0 | 78 | 78 | 1 | 1 |
| 1980 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1981 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1982 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1983 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | - | - | - | - | - | 0 | 0 |
| 1984 | 0.010 | -0.004 | 0.024 | 0.003 | 0.000 | 0.007 | 2.900 | 61 | 61 | 84 | 73.0 | 84 | 85 | 2 | 2 |
| 1985 | 0.004 | -0.004 | 0.012 | 0.002 | -0.002 | 0.005 | 2.300 | 70 | 70 | 70 | 70.0 | 70 | 70 | 1 | 1 |
| 1986 | 0.029 | -0.018 | 0.077 | 0.015 | -0.002 | 0.032 | 2.008 | 22 | 22 | 52 | 51.0 | 90 | 91 | 3 | 3 |
| 1987 | 0.014 | -0.005 | 0.032 | 0.012 | -0.004 | 0.027 | 1.200 | 53 | 53 | 63 | 58.5 | 63 | 64 | 2 | 2 |
| 1988 | 0.007 | -0.005 | 0.020 | 0.009 | -0.005 | 0.022 | 0.850 | 34 | 34 | 33 | 44.8 | 76 | 77 | 2 | 2 |
| 1989 | 0.005 | -0.005 | 0.014 | 0.002 | -0.002 | 0.007 | 2.100 | 71 | 71 | 71 | 71.0 | 71 | 71 | 1 | 1 |
| 1990 | 0.028 | -0.022 | 0.078 | 0.010 | -0.005 | 0.024 | 2.964 | 60 | 60 | 66 | 76.3 | 95 | 96 | 2 | 3 |
| 1991 | 0.031 | 0.000 | 0.062 | 0.020 | 0.000 | 0.040 | 1.579 | 54 | 54 | 61 | 61.3 | 73 | 74 | 4 | 5 |
| 1992 | 0.002 | -0.002 | 0.007 | 0.004 | -0.004 | 0.013 | 0.550 | 46 | 46 | 51 | 49.0 | 51 | 52 | 1 | 2 |
| 1993 | 0.141 | -0.040 | 0.321 | 0.023 | 0.004 | 0.042 | 6.180 | 45 | 45 | 74 | 86.6 | 127 | 128 | 5 | 6 |
| 1994 | 0.035 | 0.001 | 0.069 | 0.044 | 0.006 | 0.082 | 0.790 | 33 | 33 | 47 | 49.4 | 75 | 76 | 6 | 9 |
| 1995 | 0.111 | -0.009 | 0.231 | 0.040 | -0.006 | 0.085 | 2.810 | 48 | 48 | 62 | 70.9 | 113 | 114 | 4 | 10 |
| 1996 | 0.042 | -0.020 | 0.104 | 0.023 | 0.000 | 0.046 | 1.841 | 25 | 25 | 61 | 59.8 | 92 | 93 | 4 | 5 |
| 1997 | 0.105 | -0.024 | 0.234 | 0.026 | 0.004 | 0.047 | 4.065 | 36 | 36 | 79 | 73.3 | 124 | 125 | 5 | 5 |
| 1998 | 0.089 | -0.036 | 0.214 | 0.026 | 0.002 | 0.050 | 3.453 | 48 | 48 | 71 | 73.9 | 120 | 121 | 4 | 5 |
| 1999 | 0.300 | 0.051 | 0.549 | 0.085 | 0.041 | 0.130 | 3.511 | 23 | 23 | 54 | 68.0 | 120 | 121 | 13 | 15 |

Table 10 Abundance and Biomass from NEFSC Winter Surveys for Barndoor Skate for the Georges Bank to Mid-Atlantic Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1992-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | nonzero | | | |
|------|------------|--------|-------|------------|-------|-------|--------|--------|----|----------|---------|---------|---------|----|----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% mean | 95% max | tows | no fish | | |
| 1992 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | - | - | | | | | | 0 | 0 |
| 1993 | 0.123 | -0.066 | 0.311 | 0.052 | 0.004 | 0.100 | 2.358 | 20 | 20 | 65 | 57.3 | 119 | 120 | 4 | 6 |
| 1994 | 0.185 | -0.027 | 0.397 | 0.080 | 0.011 | 0.148 | 2.328 | 21 | 21 | 60 | 63.5 | 102 | 103 | 5 | 7 |
| 1995 | 0.362 | 0.121 | 0.603 | 0.198 | 0.056 | 0.340 | 1.828 | 33 | 33 | 62 | 63.6 | 88 | 109 | 11 | 24 |
| 1996 | 0.291 | 0.079 | 0.503 | 0.203 | 0.054 | 0.352 | 1.434 | 19 | 20 | 61 | 56.4 | 85 | 92 | 12 | 23 |
| 1997 | 0.618 | 0.208 | 1.028 | 0.275 | 0.032 | 0.519 | 2.247 | 35 | 38 | 65 | 67.7 | 112 | 117 | 10 | 28 |
| 1998 | 0.455 | 0.146 | 0.765 | 0.464 | 0.092 | 0.837 | 0.980 | 20 | 26 | 41 | 46.8 | 83 | 123 | 12 | 57 |
| 1999 | 1.053 | 0.347 | 1.760 | 0.709 | 0.318 | 1.099 | 1.486 | 23 | 27 | 46 | 53.2 | 113 | 124 | 22 | 81 |
| 2000 | 2.718 | 0.153 | 5.284 | 1.081 | 0.518 | 1.643 | 2.515 | 19 | 19 | 56 | 62.78 | 122 | 126 | 12 | 69 |

Figure 17 Abundance and Biomass of Barndoor Skate from the NEFSC Spring (Circles) and Autumn (Squares) Bottom Trawl Surveys from 1963-2000 in the Gulf of Maine – Southern New England Offshore Region




Figure 18 Abundance and Biomass of Barndoor Skate from the NEFSC Spring Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only

Mean Index in Solid Squares, 95% Confidence Interval in Open Squares





Figure 19 Abundance and Biomass of Barndoor Skate from the NEFSC Autumn Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only


Mean Index in Solid Squares, 95% Confidence Interval in Open Squares



The minimum length of barndoor skate caught in NEFSC surveys is 20 cm (8 in), and the largest individual caught was 136 cm (54 in) total length, during the 1963 autumn survey in the Gulf of Maine. The median length of the survey catch has ranged from 20 cm in the 1985 spring survey to 119 cm in the 1972 spring survey. The median length of the survey catch has been increasing in recent years in both the spring and autumn surveys, and is currently 70-75 cm (28-30 in; Figure 20). Length frequency distributions from the NEFSC spring and autumn surveys are presented in the SAW 30 documents as Figures B46 – B49 and are not reproduced in this SAFE Report. In general, the length frequency distributions show the decline in abundance of barndoor skate to survey catches of zero during the 1980s. Recent catches have included individuals as large as those recorded during the peak abundance of the 1960s, but the large number of fish between 40 and 80 cm evident during the 1960s is not apparent in recent surveys. The NEFSC winter survey length frequency distributions for 1998-1999 indicate a significant recent increase in the abundance of barndoor skate at lengths less than 80 cm (Figure B50 in SAW 30 document).

Research surveys and commercial fishery observer sampling conducted by Department of Fisheries and Oceans of Canada (DFO) in the broad geographic area between the Gulf of St. Lawrence and Georges Bank indicate two principal areas of barndoor skate concentration: Georges Bank/Fundian Channel and central Scotian Shelf (Figure B51 of the SAW 30 document). Barndoor skate were sporadically encountered throughout the 1970s, were nearly absent in the 1980s, and have shown an increase in abundance since the mid-1990s on the southwestern Scotian Shelf, on Brown's Bank, and in the Gulf of Maine (Simon and Frank 1999; Figures B52-B55 in the SAW 30 document). The DFO standardized research trawl survey begun on Georges Bank in 1986 found the abundance of barndoor skate was relatively low until the mid-1990s, but has been increasing since that time (Figures B56-B57 in the SAW 30 document). A broad range of sizes of barndoor skate have been encountered by DFO surveys on Georges Bank, ranging from 15 to 125 cm, suggesting the current population is composed of both juveniles and adults (Figure B57 in the SAW 30 document). Canadian commercial fishery observer sampling of both mobile and fixed gears indicates that commercial gear may regularly capture more and larger barndoor skate than are evident in research survey catches (Figures B58-B59 in the SAW 30 document). Recent information from commercial fisheries indicate a much greater depth distribution than previously expected (Kulka 1999; Figures B60-B62 in the SAW 30 document).

Figure 20 Percentiles of Length Composition (5, 50, and 95) of Barndoor Skate from the NEFSC Spring and Autumn Bottom Trawl Surveys from 1967-2000 in the Gulf of Maine to Southern New England Offshore Region



2.3.1.4 Thorny Skate

NEFSC bottom trawl surveys indicate that thorny skate are most abundant in the Gulf of Maine and Georges Bank offshore strata regions, with very few fish caught in inshore (< 27 meters depth), Southern New England, or Mid-Atlantic regions (**Appendix I**). In the NEFSC spring surveys (1968-2000), the annual total catch of thorny skate has ranged from 44 fish in 1999 to 574 fish in 1973. In the NEFSC autumn surveys (1963-1999), the annual total catch of thorny skate has ranged from 60 fish in 1998 to 874 fish in 1978. Calculated on a per tow basis, these spring and autumn survey catches equate to maximum stratified mean number per tow indices for the GOM-MA offshore strata of about 2 to 3 fish, or about 6.0 kg, per tow during the early 1970s (Table 11 and Table 12).

NEFSC survey indices for thorny skate have declined continuously over the last 30 years. Indices of thorny skate abundance and biomass from the NEFSC spring and autumn surveys were at a peak during the early 1970s, reaching 2.9 fish per tow (5.3 kg per tow) in the spring survey and 1.8 fish per tow (5.9 kg per tow) in the autumn survey. Kulka and Mowbray (1998) indicated a similar period of high abundance for thorny skate in Canadian waters. NEFSC indices of thorny skate abundance have declined steadily since the late 1970s, reaching historically low values in 1998 and 1999 that are only 10%-15 % of the peak observed in the 1970s (Figure 21 – Figure 23).

The minimum length of thorny skate caught in NEFSC surveys is about 10 cm (4 in), and the largest individual caught was 111 cm (44 in) total length, most recently during the 1977 spring survey on Georges Bank. The median length of the survey catch has ranged from 31 cm in the 1988 autumn survey to 63 cm in the 1971 autumn survey. The median length of the survey catch has trended downward through most of the survey time series, but has been increasing in recent years in autumn surveys, and is currently 40-50 cm (16-20 in; Figure 24). Length frequency distributions from the NEFSC spring and autumn surveys are presented in the SAW 30 documents as Figures B71 – B74 and are not reproduced in this SAFE Report. In general, the length frequency distributions show a pattern of decline in abundance of larger individuals consistent with an increase in total mortality over the survey time series.

Indices of abundance for thorny skate are available from MADMF spring and autumn research trawl surveys in the inshore waters of Massachusetts for the years 1978-1998. Thorny skate are abundant in state waters north of Cape Cod into the Gulf of Maine. MADMF indices of thorny skate biomass have been variable over the time series, but there is a decreasing trend evident in both the spring and autumn time series. The spring index has stabilized around the median of 0.3 kg/tow throughout the 1990s, while the autumn index has been below the median of 0.6 kg/tow since 1994 (Figure 25). Low sample sizes and high variances suggest that the time series of thorny skate mean lengths from the MADMF survey are not a reliable metric of trends in this stock (Figure 26).

Table 11 Abundance and Biomass from NEFSC Spring Surveys for Thorny Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1968-2000.

| | weight/tow | | | number/tow | | | ind wt | length | | | | nonzero | | | |
|------|------------|-------|-------|------------|-------|-------|--------|--------|----|----------|---------|---------|---------|----|-----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% mean | 95% max | tows | no fish | | |
| 1968 | 3.181 | 2.137 | 4.225 | 1.600 | 1.067 | 2.134 | 1.987 | 12 | 16 | 44 | 47.8 | 91 | 105 | 60 | 252 |
| 1969 | 4.526 | 3.186 | 5.865 | 1.680 | 1.161 | 2.199 | 2.694 | 12 | 13 | 47 | 51.1 | 98 | 109 | 64 | 294 |
| 1970 | 4.202 | 3.229 | 5.174 | 1.990 | 1.478 | 2.502 | 2.112 | 12 | 16 | 41 | 48.2 | 95 | 110 | 84 | 363 |
| 1971 | 3.683 | 2.475 | 4.891 | 1.974 | 1.473 | 2.475 | 1.866 | 12 | 15 | 44 | 47.8 | 95 | 116 | 81 | 424 |
| 1972 | 4.984 | 3.757 | 6.212 | 2.219 | 1.773 | 2.665 | 2.246 | 12 | 16 | 47 | 50.7 | 94 | 110 | 91 | 443 |
| 1973 | 6.622 | 4.867 | 8.377 | 3.562 | 2.640 | 4.483 | 1.859 | 12 | 15 | 44 | 47.9 | 91 | 108 | 75 | 574 |
| 1974 | 3.774 | 2.939 | 4.608 | 2.450 | 1.938 | 2.962 | 1.540 | 9 | 14 | 43 | 45.8 | 87 | 106 | 81 | 376 |
| 1975 | 3.189 | 2.222 | 4.157 | 1.360 | 0.990 | 1.731 | 2.344 | 10 | 15 | 46 | 50.5 | 95 | 102 | 62 | 192 |
| 1976 | 2.895 | 2.041 | 3.750 | 1.671 | 1.281 | 2.060 | 1.733 | 13 | 15 | 43 | 47.2 | 90 | 106 | 79 | 339 |
| 1977 | 1.623 | 1.175 | 2.070 | 0.942 | 0.675 | 1.209 | 1.722 | 12 | 15 | 42 | 48.1 | 89 | 111 | 74 | 213 |
| 1978 | 1.250 | 0.806 | 1.695 | 0.800 | 0.579 | 1.020 | 1.564 | 10 | 15 | 49 | 46.8 | 83 | 97 | 71 | 191 |
| 1979 | 1.079 | 0.729 | 1.429 | 0.582 | 0.410 | 0.754 | 1.853 | 12 | 17 | 51 | 50.5 | 84 | 102 | 68 | 163 |
| 1980 | 2.105 | 1.308 | 2.901 | 1.319 | 0.880 | 1.757 | 1.596 | 11 | 13 | 37 | 43.6 | 92 | 100 | 60 | 250 |
| 1981 | 2.700 | 2.065 | 3.335 | 1.535 | 1.139 | 1.930 | 1.760 | 9 | 13 | 47 | 48.1 | 87 | 100 | 60 | 255 |
| 1982 | 2.345 | 1.685 | 3.004 | 1.144 | 0.878 | 1.411 | 2.049 | 10 | 17 | 53 | 52.4 | 85 | 97 | 62 | 218 |
| 1983 | 2.142 | 1.398 | 2.886 | 0.968 | 0.728 | 1.209 | 2.212 | 12 | 15 | 52 | 52.3 | 91 | 103 | 55 | 156 |
| 1984 | 1.453 | 0.818 | 2.087 | 0.608 | 0.462 | 0.755 | 2.389 | 12 | 16 | 51 | 53.0 | 96 | 100 | 40 | 97 |
| 1985 | 3.074 | 2.124 | 4.024 | 1.413 | 1.060 | 1.766 | 2.175 | 11 | 14 | 44 | 48.4 | 95 | 102 | 59 | 209 |
| 1986 | 2.619 | 1.974 | 3.263 | 1.718 | 1.377 | 2.058 | 1.525 | 10 | 15 | 38 | 44.0 | 83 | 98 | 69 | 276 |
| 1987 | 1.469 | 0.805 | 2.133 | 0.852 | 0.646 | 1.058 | 1.724 | 14 | 16 | 42 | 46.6 | 87 | 109 | 53 | 141 |
| 1988 | 1.173 | 0.735 | 1.612 | 1.106 | 0.766 | 1.446 | 1.061 | 11 | 14 | 32 | 38.5 | 82 | 98 | 59 | 176 |
| 1989 | 1.481 | 0.793 | 2.169 | 1.221 | 0.801 | 1.640 | 1.213 | 11 | 15 | 34 | 40.0 | 84 | 101 | 57 | 175 |
| 1990 | 1.565 | 0.833 | 2.296 | 1.097 | 0.688 | 1.506 | 1.427 | 14 | 16 | 39 | 44.5 | 82 | 99 | 49 | 167 |
| 1991 | 1.542 | 0.945 | 2.139 | 0.858 | 0.569 | 1.147 | 1.797 | 11 | 13 | 47 | 48.5 | 89 | 99 | 47 | 132 |
| 1992 | 1.092 | 0.621 | 1.564 | 0.612 | 0.384 | 0.840 | 1.784 | 14 | 15 | 47 | 48.4 | 89 | 102 | 31 | 86 |
| 1993 | 0.700 | 0.366 | 1.034 | 0.486 | 0.327 | 0.646 | 1.440 | 13 | 13 | 36 | 42.0 | 91 | 105 | 37 | 79 |
| 1994 | 0.435 | 0.242 | 0.629 | 0.439 | 0.270 | 0.609 | 0.991 | 12 | 12 | 37 | 39.3 | 67 | 92 | 39 | 80 |
| 1995 | 0.564 | 0.307 | 0.821 | 0.384 | 0.236 | 0.533 | 1.467 | 9 | 12 | 42 | 45.8 | 84 | 92 | 31 | 66 |
| 1996 | 0.371 | 0.178 | 0.563 | 0.321 | 0.106 | 0.535 | 1.156 | 12 | 12 | 36 | 40.8 | 80 | 93 | 24 | 63 |
| 1997 | 0.422 | 0.117 | 0.727 | 0.270 | 0.153 | 0.387 | 1.560 | 15 | 20 | 47 | 47.9 | 82 | 87 | 25 | 47 |
| 1998 | 0.480 | 0.209 | 0.752 | 0.334 | 0.236 | 0.431 | 1.440 | 12 | 14 | 35 | 40.8 | 89 | 98 | 42 | 85 |
| 1999 | 0.369 | 0.093 | 0.646 | 0.255 | 0.163 | 0.347 | 1.448 | 11 | 17 | 40 | 46.2 | 83 | 89 | 26 | 44 |
| 2000 | 0.423 | 0.166 | 0.680 | 0.470 | 0.013 | 0.927 | 0.900 | 12 | 12 | 24 | 34.0 | 82 | 89 | 28 | 103 |

Table 12 Abundance and Biomass from NEFSC Autumn Surveys for Thorny Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1963-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|-------|-------|------------|-------|-------|--------|--------|----|-----|------|---------|---------|---------|-----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% | mean | 95% max | tows | no fish | |
| 1963 | 5.371 | 3.788 | 6.954 | 1.672 | 1.305 | 2.039 | 3.213 | 10 | 15 | 60 | 60.4 | 99 | 107 | 65 | 297 |
| 1964 | 4.403 | 3.273 | 5.534 | 1.651 | 1.110 | 2.192 | 2.667 | 10 | 14 | 49 | 52.7 | 96 | 110 | 66 | 278 |
| 1965 | 4.474 | 3.268 | 5.681 | 1.825 | 1.243 | 2.408 | 2.451 | 10 | 14 | 45 | 49.6 | 95 | 107 | 55 | 352 |
| 1966 | 7.971 | 6.163 | 9.780 | 2.371 | 1.855 | 2.886 | 3.362 | 9 | 13 | 61 | 59.4 | 95 | 112 | 72 | 364 |
| 1967 | 2.712 | 1.422 | 4.001 | 0.982 | 0.383 | 1.580 | 2.763 | 12 | 14 | 49 | 52.5 | 95 | 100 | 54 | 165 |
| 1968 | 4.421 | 3.321 | 5.521 | 1.440 | 1.040 | 1.840 | 3.071 | 12 | 16 | 55 | 57.5 | 97 | 107 | 59 | 217 |
| 1969 | 5.715 | 4.320 | 7.110 | 1.833 | 1.359 | 2.307 | 3.117 | 12 | 14 | 55 | 56.7 | 97 | 106 | 72 | 289 |
| 1970 | 7.347 | 5.630 | 9.065 | 2.216 | 1.474 | 2.958 | 3.316 | 8 | 19 | 57 | 60.4 | 98 | 109 | 77 | 403 |
| 1971 | 5.357 | 4.149 | 6.565 | 1.434 | 1.095 | 1.774 | 3.735 | 12 | 18 | 63 | 64.1 | 99 | 111 | 69 | 284 |
| 1972 | 4.119 | 2.974 | 5.263 | 1.717 | 1.302 | 2.132 | 2.399 | 12 | 16 | 51 | 53.1 | 94 | 105 | 75 | 306 |
| 1973 | 4.564 | 3.227 | 5.902 | 1.536 | 1.134 | 1.939 | 2.971 | 12 | 17 | 59 | 61.2 | 95 | 111 | 72 | 274 |
| 1974 | 3.038 | 2.166 | 3.910 | 1.392 | 1.025 | 1.759 | 2.182 | 10 | 14 | 50 | 51.1 | 89 | 111 | 79 | 293 |
| 1975 | 2.474 | 1.483 | 3.464 | 1.027 | 0.716 | 1.338 | 2.409 | 10 | 12 | 47 | 50.0 | 94 | 106 | 70 | 232 |
| 1976 | 1.720 | 1.003 | 2.437 | 0.798 | 0.543 | 1.052 | 2.157 | 12 | 15 | 44 | 49.1 | 91 | 103 | 57 | 143 |
| 1977 | 3.221 | 2.513 | 3.928 | 1.548 | 1.223 | 1.874 | 2.080 | 10 | 13 | 49 | 50.7 | 89 | 107 | 108 | 446 |
| 1978 | 4.291 | 3.473 | 5.109 | 2.145 | 1.643 | 2.648 | 2.000 | 10 | 16 | 49 | 51.1 | 88 | 107 | 155 | 874 |
| 1979 | 3.612 | 2.750 | 4.474 | 1.283 | 0.864 | 1.702 | 2.815 | 11 | 21 | 59 | 59.5 | 89 | 101 | 134 | 486 |
| 1980 | 4.601 | 3.344 | 5.859 | 1.882 | 1.484 | 2.280 | 2.445 | 11 | 14 | 54 | 54.4 | 90 | 100 | 84 | 416 |
| 1981 | 3.339 | 2.551 | 4.127 | 1.305 | 0.957 | 1.653 | 2.559 | 12 | 15 | 55 | 57.1 | 90 | 103 | 71 | 223 |
| 1982 | 0.646 | 0.312 | 0.981 | 0.393 | 0.194 | 0.592 | 1.644 | 11 | 13 | 33 | 43.0 | 85 | 96 | 31 | 83 |
| 1983 | 2.409 | 1.553 | 3.266 | 0.833 | 0.589 | 1.077 | 2.892 | 15 | 20 | 56 | 58.8 | 93 | 108 | 49 | 121 |
| 1984 | 2.887 | 1.978 | 3.795 | 1.270 | 0.975 | 1.565 | 2.272 | 10 | 13 | 48 | 49.8 | 94 | 107 | 70 | 211 |
| 1985 | 2.877 | 1.765 | 3.988 | 1.438 | 1.094 | 1.783 | 2.000 | 12 | 16 | 49 | 49.6 | 87 | 103 | 66 | 260 |
| 1986 | 1.629 | 1.068 | 2.189 | 1.019 | 0.771 | 1.268 | 1.598 | 11 | 15 | 35 | 44.2 | 83 | 101 | 61 | 183 |
| 1987 | 0.944 | 0.590 | 1.297 | 0.841 | 0.600 | 1.082 | 1.123 | 12 | 14 | 36 | 40.2 | 78 | 92 | 49 | 143 |
| 1988 | 1.488 | 0.998 | 1.978 | 1.099 | 0.702 | 1.497 | 1.354 | 13 | 15 | 31 | 41.5 | 84 | 101 | 56 | 208 |
| 1989 | 1.883 | 0.980 | 2.786 | 1.129 | 0.787 | 1.471 | 1.668 | 12 | 14 | 40 | 46.2 | 85 | 101 | 63 | 198 |
| 1990 | 1.704 | 1.090 | 2.318 | 1.040 | 0.744 | 1.335 | 1.639 | 12 | 17 | 42 | 47.2 | 85 | 95 | 53 | 202 |
| 1991 | 1.632 | 0.519 | 2.745 | 0.921 | 0.591 | 1.251 | 1.772 | 13 | 15 | 47 | 49.5 | 86 | 108 | 54 | 153 |
| 1992 | 0.962 | 0.551 | 1.373 | 0.775 | 0.461 | 1.088 | 1.242 | 12 | 13 | 36 | 41.2 | 83 | 99 | 48 | 144 |
| 1993 | 1.658 | 0.639 | 2.676 | 0.901 | 0.440 | 1.361 | 1.840 | 12 | 13 | 47 | 47.8 | 91 | 101 | 50 | 157 |
| 1994 | 1.509 | 0.343 | 2.675 | 0.981 | 0.311 | 1.652 | 1.538 | 13 | 17 | 45 | 46.9 | 84 | 97 | 41 | 170 |
| 1995 | 0.783 | 0.331 | 1.235 | 0.639 | 0.183 | 1.095 | 1.226 | 13 | 14 | 39 | 42.2 | 72 | 99 | 37 | 107 |
| 1996 | 0.814 | 0.360 | 1.269 | 0.602 | 0.362 | 0.842 | 1.352 | 14 | 14 | 39 | 43.3 | 85 | 99 | 37 | 102 |
| 1997 | 0.849 | 0.405 | 1.293 | 0.404 | 0.241 | 0.567 | 2.101 | 12 | 20 | 50 | 52.3 | 83 | 99 | 33 | 79 |
| 1998 | 0.648 | 0.297 | 0.999 | 0.307 | 0.145 | 0.468 | 2.113 | 13 | 14 | 51 | 52.4 | 87 | 93 | 30 | 60 |
| 1999 | 0.479 | 0.249 | 0.710 | 0.326 | 0.195 | 0.457 | 1.469 | 13 | 14 | 41 | 46.3 | 87 | 94 | 38 | 72 |

Figure 21 Abundance and Biomass of Thorny Skate from the NEFSC Spring (Circles) and Autumn (Squares) Bottom Trawl Surveys from 1963-2000 in the Gulf of Maine to Southern New England Offshore Region




Figure 22 Abundance and Biomass of Thorny Skate from the NEFSC Autumn Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only

Mean Index In Solid Squares, 95% Confidence Interval In Open Squares




Figure 23 Abundance and Biomass of Thorny Skate from the NEFSC Spring Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only

Mean Index In Solid Squares, 95% Confidence Interval In Open Squares




Figure 24 Percentiles of Length Composition (5, 50, and 95) of Thorny Skate from the NEFSC Spring and Autumn Bottom Trawl Surveys from 1967-2000 in the Gulf of Maine to Southern New England Offshore Region

**Thorny Skate: GOM-SNE Offshore
Percentiles of Length Composition**




Figure 25 Abundance and Biomass of Thorny Skate from the Massachusetts Spring (Circles) and Autumn (Squares) Finfish Bottom Trawl Survey in State Waters

Thorny Skate - Massachusetts Trawl Survey







Figure 26 Stratified Mean Total Length (cm) of Thorny Skate from the Massachusetts Spring and Autumn Bottom Trawl Surveys from 1978-1999

**Thorny Skate - Massachusetts Trawl Survey
Stratified Mean Length**



2.3.1.5 Smooth Skate

NEFSC bottom trawl surveys indicate that smooth skate are most abundant in the Gulf of Maine and Georges Bank offshore strata regions, with very few fish caught in inshore (< 27 meters depth), Southern New England, or Mid-Atlantic regions (**Appendix I**). In the NEFSC spring surveys (1968-2000), the annual total catch of smooth skate has ranged from 12 fish in 1996 to 179 fish in 1973. In the NEFSC autumn surveys (1963-1999), the annual total catch of smooth skate has ranged from 10 fish in 1976 to 130 fish in 1978. Calculated on a per tow basis, these spring and autumn survey catches equate to maximum stratified mean number per tow indices for the GOM-MA offshore strata of 0.6 to 1.6 fish, or about 0.6 to 0.9 kg, per tow during the 1970s (Table 13 and Table 14).

Indices of smooth skate abundance and biomass from the NEFSC surveys were at a peak during the early 1970s for the spring series and the late 1970s for the autumn series. NEFSC survey indices declined during the 1980s, before stabilizing during the early 1990s at about 25% of the autumn and 50% of the spring survey index values of the 1970s. There is evidence in the spring 1998-1999 indices of a recent increase in smooth skate abundance (Figure 27 – Figure 29).

The minimum length of smooth skate caught in NEFSC surveys is about 8 cm (3 in), and the largest individual caught was 71 cm (28 in) total length, during the 1984 autumn survey on Georges Bank. The median length of the survey catch has ranged from 26 cm in the 1993 autumn survey to 53 cm in the 1971 autumn survey. The median length of the survey catch in the GOM offshore region shows no trend over the full survey time series, and is currently at about 30 cm (12 in)(Figure 30). Length frequency distributions from the NEFSC spring and autumn surveys are presented in the SAW 30 documents as Figures B82 – B85 and are not reproduced in this SAFE Report. In general, the length frequency distributions show modes at 30 and 50 cm. The relatively high abundances evident in the 1969-1983 spring surveys at the larger mode may represent the accumulated abundance at several older ages. Truncation of the larger mode is evident in the spring distributions during the 1980s and most of the 1990s. The 1999 spring survey length frequency distribution may indicate strong recruitment in the region.

Table 13 Abundance and Biomass from NEFSC Spring Surveys for Smooth Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1968-2000.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|-------|-------|------------|--------|-------|--------|--------|----|-----|------|---------|---------|---------|-----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% | mean | 95% max | tows | no fish | |
| 1968 | 0.211 | 0.080 | 0.342 | 0.484 | 0.129 | 0.838 | 0.436 | 12 | 24 | 41 | 42.1 | 58 | 64 | 17 | 41 |
| 1969 | 0.377 | 0.193 | 0.562 | 0.834 | 0.521 | 1.147 | 0.452 | 11 | 19 | 48 | 43.3 | 58 | 63 | 28 | 82 |
| 1970 | 0.346 | 0.134 | 0.557 | 0.702 | 0.376 | 1.028 | 0.492 | 9 | 14 | 47 | 40.9 | 57 | 61 | 25 | 68 |
| 1971 | 0.800 | 0.395 | 1.205 | 1.185 | 0.650 | 1.719 | 0.675 | 9 | 20 | 51 | 48.2 | 61 | 63 | 40 | 114 |
| 1972 | 0.621 | 0.355 | 0.886 | 1.016 | 0.582 | 1.450 | 0.611 | 14 | 20 | 47 | 44.3 | 59 | 64 | 34 | 122 |
| 1973 | 1.000 | 0.745 | 1.255 | 1.907 | 1.401 | 2.414 | 0.524 | 9 | 24 | 45 | 44.2 | 59 | 65 | 51 | 179 |
| 1974 | 1.092 | 0.594 | 1.590 | 2.003 | 1.109 | 2.896 | 0.545 | 9 | 9 | 47 | 42.7 | 59 | 63 | 47 | 172 |
| 1975 | 0.240 | 0.133 | 0.346 | 0.383 | 0.224 | 0.543 | 0.626 | 19 | 25 | 49 | 46.8 | 59 | 61 | 22 | 37 |
| 1976 | 0.534 | 0.413 | 0.655 | 1.150 | 0.870 | 1.429 | 0.464 | 12 | 16 | 43 | 39.8 | 57 | 60 | 49 | 134 |
| 1977 | 0.122 | 0.066 | 0.178 | 0.302 | 0.158 | 0.445 | 0.405 | 15 | 18 | 40 | 41.4 | 57 | 60 | 28 | 45 |
| 1978 | 0.251 | 0.144 | 0.358 | 0.413 | 0.258 | 0.567 | 0.609 | 24 | 26 | 50 | 46.7 | 58 | 61 | 33 | 56 |
| 1979 | 0.218 | 0.097 | 0.340 | 0.410 | 0.163 | 0.657 | 0.533 | 15 | 19 | 39 | 40.2 | 54 | 61 | 27 | 54 |
| 1980 | 0.484 | 0.316 | 0.651 | 0.948 | 0.625 | 1.271 | 0.510 | 16 | 20 | 42 | 41.9 | 56 | 60 | 42 | 84 |
| 1981 | 0.358 | 0.227 | 0.489 | 0.782 | 0.513 | 1.050 | 0.458 | 8 | 13 | 38 | 37.2 | 57 | 65 | 38 | 70 |
| 1982 | 0.152 | 0.057 | 0.247 | 0.225 | 0.092 | 0.357 | 0.677 | 11 | 10 | 52 | 45.6 | 57 | 64 | 14 | 23 |
| 1983 | 0.363 | 0.219 | 0.507 | 0.531 | 0.335 | 0.727 | 0.683 | 11 | 21 | 50 | 47.9 | 57 | 69 | 25 | 50 |
| 1984 | 0.065 | 0.010 | 0.120 | 0.124 | 0.026 | 0.221 | 0.523 | 19 | 20 | 48 | 39.8 | 59 | 60 | 9 | 13 |
| 1985 | 0.211 | 0.136 | 0.286 | 0.450 | 0.298 | 0.602 | 0.469 | 18 | 20 | 41 | 40.4 | 57 | 63 | 31 | 59 |
| 1986 | 0.250 | 0.137 | 0.362 | 0.466 | 0.256 | 0.677 | 0.536 | 20 | 24 | 48 | 46.7 | 59 | 65 | 30 | 93 |
| 1987 | 0.069 | 0.029 | 0.108 | 0.105 | 0.044 | 0.166 | 0.655 | 43 | 42 | 48 | 50.2 | 59 | 62 | 12 | 15 |
| 1988 | 0.115 | 0.044 | 0.186 | 0.328 | 0.175 | 0.480 | 0.350 | 11 | 13 | 36 | 36.3 | 57 | 60 | 24 | 49 |
| 1989 | 0.225 | 0.107 | 0.343 | 0.620 | 0.402 | 0.838 | 0.363 | 13 | 15 | 37 | 38.8 | 60 | 63 | 30 | 88 |
| 1990 | 0.152 | 0.010 | 0.294 | 0.294 | 0.080 | 0.509 | 0.515 | 11 | 16 | 46 | 44.0 | 57 | 62 | 18 | 40 |
| 1991 | 0.137 | 0.073 | 0.200 | 0.237 | 0.136 | 0.337 | 0.576 | 11 | 17 | 49 | 47.1 | 59 | 62 | 22 | 34 |
| 1992 | 0.063 | 0.025 | 0.101 | 0.104 | 0.035 | 0.172 | 0.608 | 22 | 40 | 49 | 48.5 | 56 | 57 | 12 | 16 |
| 1993 | 0.086 | 0.021 | 0.151 | 0.214 | 0.020 | 0.408 | 0.403 | 21 | 23 | 42 | 41.2 | 56 | 58 | 14 | 35 |
| 1994 | 0.098 | 0.043 | 0.153 | 0.176 | 0.082 | 0.269 | 0.558 | 29 | 29 | 47 | 47.1 | 56 | 58 | 15 | 30 |
| 1995 | 0.101 | 0.050 | 0.152 | 0.234 | 0.119 | 0.349 | 0.432 | 9 | 20 | 42 | 41.9 | 55 | 59 | 18 | 33 |
| 1996 | 0.036 | 0.014 | 0.058 | 0.084 | 0.038 | 0.129 | 0.429 | 20 | 19 | 48 | 43.8 | 53 | 59 | 10 | 12 |
| 1997 | 0.037 | 0.015 | 0.059 | 0.122 | 0.035 | 0.208 | 0.307 | 17 | 20 | 36 | 38.9 | 55 | 58 | 11 | 22 |
| 1998 | 0.200 | 0.089 | 0.311 | 0.410 | 0.206 | 0.613 | 0.489 | 9 | 19 | 46 | 44.6 | 56 | 60 | 28 | 77 |
| 1999 | 0.243 | 0.068 | 0.418 | 0.925 | -0.074 | 1.924 | 0.262 | 18 | 20 | 32 | 35.6 | 51 | 65 | 23 | 111 |
| 2000 | 0.060 | 0.025 | 0.095 | 0.220 | -0.021 | 0.460 | 0.272 | 10 | 10 | 27 | 30.9 | 59 | 62 | 13 | 30 |

Table 14 Abundance and Biomass from NEFSC Autumn Surveys for Smooth Skate for the Gulf of Maine to Southern New England Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1963-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|-------|-------|------------|-------|-------|--------|--------|----|----------|---------|------|---------|----|-----|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% mean | 95% max | tows | no fish | | |
| 1963 | 0.498 | 0.306 | 0.689 | 0.543 | 0.282 | 0.804 | 0.917 | 9 | 20 | 48 | 43.9 | 58 | 62 | 26 | 53 |
| 1964 | 0.326 | 0.152 | 0.501 | 0.360 | 0.209 | 0.512 | 0.906 | 9 | 20 | 42 | 41.7 | 59 | 64 | 19 | 35 |
| 1965 | 0.475 | 0.140 | 0.811 | 1.221 | 0.440 | 2.001 | 0.389 | 11 | 16 | 35 | 38.1 | 56 | 64 | 27 | 94 |
| 1966 | 0.323 | 0.175 | 0.471 | 0.867 | 0.519 | 1.216 | 0.372 | 13 | 17 | 37 | 38.6 | 58 | 59 | 28 | 60 |
| 1967 | 0.152 | 0.036 | 0.268 | 0.293 | 0.118 | 0.469 | 0.518 | 22 | 24 | 48 | 46.5 | 62 | 69 | 16 | 27 |
| 1968 | 0.385 | 0.211 | 0.559 | 0.665 | 0.375 | 0.955 | 0.579 | 17 | 20 | 48 | 45.9 | 58 | 62 | 24 | 56 |
| 1969 | 0.290 | 0.131 | 0.449 | 0.604 | 0.282 | 0.925 | 0.481 | 12 | 16 | 41 | 39.6 | 58 | 64 | 21 | 50 |
| 1970 | 0.232 | 0.121 | 0.343 | 0.530 | 0.289 | 0.771 | 0.437 | 9 | 13 | 45 | 38.3 | 59 | 62 | 25 | 50 |
| 1971 | 0.157 | 0.077 | 0.238 | 0.250 | 0.120 | 0.379 | 0.631 | 17 | 36 | 53 | 51.0 | 57 | 59 | 18 | 27 |
| 1972 | 0.332 | 0.185 | 0.478 | 0.499 | 0.285 | 0.713 | 0.664 | 16 | 24 | 49 | 49.8 | 62 | 64 | 30 | 52 |
| 1973 | 0.311 | 0.199 | 0.423 | 0.506 | 0.344 | 0.667 | 0.614 | 17 | 22 | 48 | 46.9 | 58 | 60 | 32 | 56 |
| 1974 | 0.123 | 0.055 | 0.192 | 0.180 | 0.088 | 0.273 | 0.684 | 11 | 11 | 50 | 48.5 | 60 | 63 | 13 | 21 |
| 1975 | 0.076 | 0.029 | 0.123 | 0.104 | 0.043 | 0.165 | 0.727 | 21 | 30 | 49 | 46.7 | 56 | 57 | 12 | 15 |
| 1976 | 0.039 | 0.004 | 0.074 | 0.077 | 0.020 | 0.135 | 0.501 | 17 | 36 | 41 | 43.9 | 52 | 60 | 9 | 10 |
| 1977 | 0.376 | 0.274 | 0.478 | 0.600 | 0.443 | 0.757 | 0.627 | 19 | 24 | 48 | 44.9 | 56 | 61 | 50 | 84 |
| 1978 | 0.450 | 0.240 | 0.661 | 0.635 | 0.359 | 0.912 | 0.709 | 8 | 25 | 50 | 48.0 | 59 | 66 | 49 | 130 |
| 1979 | 0.182 | 0.075 | 0.288 | 0.239 | 0.116 | 0.362 | 0.761 | 9 | 29 | 50 | 48.7 | 60 | 62 | 31 | 60 |
| 1980 | 0.343 | 0.167 | 0.519 | 0.522 | 0.254 | 0.789 | 0.658 | 15 | 23 | 52 | 46.4 | 58 | 62 | 37 | 60 |
| 1981 | 0.119 | 0.039 | 0.199 | 0.167 | 0.069 | 0.264 | 0.715 | 23 | 26 | 49 | 48.1 | 60 | 61 | 13 | 18 |
| 1982 | 0.039 | 0.007 | 0.071 | 0.074 | 0.025 | 0.123 | 0.521 | 9 | 9 | 49 | 41.9 | 63 | 64 | 11 | 11 |
| 1983 | 0.146 | 0.056 | 0.236 | 0.255 | 0.085 | 0.426 | 0.573 | 14 | 14 | 46 | 40.9 | 57 | 59 | 12 | 24 |
| 1984 | 0.199 | 0.106 | 0.292 | 0.389 | 0.171 | 0.607 | 0.512 | 14 | 22 | 37 | 39.2 | 58 | 71 | 23 | 39 |
| 1985 | 0.210 | 0.088 | 0.332 | 0.340 | 0.180 | 0.500 | 0.617 | 12 | 15 | 51 | 45.2 | 59 | 63 | 28 | 64 |
| 1986 | 0.209 | 0.118 | 0.300 | 0.392 | 0.216 | 0.567 | 0.534 | 13 | 21 | 47 | 45.0 | 63 | 66 | 24 | 63 |
| 1987 | 0.095 | 0.045 | 0.145 | 0.164 | 0.081 | 0.247 | 0.581 | 15 | 15 | 48 | 44.8 | 60 | 61 | 19 | 28 |
| 1988 | 0.284 | 0.103 | 0.465 | 0.446 | 0.223 | 0.670 | 0.637 | 20 | 20 | 51 | 48.3 | 59 | 65 | 27 | 90 |
| 1989 | 0.128 | 0.072 | 0.185 | 0.336 | 0.194 | 0.478 | 0.382 | 13 | 16 | 33 | 36.8 | 59 | 62 | 27 | 52 |
| 1990 | 0.194 | 0.120 | 0.268 | 0.332 | 0.202 | 0.462 | 0.584 | 16 | 23 | 48 | 46.4 | 58 | 62 | 27 | 45 |
| 1991 | 0.167 | 0.070 | 0.265 | 0.335 | 0.188 | 0.482 | 0.500 | 18 | 20 | 46 | 43.9 | 57 | 62 | 25 | 59 |
| 1992 | 0.126 | 0.024 | 0.228 | 0.316 | 0.120 | 0.511 | 0.400 | 12 | 18 | 43 | 40.0 | 58 | 60 | 16 | 56 |
| 1993 | 0.227 | 0.107 | 0.346 | 0.818 | 0.273 | 1.362 | 0.277 | 13 | 13 | 26 | 32.6 | 56 | 62 | 29 | 123 |
| 1994 | 0.099 | 0.030 | 0.169 | 0.269 | 0.105 | 0.433 | 0.370 | 11 | 11 | 36 | 38.0 | 57 | 59 | 17 | 36 |
| 1995 | 0.189 | 0.115 | 0.263 | 0.764 | 0.315 | 1.214 | 0.247 | 10 | 13 | 30 | 32.6 | 56 | 59 | 29 | 119 |
| 1996 | 0.176 | 0.093 | 0.260 | 0.421 | 0.249 | 0.594 | 0.418 | 15 | 18 | 46 | 41.6 | 56 | 59 | 26 | 55 |
| 1997 | 0.232 | 0.117 | 0.347 | 0.449 | 0.232 | 0.665 | 0.517 | 16 | 21 | 47 | 45.2 | 60 | 64 | 20 | 59 |
| 1998 | 0.028 | 0.005 | 0.051 | 0.108 | 0.021 | 0.194 | 0.263 | 18 | 17 | 29 | 35.2 | 51 | 53 | 11 | 18 |
| 1999 | 0.070 | 0.032 | 0.109 | 0.110 | 0.050 | 0.171 | 0.638 | 22 | 22 | 50 | 48.7 | 60 | 62 | 16 | 22 |

Figure 27 Abundance and Biomass of Smooth Skate from the NEFSC Spring (Circles) and Autumn (Squares) Bottom Trawl Surveys from 1963-2000 in the Gulf of Maine to Southern New England Offshore Region




Figure 28 Abundance and Biomass of Smooth Skate from the NEFSC Spring Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only

Mean Index in Solid Squares, 95% Confidence Interval in Open Squares




Figure 29 Abundance and Biomass of Smooth Skate from the NEFSC Autumn Bottom Trawl Survey in the Gulf of Maine to Southern New England Region, Offshore Strata Only

Mean Index in Solid Squares, 95% Confidence Interval in Open Squares





Figure 30 Percentiles of Length Composition (5, 50, and 95) of Smooth Skate from the NEFSC Spring and Autumn Bottom Trawl Surveys from 1967-2000 in the Gulf of Maine to Southern New England Offshore Region

**Smooth Skate: GOM-SNE Offshore
Percentiles of Length Composition**



2.3.1.6 Clearnose Skate

NEFSC bottom trawl surveys indicate that clearnose skate are most abundant in the Mid-Atlantic offshore and inshore strata regions, with very few fish caught in Southern New England and no fish caught in other survey regions (**Appendix I**). In the NEFSC spring surveys (1976-2000), the annual total catch of clearnose skate has ranged from 9 fish in 1979 to 136 fish in 1993. In the NEFSC autumn surveys (1975-1999), the annual total catch of clearnose skate has ranged from 19 fish in 1983 to 129 fish in 1994. Calculated on a per tow basis, these spring and autumn survey catches equate to maximum stratified mean number per tow indices for the Mid-Atlantic offshore and inshore strata set of 1.2-1.6 fish, or about 0.8-0.9 kg, per tow during the mid 1990s (Table 15 and Table 16).

The catchability of clearnose skate in the recently instituted NEFSC winter bottom trawl survey (which substitutes a chain sweep with small cookies for the large rollers used in the spring and autumn surveys, to better target flatfish) is significantly higher than in the spring and autumn series. NEFSC winter survey (1992-1999) annual catches of clearnose skate have ranged from 343 fish in 1999 to 3,086 fish in 1996, equating to a maximum stratified mean catch per tow of 12 fish or 15 kg per tow in 1996 (Table 17). The winter survey is focused in the Southern New England and Mid-Atlantic offshore regions, with a limited number of samples on Georges Bank, and no sampling in the Gulf of Maine.

NEFSC spring and autumn survey indices for clearnose skate have been increasing since the mid-1980s (Figure 31 – Figure 33). The minimum length of clearnose skate caught in NEFSC surveys is about 10 cm (4 in), and the largest individual caught was 78 cm (31 in) total length, during the 1971 autumn survey in the Mid-Atlantic Bight region. The median length of the survey catch has ranged from 41 cm in the 1980 spring survey to 67 cm in the 1995 spring survey. The median length of the spring survey catch has increased over the time series, from about 50 cm during the late 1970s to at about 60 cm in recent years (24 in; Figure 34). The median length of the autumn survey catch has been stable over the time series, and is also at about 60 cm. Length frequency distributions from the NEFSC spring and autumn surveys are presented in the SAW 30 documents as figures B92 – B94 and are not reproduced in this SAFE Report. In general, the length frequency distributions show a consistent mode at 60-70 cm that may represent the accumulated abundance of several older ages.

Indices of abundance for clearnose skate are available from the CTDEP spring and autumn finfish trawl surveys in Long Island Sound for the years 1984-1998 (1992 and later only for biomass). The CTDEP survey has caught very few clearnose skate, with annual catches ranging from 0 to 20 skates, although the CTDEP spring survey suggests an increase in clearnose skate abundance in Long Island Sound over the times series (Figure 35).

Indices of abundance for clearnose skate are available from the Virginia Institute of Marine Science (VIMS) trawl survey in Chesapeake Bay and its' tributaries for the years 1988-1998. The VIMS trawl survey indices suggest no trend in clearnose skate abundance over the this period (Figure 36).

Table 15 Abundance and Biomass from NEFSC Spring Surveys for Clearnose Skate for the Mid-Atlantic Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1976-2000.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|--------|-------|------------|--------|-------|--------|--------|----|-----|------|-----|---------|------|---------|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% | mean | 95% | max | tows | no fish |
| 1976 | 0.100 | 0.020 | 0.179 | 0.129 | 0.040 | 0.218 | 0.770 | 26 | 26 | 43 | 48.5 | 66 | 67 | 8 | 12 |
| 1977 | 0.509 | 0.297 | 0.722 | 0.500 | 0.260 | 0.741 | 1.017 | 23 | 23 | 56 | 52.5 | 63 | 64 | 17 | 41 |
| 1978 | 0.211 | -0.094 | 0.516 | 0.237 | -0.057 | 0.530 | 0.893 | 20 | 20 | 57 | 52.2 | 68 | 69 | 8 | 21 |
| 1979 | 0.109 | 0.010 | 0.209 | 0.125 | 0.004 | 0.247 | 0.875 | 25 | 25 | 42 | 50.3 | 77 | 78 | 6 | 9 |
| 1980 | 0.319 | 0.100 | 0.538 | 0.456 | 0.136 | 0.775 | 0.700 | 25 | 25 | 41 | 45.1 | 64 | 69 | 14 | 44 |
| 1981 | 0.891 | -0.141 | 1.923 | 0.606 | 0.106 | 1.107 | 1.469 | 24 | 26 | 60 | 55.9 | 67 | 72 | 10 | 44 |
| 1982 | 0.328 | 0.165 | 0.491 | 0.368 | 0.126 | 0.610 | 0.892 | 30 | 32 | 52 | 53.6 | 66 | 71 | 14 | 40 |
| 1983 | 0.138 | 0.005 | 0.270 | 0.127 | 0.003 | 0.252 | 1.081 | 13 | 13 | 58 | 51.3 | 65 | 66 | 7 | 11 |
| 1984 | 0.380 | 0.103 | 0.658 | 0.288 | 0.018 | 0.557 | 1.321 | 48 | 48 | 62 | 60.7 | 70 | 74 | 11 | 25 |
| 1985 | 0.493 | -0.166 | 1.151 | 0.436 | -0.203 | 1.076 | 1.129 | 48 | 48 | 58 | 59.3 | 69 | 72 | 10 | 37 |
| 1986 | 0.155 | 0.035 | 0.274 | 0.232 | 0.038 | 0.427 | 0.666 | 27 | 27 | 44 | 44.8 | 68 | 69 | 11 | 15 |
| 1987 | 0.306 | 0.150 | 0.463 | 0.202 | 0.109 | 0.204 | 1.519 | 49 | 51 | 63 | 61.9 | 69 | 72 | 16 | 20 |
| 1988 | 0.340 | 0.171 | 0.508 | 0.300 | 0.097 | 0.502 | 1.134 | 44 | 44 | 58 | 57.1 | 67 | 71 | 11 | 19 |
| 1989 | 0.424 | 0.258 | 0.590 | 0.415 | 0.275 | 0.554 | 1.023 | 25 | 25 | 58 | 52.3 | 68 | 72 | 14 | 40 |
| 1990 | 0.501 | 0.283 | 0.719 | 0.420 | 0.243 | 0.597 | 1.192 | 30 | 30 | 59 | 56.2 | 67 | 72 | 15 | 52 |
| 1991 | 0.690 | 0.463 | 0.918 | 0.543 | 0.354 | 0.731 | 1.272 | 27 | 27 | 62 | 58.8 | 68 | 71 | 23 | 59 |
| 1992 | 0.748 | 0.324 | 1.172 | 0.489 | 0.218 | 0.760 | 1.529 | 46 | 46 | 63 | 63.0 | 68 | 80 | 23 | 47 |
| 1993 | 0.856 | 0.479 | 1.233 | 0.656 | 0.216 | 1.096 | 1.305 | 21 | 33 | 63 | 58.6 | 70 | 74 | 12 | 136 |
| 1994 | 0.319 | 0.052 | 0.585 | 0.188 | 0.043 | 0.333 | 1.699 | 51 | 57 | 65 | 66.0 | 73 | 74 | 8 | 24 |
| 1995 | 0.669 | 0.361 | 0.977 | 0.464 | 0.261 | 0.666 | 1.443 | 46 | 46 | 67 | 62.4 | 68 | 74 | 18 | 32 |
| 1996 | 1.224 | 0.194 | 2.254 | 0.948 | 0.255 | 1.641 | 1.291 | 13 | 27 | 62 | 59.8 | 70 | 75 | 30 | 95 |
| 1997 | 1.290 | 0.885 | 1.695 | 0.972 | 0.542 | 1.403 | 1.326 | 33 | 39 | 63 | 61.3 | 71 | 78 | 22 | 80 |
| 1998 | 0.903 | 0.674 | 1.133 | 0.667 | 0.369 | 0.964 | 1.355 | 26 | 38 | 62 | 60.2 | 70 | 74 | 29 | 81 |
| 1999 | 0.943 | 0.647 | 1.238 | 0.862 | 0.470 | 1.255 | 1.093 | 26 | 28 | 59 | 57.3 | 67 | 72 | 19 | 54 |
| 2000 | 1.391 | 1.046 | 1.736 | 1.140 | 0.789 | 1.491 | 1.221 | 24 | 40 | 59 | 59.4 | 70 | 76 | 31 | 126 |

Table 16 Abundance and Biomass from NEFSC Autumn Surveys for Clearnose Skate for the Mid-Atlantic Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1975-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|--------|-------|------------|-------|-------|--------|--------|----|-----|------|-----|---------|------|---------|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% | mean | 95% | max | tows | no fish |
| 1975 | 0.237 | 0.086 | 0.388 | 0.246 | 0.133 | 0.360 | 0.961 | 21 | 21 | 53 | 50.3 | 63 | 66 | 31 | 49 |
| 1976 | 0.302 | 0.189 | 0.415 | 0.348 | 0.236 | 0.459 | 0.869 | 18 | 34 | 52 | 52.1 | 64 | 69 | 26 | 54 |
| 1977 | 0.768 | 0.288 | 1.248 | 0.742 | 0.281 | 1.203 | 1.035 | 15 | 37 | 57 | 55.4 | 65 | 68 | 32 | 106 |
| 1978 | 0.156 | 0.073 | 0.240 | 0.224 | 0.086 | 0.363 | 0.697 | 10 | 10 | 44 | 40.8 | 64 | 66 | 14 | 23 |
| 1979 | 0.419 | 0.116 | 0.721 | 0.346 | 0.146 | 0.545 | 1.211 | 22 | 24 | 56 | 55.4 | 67 | 71 | 27 | 46 |
| 1980 | 0.685 | 0.408 | 0.961 | 0.549 | 0.322 | 0.775 | 1.248 | 33 | 37 | 59 | 58.1 | 69 | 72 | 32 | 80 |
| 1981 | 0.171 | 0.081 | 0.260 | 0.179 | 0.087 | 0.271 | 0.954 | 27 | 27 | 55 | 51.5 | 65 | 68 | 19 | 28 |
| 1982 | 0.213 | 0.099 | 0.326 | 0.183 | 0.095 | 0.271 | 1.163 | 32 | 43 | 59 | 58.3 | 67 | 72 | 26 | 37 |
| 1983 | 0.141 | 0.027 | 0.254 | 0.127 | 0.043 | 0.210 | 1.110 | 16 | 16 | 57 | 52.2 | 64 | 70 | 15 | 19 |
| 1984 | 0.178 | 0.064 | 0.293 | 0.189 | 0.063 | 0.315 | 0.945 | 34 | 37 | 53 | 54.0 | 67 | 83 | 20 | 32 |
| 1985 | 0.306 | 0.173 | 0.439 | 0.315 | 0.182 | 0.447 | 0.974 | 32 | 41 | 56 | 54.9 | 66 | 71 | 23 | 42 |
| 1986 | 0.545 | -0.038 | 1.027 | 0.591 | 0.091 | 1.092 | 0.921 | 23 | 23 | 59 | 52.6 | 64 | 71 | 31 | 62 |
| 1987 | 0.320 | 0.176 | 0.465 | 0.289 | 0.167 | 0.412 | 1.107 | 15 | 41 | 56 | 55.5 | 69 | 70 | 23 | 42 |
| 1988 | 0.335 | 0.157 | 0.513 | 0.329 | 0.163 | 0.495 | 1.019 | 33 | 37 | 57 | 56.0 | 66 | 71 | 19 | 60 |
| 1989 | 0.273 | 0.075 | 0.471 | 0.324 | 0.064 | 0.584 | 0.843 | 37 | 37 | 52 | 52.7 | 63 | 70 | 20 | 39 |
| 1990 | 0.402 | 0.157 | 0.646 | 0.306 | 0.114 | 0.499 | 1.311 | 16 | 41 | 60 | 57.9 | 69 | 72 | 17 | 50 |
| 1991 | 0.922 | 0.279 | 1.566 | 0.816 | 0.339 | 1.294 | 1.130 | 35 | 39 | 58 | 57.1 | 69 | 71 | 35 | 119 |
| 1992 | 0.345 | 0.185 | 0.505 | 0.312 | 0.185 | 0.440 | 1.104 | 16 | 42 | 59 | 56.7 | 67 | 69 | 22 | 48 |
| 1993 | 0.495 | 0.145 | 0.844 | 0.474 | 0.188 | 0.759 | 1.044 | 35 | 40 | 57 | 56.8 | 66 | 73 | 27 | 104 |
| 1994 | 0.938 | 0.479 | 1.398 | 0.842 | 0.494 | 1.190 | 1.115 | 35 | 40 | 57 | 57.1 | 66 | 73 | 35 | 129 |
| 1995 | 0.331 | 0.189 | 0.473 | 0.426 | 0.233 | 0.618 | 0.777 | 14 | 14 | 51 | 45.5 | 66 | 72 | 25 | 63 |
| 1996 | 0.430 | 0.194 | 0.666 | 0.369 | 0.163 | 0.576 | 1.165 | 29 | 45 | 59 | 58.8 | 68 | 72 | 20 | 42 |
| 1997 | 0.614 | 0.296 | 0.932 | 0.484 | 0.281 | 0.688 | 1.269 | 43 | 43 | 61 | 60.2 | 69 | 77 | 27 | 60 |
| 1998 | 1.121 | 0.115 | 2.128 | 1.096 | 0.124 | 2.068 | 1.023 | 34 | 43 | 57 | 57.5 | 68 | 73 | 32 | 98 |
| 1999 | 1.053 | 0.536 | 1.570 | 0.928 | 0.525 | 1.332 | 1.134 | 15 | 32 | 61 | 57.8 | 69 | 71 | 41 | 84 |

Table 17 Abundance and Biomass from NEFSC Winter Surveys for Clearnose Skate for the Georges Bank to Mid-Atlantic Region

The mean index, 95% confidence intervals, individual fish weight, minimum, mean, and maximum length, 5th, 50th, and 95th percentiles of length, number of nonzero tows, and number of fish caught are presented for 1992-1999.

| | weight/tow | | | number/tow | | | ind wt | length | | | | | nonzero | | |
|------|------------|-------|--------|------------|-------|--------|--------|--------|----|-----|------|-----|---------|------|---------|
| | mean | lower | upper | mean | lower | upper | | min | 5% | 50% | mean | 95% | max | tows | no fish |
| 1992 | 5.622 | 3.247 | 7.997 | 5.247 | 2.974 | 7.519 | 1.072 | 23 | 26 | 59 | 54.7 | 67 | 93 | 22 | 551 |
| 1993 | 6.013 | 3.818 | 8.208 | 5.973 | 3.852 | 8.093 | 1.007 | 22 | 33 | 57 | 54.3 | 67 | 81 | 23 | 716 |
| 1994 | 8.854 | 4.037 | 13.672 | 7.692 | 2.152 | 13.233 | 1.151 | 27 | 33 | 60 | 57.5 | 69 | 77 | 16 | 639 |
| 1995 | 7.924 | 2.521 | 13.327 | 6.247 | 1.301 | 11.194 | 1.268 | 24 | 45 | 61 | 60.2 | 69 | 76 | 23 | 737 |
| 1996 | 14.725 | 8.266 | 21.183 | 11.555 | 6.347 | 16.762 | 1.274 | 22 | 40 | 61 | 60.0 | 69 | 77 | 32 | 3086 |
| 1997 | 5.522 | 3.154 | 7.890 | 5.069 | 2.158 | 7.980 | 1.089 | 22 | 35 | 59 | 56.2 | 70 | 76 | 32 | 682 |
| 1998 | 6.031 | 4.470 | 7.592 | 4.878 | 3.195 | 6.560 | 1.236 | 22 | 36 | 60 | 58.3 | 71 | 88 | 32 | 1091 |
| 1999 | 3.826 | 2.335 | 5.317 | 3.022 | 1.586 | 4.459 | 1.266 | 23 | 37 | 61 | 59.6 | 70 | 76 | 30 | 343 |
| 2000 | 10.102 | 5.693 | 14.510 | 8.864 | 4.579 | 13.150 | 1.140 | 25 | 42 | 59 | 58.2 | 69 | 93 | 43 | 1449 |

Figure 31 Abundance and Biomass of Clearnose Skate from the NEFSC Spring (Circles) and Autumn (Squares) Bottom Trawl Surveys from 1975-2000 in the Mid-Atlantic Offshore and Inshore Regions

**Clearnose Skate
Mid-Atlantic All strata**




Figure 32 Abundance and Biomass of Clearnose Skate from the NEFSC Spring Bottom Trawl Survey in the Mid-Atlantic Region, Offshore and Inshore Regions

Mean Index In Solid Squares, 95% Confidence Interval In Open Squares

