


Monkfish-L mesh gillnet; January


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; January

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

JAN_MNK_LGNS

1

2

3 - 12

13 - 20


21 - 44

Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; January


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; January

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

JAN_MNK_XGNS

1 - 4

5 - 11

12 - 18

19 - 27


28 - 94

Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; January



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; January

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; February

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; February


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch


MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; February

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-XL mesh gillnet; February

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

FEB_MNK_XGNS

1-2

3-5

6-8

9-31

32-81

Atlantic sturgeon bycatch

MONTH

1

2

3

4

5

6

7

8


9

10

11

12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; February



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; February

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; March

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-L mesh gillnet; March

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; March

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
- TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-XL mesh gillnet; March

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-XL mesh gillnet; March


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch



MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; March

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-L mesh gillnet; April

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

APR_MNK_LGNS

2

3

4

5 - 15

16 - 81

Atlantic sturgeon bycatch

MONTH

1

2

3

4

5

6

7

8

9


10

11

12

● TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-L mesh gillnet; April


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; April

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; April

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-XL mesh gillnet; April


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; April


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; May

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish


Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-L mesh gillnet; May


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; May

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; May

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.



Monkfish-XL mesh gillnet; May


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; May

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; June

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; June


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; June

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; June


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; June

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12




TripTarget1=Monkfish

Statistical_Areas

JUN_MNK_XGNS

- 1 - 5
- 6 - 18
- 19 - 26
- 27 - 60
- 61 - 119




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; June

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; July

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; July


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; July

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

TripTarget1=Monkfish



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; July

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

JUL_MNK_XGNS

1 - 5

6 - 12

13 - 19

20 - 33


34 - 83

Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish



Monkfish-XL mesh gillnet; July

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

JUL_MNK_XGNS

1 - 5

6 - 12

13 - 19

20 - 33

34 - 83

Atlantic sturgeon bycatch

MONTH

1

2

3

4

5

6

7

8


9

10

11

12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; July

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; August

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12



TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; August


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; August

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12



TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; August

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; August

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12



TripTarget1=Monkfish



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; August

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0

Statistical_Areas

AUG_MNK_XGNS

1

2 - 3

4 - 13

14 - 57

58 - 80


Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12



TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; September


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; September


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; September

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; September

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; September


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010

0


Statistical_Areas

SEP_MNK_XGNS



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; September

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; October

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; October

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; October

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; October

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; October



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; October

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; November

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010




Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12

● TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; November


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; November

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; November



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; November



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; November


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11 
- 12
-  TripTarget1=Monkfish



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; December

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; December



To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-L mesh gillnet; December

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; December

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; December


To Read Labels:
567 << Stat Area ID
123 << Average Days Absent

Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH




Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.

Monkfish-XL mesh gillnet; December

To Read Labels:
567 << Stat Area ID
123 << Average Days Absent


Average Annual
Time Absent in
Days, 2006-2010



Atlantic sturgeon bycatch

MONTH

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- TripTarget1=Monkfish



Effort data (shaded statistical areas) were derived from VTR records and are an average of data from 2006-2010. The effort is expressed as the average annual time vessels were absent from port and reported catching the indicated species on the VTR. All sturgeon records were obtained from the NEFOP + ASM database and represent observations from 2006-2010. These sturgeon records were only filtered by month where indicated (hollow points and blue points) and the declared trip target (blue points only). Thus, the sturgeon records will be identical between months, with changes to the blue points based on the target species identified, not by gear type or mesh size.