

#3



New England Fishery Management Council

50 WATER STREET | NEWBURYPORT, MASSACHUSETTS 01950 | PHONE 978 465 0492 | FAX 978 465 3116

John Pappalardo, *Chairman* | Paul J. Howard, *Executive Director*

MEMORANDUM

DATE: January 14, 2008
TO: Science and Statistical Committee
FROM: Skate PDT
SUBJECT: Skate rebuilding catch limit re-analysis

This analysis incorporates the new Data Poor Assessment Workshop¹ skate catch time series into the previous PDT evaluation of skate rebuilding potential. While total landings were updated and new methods to allocate unclassified skate landings to species were developed in the DPWS, new discard estimates were completely revised using observer data which had not previously been included. As a result, the re-assignment of catches to skate species were revised and total discard estimates are substantially different than previous data used in the Draft Amendment 3 analysis.

Like the previous assessment, the new analyses evaluate the relationship between catch, relative exploitation (catch/biomass) and changes in stratified mean biomass estimated by the surveys (spring for little skate, fall for the remaining six species). To smooth out noise from annual indices, a 3-year moving average for catch and biomass with no lags was evaluated². Based on this type of analysis, the PDT recommended and the SSC approved using the median relative exploitation ratio (C/B) applied to the latest three year stratified mean biomass as an interim catch limit to initiate rebuilding of smooth, thorny, and winter skates.

The median values (2005-2007) for each species were summed and applied as an aggregate skate ABC/ACT, accounting for the partial effectiveness (assumed 90%) of barndoor, smooth, and thorny skate landings prohibitions. A value of 75% of the threshold catch limit was recommended to account for scientific and management uncertainty, approved for a management target, and applied as an ACT in the Draft Amendment 3 document. The average discard rate for 2004-2006 and two different historic landings splits between the skate wing and bait fisheries was then applied to estimate TALs for each fishery. The same procedure was applied in this analysis, except that the 2005-2007 discard rate was

¹ A Data Poor Assessment Workshop (DPWS) was conducted by the Northeast Fisheries Science Center during October 2008 to January 2009, focusing on exploratory assessment analyses of model-resistant species, including the seven managed skate stocks. While the survey time series is believed to be a good representation of changes in skate abundance and biomass, there has been considerable uncertainty in the skate species landings and in discard estimates. One of the important outcomes of the DPWS was two methods to allocate skate catches to species based on where the fishing activity occurred and the observed lengths of skate catches.

² Other lags and moving average durations were evaluated in the Draft Amendment 3 technical analyses and were not informative, i.e. correlations between catch or relative exploitation and biomass changes were worse.

applied, now that 2007 discard estimates have been calculated in the DPWS. A summary of comparative results are given in the table below.

Data source	Catch limit, mt (ABC/ACL)	Catch target, mt (ACT)	Discard rate	Total allowable landings, mt (TAL)	MSY (landings with biomass @ target)	Landings reduction from 2007 to achieve TAL	
						Wing	Bait
Draft Amendment 3	22,612	16,959	38%	10,484	53,731	-45%	-43%
DPWS Length composition method.	24,688	15,546	58%	7,786	64,196	-57%	-63%
DPWS selectivity ogive method	23,826	17,864	59%	7,328	63,240	-65%	-60%

Although they were initially different and derived independently, the two DPWS method catch series have become similar with refinement. As indicated above, there really is little difference between them in the context of the Amendment 3 rebuilding prospects at catch rates below and above the median values for the time series. Even the overall catch limit (landings and discards) are similar to the Draft Amendment 3 results, but the higher discard estimates result in a lower fraction (41-42%) of the total catch being allocated to landings (i.e. TAL).

Analysis of rebuilding potential, however, shows that the linkage between low exploitation rates and increases in biomass is either non-existent or not significant. None of the relationships are very strong and are probably not very predictive of rebuilding potential at lower catch levels. There is little or no relationship between the C/B ratio and changes in biomass for barndoor, clearnose, little, or thorny skates. The relationship for smooth and rosette seem entirely attributable to a few number of points which may be related to transient oceanographic events or sampling variability, while the relationship for winter seems to be related to serial autocorrelation.

Catch time series

For the Amendment 3 DEIS, the PDT estimated landings species composition by applying the survey biomass proportions for exploitable skates in each three-digit statistical area, as determined by a fitted logistic selectivity curve (fitting observed commercial kept skates to the survey in equivalent areas and seasons) of observed kept skate lengths on survey length frequencies in each region and season. Although known at the time, this procedure had a technical flaw and inconsistency with the survey design, but was not thought to significantly skew the species allocations. During the DEIS comment period, NMFS commented on this flaw in the analysis and it would be addressed in the DPWS³. Although the Council was slated to take final action at the November 2008 meeting, NMFS recommended that the Council wait to receive these results to determine whether to proceed with Amendment 3.

³ Analyses were presented at the DPWS that the previous Amendment 3 assumption did not badly violate the survey statistical design and did not skew the biomass proportions or the calculated mean biomass of each species in a statistical area.

During the Amendment 3 development, the PDT also only had regional estimates (Georges Bank/Southern New England and Mid-Atlantic) of aggregate skate discards to use in the Amendment 3 analyses. These discard estimates used SAW44-reviewed procedures, but used the Groundfish Assessment Review Meeting (GARM) area allocation tables⁴ to assign landings to statistical area and region. Because species composition of discards was not available at the time, the PDT used the regional skate discard estimates as a catch index for species by region (Georges Bank/Southern New England for thorny, smooth, winter, and little skates; Mid-Atlantic for clearnose and rosette skates).

The new catch series for this analysis allocate skate landings and discards to species based on surveyed biomass fractions using two different methods. These two methods were developed simultaneously, and independently arrived at similar results to one another. The details are described in the DPWS documents, but are summarized below. Each method has its pros and cons and both methods were accepted by the DPWS.

For the length composition method, the skate lengths of kept and discarded skates were binned into 5 cm intervals and applied to the survey biomass fractions by region. These biomass fractions were applied to total landings and total discard estimates by year, half-year, gear, and region (Gulf of Maine, Southern New England, and Mid-Atlantic). Discard to kept ratios were applied to total landings on all trips, also by year, half-year, gear, and region. The discard species composition was calculated in the same fashion as that for landings, using the length composition of discarded skates on observed trips. For both landings and discards, the species composition could only be determined since 1989, the first year of sea sampling data. Total discards were however hind-casted by applying the 1989-1991 DK ratio to dealer reported landings in earlier years.

For the selectivity ogive method of assigning species composition to skate landings and discards, the fraction of skate biomass for skate species were determined by estimating a selectivity ogive from kept skates on observed trips during 2004-2006 applied to surveyed skates in each three digit statistical area by year, gear, and season (spring, fall, and winter corresponding to the three trawl surveys). The survey biomass fractions were determined by applying the 2004-2006 selectivity ogive by year, season, sub-region, gear, and mesh (small, large, and extra-large for trawls and gillnets). Subsequent to the DPWS, these fractions were re-estimated by treating the fractions kept for vessels using gillnets separately for the skate wing and whole/bait fisheries⁵. This is the same procedure as the DPWS approved for trawls and recommended that it be used for the gillnet fishery as well.

Total discards were independently estimated for the DPWS by estimating the DK ratios⁶ for each year, gear, sub-region, season, and mesh and applying them to total landings on every trip reported by dealers and included in the GARM area allocation tables. Trips in these tables with unknown area allocations were distributed to areas, based on fishing activity for assigned trips in each state, year, and gear combination. Skate species allocations were made using the same procedure that the DPWS accepted for landings, using the selectivity ogive method, except that for trips landing skates the species composition of discards were determined by $A - B$, where A is the selectivity ogive fitted for catch and B is the selectivity ogive fitted for kept skates. For trips with no skate landings in the dealer data, only the

⁴ These area allocation tables use a peer-reviewed method to allocate dealer reported landings to statistical area level fishing locations.

⁵ The DPWS estimates use a single selectivity ogive for all skate landings by vessels using gillnets and it was discovered during the review that a substantial fraction of gillnet landings are landed in whole form, presumably targeting little skates for bait.

⁶ Skates discarded to total live weight of landings of all species on observed trips, which are then applied to total live weight of dealer reported landings of all species.

selectivity ogive for catch (A) was used to assign species composition to skate discards. Since the method uses the length distribution of skates in the survey to determine species composition, the species allocations could be assigned as far back as 1977.

This analysis of rebuilding potential described below uses the species composition of 1994-2007 landings and discards from each species allocation method independently to examine the effect of catch on changes in survey skate biomass. For 1977-1993, the total skate landings and both discard estimates were assigned the species composition determined by the selectivity ogive method (because there were no observed trips before 1989 and no GARM area allocation tables before 1994). Before the advent of the sea sampling program in 1989, both methods apply the 1989-1991 DK ratio to total landings. The data used in this analysis are shown in the following tables.

Table 1. Landings time series used in the rebuilding potential analysis and in estimating catch limits and targets associated with the median C/B exploitation ratio. Landings before 1994 were derived from the same time series of aggregate landings and species composition was assigned via the selectivity ogive method.

Length composition method												
Landings Allocations												
Year	barndoor	chamose	littl	rossette	smooth	thorny	winter	barndoor	chamose	littl	rossette	winter
1984	4.52	0.00	1.14	0.03	0.29	20.05	4.57	4.52	0.00	1.14	0.03	0.29
1985	5.78	0.00	1.87	0.02	0.44	25.03	5.46	5.78	0.00	1.87	0.02	0.44
1986	3.41	0.00	1.26	0.00	0.20	21.49	4.54	3.41	0.00	1.26	0.00	0.20
1987	3.68	1.23	5.48	0.12	0.36	42.38	18.24	3.68	1.23	5.48	0.12	0.36
1988	2.62	1.35	2.45	0.00	0.40	20.84	7.83	2.62	1.35	2.45	0.00	0.40
1989	2.62	1.30	3.54	0.01	0.48	32.43	11.21	2.62	1.30	3.54	0.01	0.48
1990	1.95	0.47	4.79	0.03	0.52	46.86	14.97	1.95	0.47	4.79	0.03	0.52
1991	0.78	0.64	5.24	0.05	0.70	37.77	17.92	0.78	0.64	5.24	0.05	0.70
1992	1.61	0.00	6.05	0.00	0.94	51.57	25.93	1.61	0.00	6.05	0.00	0.94
1993	1.43	0.71	6.82	0.03	0.77	47.47	29.67	1.43	0.71	6.82	0.03	0.77
1994	1.19	0.75	6.42	0.03	0.66	40.68	30.37	1.19	0.75	6.42	0.03	0.66
1995	1.51	1.20	9.34	0.04	0.87	53.56	47.58	1.51	1.20	9.34	0.04	0.87
1996	1.30	1.31	9.37	0.04	0.79	48.36	51.03	1.30	1.31	9.37	0.04	0.79
1997	1.47	1.91	12.64	0.05	0.96	58.22	73.05	1.47	1.91	12.64	0.05	0.96
1998	2.51	4.26	26.30	0.10	1.78	107.16	160.49	2.51	4.26	26.30	0.10	1.78
1999	3.27	7.52	43.61	0.15	2.80	155.39	279.66	3.27	7.52	43.61	0.15	2.80
2000	3.08	10.14	55.63	0.19	2.89	170.84	373.34	3.08	10.14	55.63	0.19	2.89
2001	1.10	8.80	30.21	0.09	1.34	78.44	211.42	1.10	8.80	30.21	0.09	1.34
2002	0.02	8.68	50.86	0.21	1.14	88.77	279.02	0.02	8.68	50.86	0.21	1.14
2003	0.00	11.38	76.64	0.01	3.42	124.33	666.91	0.00	11.38	76.64	0.01	3.42
2004	0.28	22.71	69.73	0.28	1.59	114.00	822.20	0.28	22.71	69.73	0.28	1.59
2005	0.06	12.71	51.44	0.08	2.32	72.72	623.98	0.06	12.71	51.44	0.08	2.32
2006	0.36	13.84	30.73	0.10	4.26	78.53	858.28	0.36	13.84	30.73	0.10	4.26
2007	0.32	40.34	84.38	0.26	4.52	107.37	1202.52	0.32	40.34	84.38	0.26	4.52
1984	0.01	63.95	99.07	0.51	10.73	163.98	1775.47	0.01	63.95	99.07	0.51	10.73
1985	1.03	112.20	550.87	0.70	27.51	692.92	5322.07	1.03	112.20	550.87	0.70	27.51
1986	14.39	322.67	830.97	1.07	65.07	859.75	9308.58	14.39	322.67	830.97	1.07	65.07
1987	16.47	983.65	1332.93	3.82	51.05	1173.66	7770.72	16.47	983.65	1332.93	3.82	51.05
1988	471.60	746.18	1379.39	4.11	77.86	2089.16	7757.00	471.60	746.18	1379.39	4.11	77.86
1989	70.90	1054.90	2915.57	2.20	117.38	1581.75	7161.30	70.90	1054.90	2915.57	2.20	117.38
1990	134.20	973.71	1794.69	6.62	89.99	1966.44	3818.55	134.20	973.71	1794.69	6.62	89.99
1991	83.11	346.48	1626.66	6.39	0.77	314.57	4453.06	83.11	346.48	1626.66	6.39	0.77
1992	336.39	539.89	2398.89	11.01	0.37	759.51	10031.54	336.39	539.89	2398.89	11.01	0.37
1993	281.04	748.73	3792.04	12.90	6.89	510.38	5355.70	281.04	748.73	3792.04	12.90	6.89
1994	161.12	447.45	4028.73	27.33	7.83	628.19	8344.25	161.12	447.45	4028.73	27.33	7.83
1995	452.37	324.36	3980.41	15.35	2.09	203.71	6966.57	452.37	324.36	3980.41	15.35	2.09
1996	494.42	501.95	3390.02	19.96	7.67	468.39	8372.99	494.42	501.95	3390.02	19.96	7.67
2000	1538.85	1860.07	1700.99	5.61	18.78	195.42	7855.28	1538.85	1860.07	1700.99	5.61	18.78
2001	2123.66	640.20	2371.81	10.72	17.24	401.63	9986.12	2123.66	640.20	2371.81	10.72	17.24
2002	854.82	335.61	3302.87	5.82	8.55	302.94	7099.12	854.82	335.61	3302.87	5.82	8.55
2003	844.52	344.54	1955.26	6.80	5.63	511.56	11787.82	844.52	344.54	1955.26	6.80	5.63
2004	1976.34	168.47	3056.36	8.97	10.39	439.86	7650.58	1976.34	168.47	3056.36	8.97	10.39
2005	2632.83	384.49	2392.33	8.63	21.51	642.87	8256.81	2632.83	384.49	2392.33	8.63	21.51
2006	2011.46	361.73	3078.31	22.41	17.84	351.91	12860.80	2011.46	361.73	3078.31	22.41	17.84
2007												
1985-2007	7.9%	4.0%	21.3%	0.1%	0.1%	3.3%	63.2%	7.9%	4.0%	21.3%	0.1%	0.1%

Selectivity ogive method												
Landings Allocations												
Year	barndoor	chamose	littl	rossette	smooth	thorny	winter	barndoor	chamose	littl	rossette	winter
1984	4.52	0.00	1.14	0.03	0.29	20.05	4.57	4.52	0.00	1.14	0.03	0.29
1985	5.78	0.00	1.87	0.02	0.44	25.03	5.46	5.78	0.00	1.87	0.02	0.44
1986	3.41	0.00	1.26	0.00	0.20	21.49	4.54	3.41	0.00	1.26	0.00	0.20
1987	3.68	1.23	5.48	0.12	0.36	42.38	18.24	3.68	1.23	5.48	0.12	0.36
1988	2.62	1.35	2.45	0.00	0.40	20.84	7.83	2.62	1.35	2.45	0.00	0.40
1989	2.62	1.30	3.54	0.01	0.48	32.43	11.21	2.62	1.30	3.54	0.01	0.48
1990	1.95	0.47	4.79	0.03	0.52	46.86	14.97	1.95	0.47	4.79	0.03	0.52
1991	0.78	0.64	5.24	0.05	0.70	37.77	17.92	0.78	0.64	5.24	0.05	0.70
1992	1.61	0.00	6.05	0.00	0.94	51.57	25.93	1.61	0.00	6.05	0.00	0.94
1993	1.43	0.71	6.82	0.03	0.77	47.47	29.67	1.43	0.71	6.82	0.03	0.77
1994	1.19	0.75	6.42	0.03	0.66	40.68	30.37	1.19	0.75	6.42	0.03	0.66
1995	1.51	1.20	9.34	0.04	0.87	53.56	47.58	1.51	1.20	9.34	0.04	0.87
1996	1.30	1.31	9.37	0.04	0.79	48.36	51.03	1.30	1.31	9.37	0.04	0.79
1997	1.47	1.91	12.64	0.05	0.96	58.22	73.05	1.47	1.91	12.64	0.05	0.96
1998	2.51	4.26	26.30	0.10	1.78	107.16	160.49	2.51	4.26	26.30	0.10	1.78
1999	3.27	7.52	43.61	0.15	2.80	155.39	279.66	3.27	7.52	43.61	0.15	2.80
2000	3.08	10.14	55.63	0.19	2.89	170.84	373.34	3.08	10.14	55.63	0.19	2.89
2001	1.10	8.80	30.21	0.09	1.34	78.44	211.42	1.10	8.80	30.21	0.09	1.34
2002	0.02	8.68	50.86	0.21	1.14	88.77	279.02	0.02	8.68	50.86	0.21	1.14
2003	0.00	11.38	76.64	0.01	3.42	124.33	666.91	0.00	11.38	76.64	0.01	3.42
2004	0.28	22.71	69.73	0.28	1.59	114.00	822.20	0.28	22.71	69.73	0.28	1.59
2005	0.06	12.71	51.44	0.08	2.32	72.72	623.98	0.06	12.71	51.44	0.08	2.32
2006	0.36	13.84	30.73	0.10	4.26	78.53	858.28	0.36	13.84	30.73	0.10	4.26
2007	0.32	40.34	84.38	0.26	4.52	107.37	1202.52	0.32	40.34	84.38	0.26	4.52
1984	0.01	63.95	99.07	0.51	10.73	163.98	1775.47	0.01	63.95	99.07	0.51	10.73
1985	1.03	112.20	550.87	0.70	27.51	692.92	5322.07	1.03	112.20	550.87	0.70	27.51
1986	14.39	322.67	830.97	1.07	65.07	859.75	9308.58	14.39	322.67	830.97	1.07	65.07
1987	16.47	983.65	1332.93	3.82	51.05	1173.66	7770.72	16.47	983.65	1332.93	3.82	51.05
1988	471.60	746.18	1379.39	4.11	77.86	2089.16	7757.00	471.60	746.18	1379.39	4.11	77.86
1989	70.90	1054.90	2915.57	2.20	117.38	1581.75	7161.30	70.90	1054.90	2915.57	2.20	117.38
1990	134.20	973.71	1794.69	6.62	89.99	1966.44	3818.55	134.20	973.71	1794.69	6.62	89.99
1991	83.11	346.48	1626.66	6.39	0.77	314.57	4453.06	83.11	346.48	1626.66	6.39	0.77
1992	336.39	539.89	2398.89	11.01	0.37	759.51	10031.54	336.39	539.89	2398.89	11.01	0.37
1993	281.04	748.73	3792.04	12.90	6.89	510.38	5355.70	281.04	748.73	3792.04	12.90	6.89
1994	161.12	447.45	4028.73	27.33	7.83	628.19	8344.25	161.12	447.45	4028.73	27.33	7.83
1995	452.37	324.36	3980.41	15.35	2.09	203.71	6966.57	452.37	324.36	3980.41	15.35	2.09
1996	494.42	501.95	3390.02	19.96	7.67	468.39	8372.99	494.42	501.95	3390.02	19.96	7.67
2000	1538.85	1860.07	1700.99	5.61	18.78	195.42	7855.28	1538.85	1860.07	1700.99	5.61	18.78
2001	2123.66	640.20	2371.81	10.72	17.24	401.63	9986.12	2123.66	640.20	2371.81	10.72	17.24
2002	854.82	335.61	3302.87	5.82	8.55	302.94	7099.12	854.82	335.61	3302.87	5.82	8.55
2003	844.52	344.54	1955.26	6.80	5.63	511.56	11787.82	844.52	344.54	1955.26	6.80	5.63
2004	1976.34	168.47	3056.36	8.97	10.39	439.86	7650.58	1976.34	168.47	3056.36	8.97	10.39
2005	2632.83	384.49	2392.33	8.63	21.51	642.87	8256.81	2632.83	384.49	2392.33	8.63	21.51
2006	2011.46	361.73	3078.31	22.41	17.84	351.91	12860.80	2011.46	361.7			

Table 2. Discard time series used in the rebuilding potential analysis and in estimating catch limits and targets associated with the median C/B exploitation ratio. Discards before 1993 were derived from the same source using DK ratios from the DPWS and the species composition using the selectivity ogive method.

Length composition method										Selectivity ogive method										Draft Amendment 3 Discards Allocations									
Year	Bamdoor	Clearnose Little	Rosette	Smooth	Thorny	Winter	Year	Bamdoor	Clearnose Little	Rosette	Smooth	Thorny	Winter	Year	Bamdoor	Clearnose Little	Rosette	Smooth	Thorny	Winter									
1964	13,820	-	3,422	168	1,551	61,225	1964	13,820	-	3,422	168	1,551	61,225	1964	13,820	-	3,422	168	1,551	61,225	17,508								
1965	14,511	-	12,760	270	2,074	65,717	1965	14,511	-	12,760	270	2,074	65,717	1965	14,511	-	12,760	270	2,074	65,717	16,393								
1966	10,627	-	16,659	1,100	1,113	70,109	1966	10,627	-	16,659	1,100	1,113	70,109	1966	10,627	-	16,659	1,100	1,113	70,109	16,902								
1967	4,146	2,302	16,899	1,000	1,003	48,940	1967	4,146	2,302	16,899	1,000	1,003	48,940	1967	4,146	2,302	16,899	1,000	1,003	48,940	23,819								
1968	5,752	3,702	16,402	80	1,525	49,761	1968	5,752	3,702	16,402	80	1,525	49,761	1968	5,752	3,702	16,402	80	1,525	49,761	19,941								
1969	3,654	2,127	14,621	113	1,166	49,507	1969	3,654	2,127	14,621	113	1,166	49,507	1969	3,654	2,127	14,621	113	1,166	49,507	17,934								
1970	1,704	720	11,916	129	816	44,796	1970	1,704	720	11,916	129	816	44,796	1970	1,704	720	11,916	129	816	44,796	14,752								
1971	708	526	12,497	45	1,157	30,091	1971	708	526	12,497	45	1,157	30,091	1971	708	526	12,497	45	1,157	30,091	15,233								
1972	845	-	9,328	-	732	29,592	1972	845	-	9,328	-	732	29,592	1972	845	-	9,328	-	732	29,592	16,374								
1973	825	498	11,256	53	751	28,273	1973	825	498	11,256	53	751	28,273	1973	825	498	11,256	53	751	28,273	16,196								
1974	739	539	11,467	53	698	26,139	1974	739	539	11,467	53	698	26,139	1974	739	539	11,467	53	698	26,139	19,910								
1975	559	522	9,979	45	552	20,530	1975	559	522	9,979	45	552	20,530	1975	559	522	9,979	45	552	20,530	18,464								
1976	529	619	11,031	48	551	20,357	1976	529	619	11,031	48	551	20,357	1976	529	619	11,031	48	551	20,357	21,615								
1977	536	794	13,315	57	597	21,869	1977	536	794	13,315	57	597	21,869	1977	536	794	13,315	57	597	21,869	27,478								
1978	553	1,062	16,834	71	673	24,380	1978	553	1,062	16,834	71	673	24,380	1978	553	1,062	16,834	71	673	24,380	36,423								
1979	489	1,259	18,960	78	669	23,929	1979	489	1,259	18,960	78	669	23,929	1979	489	1,259	18,960	78	669	23,929	42,838								
1980	375	1,372	19,726	80	607	21,374	1980	375	1,372	19,726	80	607	21,374	1980	375	1,372	19,726	80	607	21,374	46,381								
1981	252	1,467	20,196	80	534	18,421	1981	252	1,467	20,196	80	534	18,421	1981	252	1,467	20,196	80	534	18,421	49,266								
1982	6	1,814	27,399	151	365	17,074	1982	6	1,814	27,399	151	365	17,074	1982	6	1,814	27,399	151	365	17,074	52,965								
1983	-	1,175	22,731	4	622	12,738	1983	-	1,175	22,731	4	622	12,738	1983	-	1,175	22,731	4	622	12,738	64,438								
1984	27	2,400	20,160	124	259	11,205	1984	27	2,400	20,160	124	259	11,205	1984	27	2,400	20,160	124	259	11,205	63,077								
1985	6	1,304	13,956	33	343	7,026	1985	6	1,304	13,956	33	343	7,026	1985	6	1,304	13,956	33	343	7,026	56,648								
1986	32	1,231	7,755	37	540	6,826	1986	32	1,231	7,755	37	540	6,826	1986	32	1,231	7,755	37	540	6,826	65,596								
1987	25	2,459	15,778	61	412	6,454	1987	25	2,459	15,778	61	412	6,454	1987	25	2,459	15,778	61	412	6,454	64,335								
1988	2	2,841	11,538	118	732	8,856	1988	2	2,841	11,538	118	732	8,856	1988	2	2,841	11,538	118	732	8,856	67,492								
1989	15	1,859	22,280	59	550	8,852	1989	15	1,859	22,280	59	550	8,852	1989	15	1,859	22,280	59	550	8,852	67,492								
1990	129	3,595	26,349	47	1,016	9,050	1990	129	3,595	26,349	47	1,016	9,050	1990	129	3,595	26,349	47	1,016	9,050	85,647								
1991	104	6,033	27,316	119	536	7,561	1991	104	6,033	27,316	119	536	7,561	1991	104	6,033	27,316	119	536	7,561	47,670								
1992	1,766	3,371	18,290	90	577	9,299	1992	1,766	3,371	18,290	90	577	9,299	1992	1,766	3,371	18,290	90	577	9,299	34,270								
1993	178	3,023	24,196	23	546	4,371	1993	178	3,023	24,196	23	546	4,371	1993	178	3,023	24,196	23	546	4,371	21,616								
1994	871	6,956	40,319	204	1,037	14,161	1994	871	6,956	40,319	204	1,037	14,161	1994	871	6,956	40,319	204	1,037	14,161	31,239								
1995	349	4,006	32,697	223	565	745	1995	349	4,006	32,697	223	565	745	1995	349	4,006	32,697	223	565	745	27,574								
1996	108	5,315	33,937	407	350	482	1996	108	5,315	33,937	407	350	482	1996	108	5,315	33,937	407	350	482	21,953								
1997	353	761	19,277	69	491	568	1997	353	761	19,277	69	491	568	1997	353	761	19,277	69	491	568	11,205								
1998	265	3,218	34,173	218	755	1,134	1998	265	3,218	34,173	218	755	1,134	1998	265	3,218	34,173	218	755	1,134	25,728								
1999	221	776	17,262	101	291	440	1999	221	776	17,262	101	291	440	1999	221	776	17,262	101	291	440	12,058								
2000	1,902	1,581	18,272	176	342	582	2000	1,902	1,581	18,272	176	342	582	2000	1,902	1,581	18,272	176	342	582	13,982								
2001	2,398	1,411	17,268	145	684	923	2001	2,398	1,411	17,268	145	684	923	2001	2,398	1,411	17,268	145	684	923	16,982								
2002	1,484	1,196	28,756	38	1,207	1,178	2002	1,484	1,196	28,756	38	1,207	1,178	2002	1,484	1,196	28,756	38	1,207	1,178	17,523								
2003	1,450	1,521	17,493	48	1,590	721	2003	1,450	1,521	17,493	48	1,590	721	2003	1,450	1,521	17,493	48	1,590	721	20,041								
2004	4,247	1,325	20,101	90	1,839	761	2004	4,247	1,325	20,101	90	1,839	761	2004	4,247	1,325	20,101	90	1,839	761	18,922								
2005	4,254	738	13,486	46	1,126	585	2005	4,254	738	13,486	46	1,126	585	2005	4,254	738	13,486	46	1,126	585	19,890								
2006	4,065	2,002	19,014	43	868	444	2006	4,065	2,002	19,014	43	868	444	2006	4,065	2,002	19,014	43	868	444	21,023								
2007							2007							2007															

1995-2007 proportions 3.7% 4.1% 47.7% 0.3% 1.8% 1.6% 40.8% 2.1% 7.6% 54.4% 0.5% 0.8% 3.2% 31.4%

Table 3. Survey stratified mean biomass time series used in the rebuilding potential analysis and in estimating catch limits and targets associated with the median C/B exploitation ratio.

Change in Biomass		Change in Biomass 3 year moving average					Stratified mean biomass (kg/tow)						
Year	Bamdoor	Clearnose Little	Rosette	Smooth	Thorny	Winter	Year	Bamdoor	Clearnose Little	Rosette	Smooth	Thorny	Winter
1964							1964	1.21			0.33		
1965	50%			-34%			1965	1.82			0.48		
1966	-55%			46%			1966	0.81			0.32		
1967	-46%			-32%			1967	0.44		0.02	0.15		2.16
1968	-35%			53%		6%	1968	0.28		0.00	0.39		4.42
1969	-81%			-25%		-34%	1969	0.05		0.00	0.23		1.86
1970	23%			29%		-26%	1970	0.07		0.00	0.29		5.71
1971	157%			-84%		-5%	1971	0.10		0.00	0.16		3.00
1972	-44%			-27%		15%	1972	0.10		0.00	0.33		1.08
1973	-96%			-32%		-23%	1973	0.00		0.02	0.13		4.12
1974	-100%			111%		31%	1974	0.00		0.01	0.31		4.56
1975				-6%		11%	1975	0.02	0.24	0.01	0.12		2.10
1976	181%			-33%		-20%	1976	0.02	0.30	0.00	0.08		3.04
1977	-100%			-39%		-36%	1977	0.05	0.30	0.02	0.04		1.31
1978				478%		-49%	1978	-	0.77	0.02	0.38		4.10
1979				87%		-17%	1979	-	0.16	1.35	0.01		4.28
1980				20%		-37%	1980	0.01	0.42	0.01	0.45		5.12
1981				-64%		70%	1981	-	0.68	0.21	0.18		4.60
1982				766%		324%	1982	-	0.17	1.50	0.08		6.22
1983				-81%		25%	1983	-	0.21	3.63	0.01		3.34
1984				-55%		45%	1984	-	0.14	5.72	0.00		0.65
1985	-60%			2781%		-48%	1985	0.01	0.16	0.00	0.15		12.85
1986	642%			6%		16%	1986	0.00	0.31	4.09	0.03		13.32
1987	-53%			0%		43%	1987	0.00	0.54	2.75	0.00		2.88
1988	-46%			-54%		34%	1988	0.01	0.32	4.63	0.00		1.63
1989	-35%			196%		6%	1989	0.00	0.34	5.08	0.02		1.56
1990	479%			-35%		-42%	1990	0.00	0.27	6.63	0.02		1.49
1991	10%			27%		-7%	1991	0.03	0.40	4.99	0.01		1.70
1992	-92%			-14%		29%	1992	0.00	0.92	5.89	0.02		1.63
1993	569%			-41%		-4%	1993	0.00	0.34	5.30	0.03		0.96
1994	-75%			79%		-14%	1994	0.03	0.48	7.52	0.02		3.58
1995	220%			240%		-7%	1995	0.11	0.33	2.87	0.07		1.51
1996	-62%			-90%		-21%	1996	0.04	0.43	3.62	0.04		2.12
1997	149%			10%		-55%	1997	0.10	0.81	7.57	0.18		0.81
1998	-15%			-48%		-27%	1998	0.09	1.12	7.47	0.01		0.23
1999	237%			32%		-36%	1999	0.30	1.05	9.88	0.07		0.65
2000	-4%			-24%		-41%	2000	0.29	1.03	8.60	0.03		0.48
2001	89%			149%		77%	2001	0.54	1.61	6.84	0.12		0.33
2002	43%			118%		-20%	2002	0.78	0.89	6.44	0.05		0.83
2003	-29%			-61%		-19%	2003	0.55	0.66	6.48	0.03		0.19
2004	134%			71%		-23%	2004	1.29	0.71	7.22	0.05		0.74
2005	-20%			-39%		15%	2005	1.04	0.52	3.24	0.06		4.03
2006	13%			61%		-5%	2006	1.17	0.53	3.32	0.06		2.48
2007	-32%			-58%		-32%	2007	0.80	0.85	0.07	0.09		3.71

Rebuilding prospects

For each managed skate species, the response of survey biomass to changes in catch was examined using the same procedures described in Document 5 of the DEIS Appendix I and the above three catch time series. The results are shown in Figure 1 to Figure 14. This analysis shows whether catch or the relative exploitation ratio (C/B) had any measurable effect on biomass. The top graphs of each panel show a linear least squares regression line and the median value. A negative slope is indicative that high catches lead to low biomass, and vice versa, as would be expected. Positive slopes or no slope are counterintuitive meaning among other things that other factors had more influence over changes in biomass than did the estimated catches.

As was recognized in Document 5 of Appendix I, the relationship between changes in biomass and the catch/biomass ratio are not completely independent, because biomass appears in the denominator of ordinate and the numerator of the abscissa. As a result, the null hypothesis that the slope is significantly different than zero is invalid. Instead, an alternative null hypothesis was developed using a randomization procedure to estimate a slope that resulted if the data were chosen on the basis of random choice alone, but are not truly independent variables.

A randomization test was performed where the change in the three year moving average of biomass and the three year moving average catch/biomass ratio were randomly chosen with replacement, over 1000 iterations in a 20 year artificial time series. The red dashed regression line in each time series represents a threshold where the null hypothesis should be rejected with 95% confidence when the realized slope is less (i.e. more negative). The red dot in each figure represents 2007.

In the Amendment 3 draft, smooth, thorny, and winter skates (all three overfished species) were thought to have a significant relationship between catches and changes in biomass, based on the preponderance of data that biomass increased more frequently when the C/B ratio was below the median value. Other skate species had no such relationship or the slope was counter intuitively positive. The lack of a relationship was attributed to uncertainty in the catch time series, or potentially lagged and poorly understood population dynamics.

The new catch time series and the randomizing test for a significant slope changes this perception. For both sets of winter skate catch estimates (Figure 1 and Figure 2), the C/B slope is not significantly different than no relationship (i.e. cannot reject the null hypothesis with 95% confidence), although the biomass increased 17 out of 19 times for an average of a 54% annual increase when the C/B ratio was below the median. This might have more to do with autocorrelations, because the years with high biomass in the mid-1980s are all clustered below the C/B median.

Thorny skate (Figure 3 and Figure 4) exhibits a flat slope and essentially no relationship between these values, for either catch time series. On the other hand, there does seem to be a significant relationship between C/B and changes in survey biomass for smooth skate (Figure 5 and Figure 6). This relationship appears to be driven by just five years of data, and for the rest of the time series there appears to be no difference in changes in biomass at high catch rates vs. low catch rates.

For rosette skate, the slope between the C/B ratio and changes in biomass are significant and negative, but again this appears to be driven by just two points, which may be related to transient oceanographic conditions in two survey years. Little, clearnose, and barndoor skate all exhibit a flat, non-significant slope.

Even though the relationship between the C/B ratio and changes in skate biomass appear in some cases to make sense and indicate that low catches are more likely than not to cause increases in biomass

and rebuilding, none of the relationships are very strong and are probably not very predictive of rebuilding potential at lower catch levels.

Calculation of catch limits




Catch limits and targets defined by the median catch/biomass ratio applied to the annual 3 year moving averages for survey biomass and aggregated over species are shown in Table 4 to Table 8. Using the Draft Amendment 3 catch time series, the perception was that in 2006, catch was close to the target (ACT) and landings were slightly above the TAL. Landings in 2007 had however exceeded the 2007 TAL (the TAL declined due to lower stratified mean biomass⁷ values) and landings were approaching the catch target (which includes both landings and discards). It was anticipated that the discard rate in 2007 would be the same as that in 2006, or might have declined from the effects of Framework 42. Due to the increasing landings in 2007 it was however anticipated that the total catch would be above the ABC and that reductions in landings and catch were required. Amendment 3 proposed alternatives to reduce 2007 landings to the TAL. To meet the target, wing fishery landings would need to decline by 45% and bait fishery landings by 43%⁸.

In contrast, the new discard estimates for 2004-2006 are substantially higher than previous estimates. As in the Amendment 3 DEIS, 2007 landings are near the catch target (or ACT). Instead of declining by 65%, the new discard estimates are flat or even increasing in recent years. Thus the fraction of total catch attributable to discards is much higher using these new estimates and results in a much lower TAL. Without action to reduce skate discards, the analyses using the new catch data (Figure 16 and Figure 17) indicate that it would take a 57-60% reduction in skate wing landings and a 63-65% reduction in skate bait landings to prevent the catch from exceeding the ACT.

⁷ A considerable portion of the survey biomass decline arises from 2004 dropping out of the three year moving average.

⁸ These re-estimated TALs are slightly

Figure 1. Relationship for winter skate between three year moving average of catch (length composition method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change
41,320	All	Up 19 Down 16
18,255	Above median	Average Δ 2004-2006
14,604	Below Median	16 16.5%
Percentile	Below 80%	8 12 0.7%
		11 4 42.9%
		5 2 30.5%
		-6%
		-25%

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change
9.26	All	Up 19 Down 16
4.12	Above median	Average Limit (mt) Δ 2004-2006
3.09	Below Median	16 16.5%
Percentile	Below 75%	2 14 -27.2%
		17 2 54.5%
		6 1 49.5%
		12,087
		9,065
		-38%
		-53%

Figure 2. Relationship for winter skate between three year moving average of catch (selectivity_ogive_method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change
All	All	Up 19, Down 16
41,320	Above median	Average Δ 2004-2006 16.5%
16,586	Below median	10.4%
13,269	Below 80%	5 33.9% -7%
Percentile	35%	7 2 44.5% -25%

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change
All	All	Up 19, Down 16
8.02	Above median	Average Limit (mt) Δ 2004-2006 16.5%
4.12	Below Median	2 14 -27.2%
3.09	Below 75%	17 2 54.5% 12,087 -32%
Percentile	24%	9 1 52.4% 9,065 -49%

Figure 3. Relationship for thorny skate between three year moving average of catch (length composition method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change
	All	Up 11 Down 26 Average Δ 2004-2006 -11.8%
32,864	Above median	6 10 -9.5%
Median	Below Median	5 16 3.5% 524%
80% of median	Below 80%	4 12 2.8% 399%
Percentile	38%	

Catch/biomass ratio

Limit	Threshold	Biomass change
	All	Up 11 Down 26 Average Limit (mt) Δ 2004-2006 -11.8%
Maximum	Above median	5 13 -12.9%
Median	Below Median	6 13 7.3% 1,258
75% of median	Below 75%	4 10 3.2% 944
Percentile	35%	8%

Figure 4. Relationship for thorny skate between three year moving average of catch (selectivity ogive method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change	Average Δ 2004-2006		
All	All	Up 11	Down 26	-11.8%	
Maximum	32,864	Above median	6	10	-8.5%
Median	5,209	Below Median	5	16	3.5%
80% of med	4,167	Below 80%	3	13	1.9%
Percentile	38%				472%

Catch/biomass ratio

Limit	Threshold	Biomass change	Average Limit (mt)	Δ 2004-2006	
All	All	Up 11	26	-11.8%	
Maximum	7.32	Above median	3	15	-20.7%
Median	3.14	Below Median	8	11	10.5%
75% of med	2.36	Below 75%	4	3	6.5%
Percentile	18%				83%
					37%

Figure 5. Relationship for smooth skate between three year moving average of catch (length composition method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change
All	All	Up 15, Down 22, Average Δ 2004-2006 9.6%
Maximum	790	Above median 5, Below median 11, Average Δ 2004-2006 4.2%
Median	345	Above median 10, Below median 11, Average Δ 2004-2006 36.3%
80% of median	276	Above median 5, Below median 6, Average Δ 2004-2006 17.9%
Percentile	25%	

Catch/biomass ratio

Limit	Threshold	Biomass change
All	All	Up 15, Down 22, Average Δ 2004-2006 9.6%
Maximum	4.55	Above median 4, Below median 14, Average Δ 2004-2006 -11.2%
Median	1.95	Above median 11, Below median 8, Average Δ 2004-2006 40.8%
75% of median	1.46	Above median 9, Below median 1, Average Δ 2004-2006 70.2%
Percentile	26%	

Figure 6. Relationship for smooth skate between three year moving average of catch (selectivity ogive method) and biomass with no lag.






Catch thresholds and historic change in biomass

Catch	Limit (mt)	Threshold	Biomass change		Average	Δ2004-2006
			Up	Down		
Maximum	790	Above median	5	11	23.8%	-6%
Median	303	Below Median	10	11	19.5%	-25%
80% of medi:	242	Below 80%	8	2	39.7%	
Percentile		24%				

Catch/biomass ratio	Limit (mt)	Threshold	Biomass change		Average	Limit (mt)	Δ2004-2006
			Up	Down			
Maximum	3.79	Above median	1	17	-22.9%	243	-25%
Median	1.69	Below Median	14	5	48.1%	183	-43%
75% of medi:	1.27	Below 75%	9	0	95.0%		
Percentile		24%					

Figure 7. Relationship for rosette skate between three year moving average of catch (length composition method) and biomass with no lag.






Catch thresholds and historic change in biomass

Catch	Limit (mt)	Threshold	Biomass change		Average	Δ2004-2006
			Up	Down		
Maximum	228	All	22	15	46.5%	
Median	41	Above median	10	6	60.8%	
80% of medicit	33	Below 80%	12	9	50.4%	6%
Percentile		22%	7	3	49.7%	-16%

Catch/biomass ratio		Biomass change		Average	Limit (mt)	Δ2004-2006
Maximum	Median	Up	Down			
26.36	1.81	22	15	46.5%		
1.36	1.36	7	11	2.0%		
	31%	15	4	96.8%	118	203%
		9	3	116.1%	88	128%

Figure 8. Relationship for rosette skate between three year moving average of catch (selectivity_ogive_method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change Up	Biomass change Down	Average Δ 2004-2006
Maximum	All	22	15	46.5%
Median	Above median	10	6	60.9%
80% of median	Below Median	12	9	50.4%
Percentile	Below 80%	8	4	45.3%
		29%		-43%
				-55%

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change Up	Biomass change Down	Average Limit (mt)	Δ 2004-2006
Maximum	All	22	15	46.5%	70%
Median	Above median	7	11	8.4%	143
75% of median	Below Median	15	4	87.1%	107
Percentile	Below 75%	6	2	129.6%	28%
		21%			

Figure 9. Relationship for little skate between three year moving average of catch (length composition method) and biomass with no lag.






Catch thresholds and historic change in biomass

Catch	Limit (mt)	Threshold	Biomass change		Average	Δ2004-2006
			Up	Down		
Maximum	19,866	All	27	10	20.1%	
Median	10,189	Above median	22	9	19.8%	
80% of medi	8,151	Below Median	5	1	29.0%	-7%
Percentile		Below 80%	2	1	7.5%	-26%

Catch/biomass ratio		Biomass change		Average	Limit (mt)	Δ2004-2006
Up	Down	Up	Down			
7.27	2.39	16	7	20.1%		
1.79	1.79	11	3	17.2%	5,230	-52%
25%	25%	7	1	31.1%	3,922	-64%

Figure 10. Relationship for little skate between three year moving average of catch (selectivity ogive method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change	Average Δ 2004-2006
Maximum	All	Up 27	20.1%
Median	Above median	Down 22	19.8%
80% of medi	Below Median	Up 5	29.0%
Percentile	Below 80%	Down 2	7.5%

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change	Average Limit (mt)	Δ 2004-2006
Maximum	All	Up 27	10	20.1%
Median	Above median	Down 16	7	17.2%
75% of medi	Below Median	Up 11	3	31.1%
Percentile	Below 75%	Down 6	1	31.1%

Figure 11. Relationship for cleamose skate between three year moving average of catch (length composition method) and biomass with no lag.






Catch thresholds and historic change in biomass

Catch	Limit (mt)	Threshold	Biomass change		Average Δ 2004-2006
			Up	Down	
Maximum	3,334	All	24	13	12.0%
Median	969	Above median	21	6	18.8%
80% of med	776	Below Median	3	7	22.1%
Percentile		Below 80%	1	2	1.3%
					-13%

Catch/biomass ratio		Biomass change		Average Limit (mt)	Δ 2004-2006
Limit (mt)	Threshold	Up	Down		
Maximum	5.89	Above median	18	7	7.5%
Median	2.25	Below Median	6	6	34.9%
75% of med	1.69	Below 75%	5	6	27.2%
Percentile					59%
					19%

Figure 12. Relationship for cleansse skate between three year moving average of catch (selectivity ogive method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change	Up	Down	Average	Δ2004-2006
Maximum	All	Down	24	13	12.0%	
Median	Above median	Up	21	8	14.4%	
80% of medic	Below Median	Down	3	5	27.6%	-2%
Percentile	Below 80%	Down	3	4	31.5%	-22%

Catch/biomass ratio

Limit	Threshold	Biomass change	Up	Down	Average	Limit (mt)	Δ2004-2006
Maximum	All	Down	24	13	12.0%		
Median	Above median	Up	18	7	7.7%		
75% of medic	Below Median	Down	6	6	34.9%	1,551	36%
Percentile	Below 75%	Down	3	3	16.9%	1,164	2%

Figure 13. Relationship for barndoor skate between three year moving average of catch (length composition method) and biomass with no lag.



Catch thresholds and historic change in biomass




Catch

Limit (mt)	Threshold	Biomass change
All	All	Up 21, Down 16, Average Δ 2004-2006 58.5%
6,514	Above median	4 108.0%
400	Below Median	7 12 63.9%
320	Below 80%	5 9 57.6%
38%	Percentile	-88%
		-91%

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change
All	All	Up 21, Down 16, Average Limit (mt) Δ 2004-2006 58.5%
64.47	Above median	8 14 0.7%
4.32	Below Median	13 2 150.9%
3.24	Below 75%	8 2 128.5%
34%	Percentile	4,332 25%
		3,249 -7%

Figure 14. Relationship for barndoor skate between three year moving average of catch (selectivity ogive method) and biomass with no lag.



Catch thresholds and historic change in biomass

Catch

Limit (mt)	Threshold	Biomass change
All	All	Up 21, Down 16, Average Δ 2004-2006 58.5%
Maximum	6,514	Above median 13, Below median 5, Average Δ 2004-2006 106.6%
Median	400	Below median 8, Above median 11, Average Δ 2004-2006 64.3%
80% of median	320	Below 80% 8, Above 80% 9, Average Δ 2004-2006 71.8%
Percentile	45%	

Catch/biomass ratio

Limit (mt)	Threshold	Biomass change
All	All	Up 21, Down 16, Average Δ 2004-2006 58.5%
Maximum	64.47	Above median 8, Below median 14, Average Δ 2004-2006 -2.2%
Median	3.23	Below median 13, Above median 2, Average Δ 2004-2006 155.1%
75% of median	2.42	Below 75% 11, Above 75% 2, Average Δ 2004-2006 163.0%
Percentile	42%	

Table 4. Calculation of alternative skate catch limits using catch and catch/biomass medians from Draft Amendment 3, using corrected discards.

Species	Catch	C/B derived catch limits		
	Median	80% of medi	Median	75% of medi
Barndoor	290	232	2,685	2,014
Clearnose	521	417	614	460
Little	17,524	14,019	7,649	5,737
Rosette	26	21	56	42
Smooth	33	26	27	20
Thorny	155	124	50	38
Winter	17,422	13,938	11,530	8,648
Total	35,971	28,777	22,612	16,959
Discards	13,734	10,987	8,634	6,475
Prohibited species	430	344	2,486	1,865
Legal species	13,305	10,644	6,147	4,611
Discard rate legal sp	37%	37%	31%	31%
Allowable landings	22,237	17,789	13,978	10,484
Prohibited species	48	38	276	207
Legal species	22,189	17,751	13,702	10,277
Wing fishery TAL	15,502	12,402	10,351	7,763
Change from 2007	10%	-12%	-26%	-45%
Bait fishery TAL	6,735	5,388	3,627	2,721
Change from 2007	41%	13%	-24%	-43%
TAL	22,237	17,789	13,978	10,484
Discards	13,734	10,987	8,634	6,475
TAC	35,971	28,777	22,612	16,959
Change from 2007	39%	11%	-13%	-34%

Figure 15. Trend in annual ABC, ACT, and TALs derived from applying the median catch/biomass ratio from Draft Amendment 3 catches to historic stratified mean biomass by skate species.




Table 5. Input variables and results application of catch/biomass ratios derived from Draft Amendment 3 catches and applied to stratified mean survey biomass.

Species	Catch/biomass index (thousand mt catch/kg per tow)		Stratified mean survey weight (kg/tow)		
	Median	75% of median	2004-2006	2005-2007	Target
Barndoor	2.68	2.01	1.17	1.00	1.62
Clearnose	0.97	0.73	0.59	0.63	0.56
Little	3.50	2.62	4.59	3.67	6.54
Rosette	0.86	0.65	0.06	0.06	0.03
Smooth	0.19	0.14	0.19	0.14	0.31
Thorny	0.12	0.09	0.55	0.42	4.41
Winter	3.93	2.95	3.04	2.93	6.46
Annual catch limit (ACL/ABC)			31,858	22,612	53,731
Annual catch target (ACT)			23,893	16,959	40,298
Total allowable landings (TAL)			14,770	10,484	29,912

Table 6. Calculation of alternative skate catch limits using catch and catch/biomass medians from the Data Poor Assessment Workshop length composition method.

Species	Catch		C/B derived catch limits	
	Median	80% of medi	Median	75% of medi
Barndoor	400	320	4,328	3,246
Clearnose	969	776	1,428	1,072
Little	10,189	8,151	5,230	3,917
Rosette	41	33	77	88
Smooth	345	276	281	210
Thorny	5,479	4,383	1,257	943
Winter	18,255	14,604	12,092	9,069
Total	35,678	28,543	24,692	18,546
Discards	20,699	16,559	14,325	10,759
Prohibited species	5,602	4,481	5,280	3,959
Legal species	15,097	12,078	9,046	6,800
Discard rate legal sp	51%	51%	48%	48%
Allowable landings	14,979	11,984	10,367	7,786
Prohibited species	622	498	587	440
Legal species	14,357	11,486	9,780	7,346
Wing fishery TAL	11,399	9,120	8,022	6,027
Change from 2007	-19%	-35%	-43%	-57%
Bait fishery TAL	3,580	2,864	2,341	1,759
Change from 2007	-25%	-40%	-51%	-63%
TAL	14,979	11,984	10,363	7,786
Discards	20,699	16,559	14,325	10,759
TAC	35,678	28,543	24,688	18,546
Change from 2007	-16%	-33%	-42%	-56%

Figure 16. Trend in annual ABC, ACT, and TALs derived from applying the median catch/biomass ratio from catches using the length composition method to assign catches and apply them to historic stratified mean biomass by skate species.




Table 7. Input variables and results application of catch/biomass ratios derived from length composition method catches and applied to stratified mean survey biomass.

Species	Catch/biomass index (thousand mt catch/kg per tow)		Stratified mean survey weight (kg/tow)		
	Median	75% of median	2004-2006	2005-2007	Target
Barndoor	4.32	3.24	1.17	1.00	1.62
Clearnose	2.25	1.69	0.59	0.63	0.56
Little	2.39	1.79	4.59	3.67	6.54
Rosette	1.18	1.36	0.06	0.06	0.03
Smooth	1.95	1.46	0.19	0.14	0.31
Thorny	2.96	2.22	0.55	0.42	4.41
Winter	4.12	3.09	3.04	2.93	6.46
Annual catch limit (ACL/ABC)			31,945	24,688	64,196
Annual catch target (ACT)			23,977	18,546	48,145
Total allowable landings (TAL)			10,067	7,786	20,213

Table 8. Calculation of alternative skate catch limits using catch and catch/biomass medians from the Data Poor Assessment Workshop selectivity ogive method.

Species	Catch Median	C/B derived catch limits		
		80% of medi	Median	75% of medi
Barndoor	400	320	3,236	2,425
Clearnose	1,110	888	1,548	1,161
Little	10,189	8,151	5,230	3,917
Rosette	47	38	142	107
Smooth	303	242	243	183
Thorny	5,209	4,167	1,334	1,002
Winter	16,586	13,269	12,092	9,069
Total	33,844	27,075	23,825	17,864
Discards	19,962	15,969	14,052	10,536
Prohibited species	5,321	4,256	4,332	3,249
Legal species	14,641	11,713	9,720	7,287
Discard rate legal sp	52%	52%	51%	51%
Allowable landings	13,882	11,106	9,773	7,328
Prohibited species	591	473	481	361
Legal species	13,291	10,633	9,292	6,967
Wing fishery TAL	10,419	8,336	7,532	5,648
Change from 2007	-26%	-41%	-47%	-60%
Bait fishery TAL	3,463	2,770	2,241	1,679
Change from 2007	-27%	-42%	-53%	-65%
TAL	13,882	11,106	9,773	7,328
Discards	19,962	15,969	14,052	10,536
TAC	33,844	27,075	23,826	17,864
Change from 2007	-19%	-35%	-43%	-57%

Figure 17. Trend in annual ABC, ACT, and TALs derived from applying the median catch/biomass ratio from catches using the selectivity ogive method to assign catches and apply them to historic stratified mean biomass by skate species.




Table 9. Input variables and results application of catch/biomass ratios derived from selectivity ogive method catches and applied to stratified mean survey biomass.

Species	Catch/biomass index (thousand mt catch/kg per tow)		Stratified mean survey weight (kg/tow)		
	Median	75% of median	2004-2006	2005-2007	Target
Barndoor	3.23	2.42	1.17	1.00	1.62
Clearnose	2.44	1.83	0.59	0.63	0.56
Little	2.39	1.79	4.59	3.67	6.54
Rosette	2.19	1.65	0.06	0.06	0.03
Smooth	1.69	1.27	0.19	0.14	0.31
Thorny	3.14	2.36	0.55	0.42	4.41
Winter	4.12	3.09	3.04	2.93	6.46
Annual catch limit (ACL/ABC)			30,898	23,826	63,240
Annual catch target (ACT)			23,162	17,864	47,462
Total allowable landings (TAL)			9,501	7,328	19,469

