

#2-SALMON

**US Atlantic Salmon Stock Structure, Status, and Metrics Information Brief
For New England Council to amend the Atlantic Salmon FMP in 2010
DRAFT: 4 June 2010**

John F. Kocik
Supervisory Research Fishery Biologist
DOC/NOAA/NMFS/NEFSC F/NEC311
NOAA Fisheries Service Maine Field Station
17 Godfrey Drive Suite 1
Orono, ME 04473-3693
John.Kocik@NOAA.GOV

SUMMARY: This species, unlike other NEFMC managed stocks, is not subject to a plan review by the National Marine Fisheries Service because the current FMP prohibits possession of Atlantic salmon and any directed or incidental (bycatch) commercial fishery for Atlantic salmon in federal waters. Due primarily to significant ongoing threats from poor marine survival, dams, and the inadequacy of existing regulatory mechanism for dams, the Gulf of Maine Distinct Population Segment (GOM DPS) of Atlantic salmon was listed as endangered under the Endangered Species Act (ESA) in June 2010. The other stock complexes in the US - Long Island Sound and Central New England and the trans-boundary Outer Bay of Fundy complex are also well below conservation limits. A metric used by salmon managers, Conservation Spawning Escapement, is comparable to minimum stock size threshold (MSST) and under current natural mortality regimes US salmon stock complexes are below the minimum stock size at which rebuilding to BMSY will occur within 10 years. Since 1967, CSE has not been achieved for a stock complex and the highest abundances noted are less than 25% of these conservation minima and the median values are less than 10% -even with extensive hatchery support.

Stock Structure and Management Units

Atlantic salmon (*Salmo salar*) are circumpolar in distribution and in North America range from Ungava Bay southward to Long Island Sound. As a consequence of human development, many native United States populations were extirpated. Salmon life history is complex owing to its use habitat as diverse as headwater streams and distant water marine habitats. US Atlantic salmon spawn in rivers, spending 1-3 years in nursery streams. When juveniles (parr) exceed 13 cm in the autumn, they develop into smolts and migrate to the Atlantic Ocean the following spring. Tagging data indicates that US salmon commonly migrate as far north as West Greenland and return after 1-3 winters at sea. Non-maturing adults remain at sea feeding in the coastal waters of West Greenland, Newfoundland and Labrador.

Strong homing capabilities of Atlantic salmon foster the formation and maintenance of local breeding groups- stocks. These stocks exhibit heritable adaptations to their home range in rivers and likely at sea. The importance of maintaining local adaptations has

demonstrated utility in salmon conservation. Because of significant declines in Atlantic salmon populations in the US, analyses of population structure was conducted and some populations are managed under the Endangered Species Act (ESA, 74 Federal Register 29346, June 19, 2009). The Act required that subgroups must be separable from the remainder of and significant to the species to which it belongs to warrant ESA protection (Fay et al. 2006). Biologists then delineated US Atlantic salmon populations into four discrete stock complexes that are now managed in differing manners: 1) Long Island Sound stock complexes; 2) Central New England stock complexes; 3) Gulf of Maine Distinct Population Segment and the 4) Outer Bay of Fundy Salmon Fishing Area (Figure 1). Both the Long Island Sound and Central New England stock complexes native stocks were extirpated in the 1800's. Remnant native populations of Atlantic salmon in the United States now persist only in Maine in the endangered Gulf of Maine DPS and cross-boundary headwater populations from the Outer Bay of Fundy SFA.

Currently, the Gulf of Maine DPS is managed under a recovery program under the ESA (NMFS and FWS 2005). The Long Island Sound and Central New England stock complexes are under restoration using donor stocks from Maine and are managed under coordinated Federal and Interstate restoration efforts, in the form of stocking and fish passage management. While, Atlantic salmon stocks from the Penobscot River in Maine were used in the restoration programs in the Connecticut (Long Island Sound DPS) and in the Merrimack and Saco in the (Central New England DPS) these programs are now genetically isolated. US watersheds in the Outer Bay of Fundy region are supplemented by St. John River Atlantic salmon broodstock and the core populations of this management unit have freshwater nursery areas primarily in Canadian watersheds. Within Canada, the Committee on the Status of Endangered Wildlife in Canada (COSEWIC) assessed population structure and status and is moving forward with a species level assessment of Atlantic salmon in eastern Canada expected to be complete by November, 2010 and the outer Bay of Fundy (oBoF) stock complex is currently noted as of particular concern warranting proactive recovery planning.

Stock Status

US Atlantic salmon populations are assessed by the US Atlantic Salmon Assessment Committee (USASAC), a team of state and federal biologists tasked with compiling data on the species throughout New England and reporting population status. Annual assessments are available on-line (<http://www.nefsc.noaa.gov/USASAC/> USASAC 2010). Currently population status of salmon is determined by 1) counting returning adults either directly (traps and weirs) or 2) indirectly using redd surveys. Total returns also include fish retained from angling in some states and historical US time series includes these data (USASAC 2010).

The modern time series of salmon returns to US Rivers starts in 1967 (Figure 2). Average annual Atlantic salmon returns to US rivers from 1967 to present was 2,173 and the median is 1,670. Because many of the populations in southern New England were extirpated and the Penobscot River was at very low levels, the salmon returns

graph illustrates the sequential rebuilding of the populations through restoration efforts in the 1970s – with increased abundance first seen in the Penobscot River then the Merrimack and Connecticut Rivers. Unfortunately, the trajectory of this recovery did not continue in the late 1980s and early 1990s. Starting in the early 1990s there was a phase shift in marine survival and an overall reduction in marine survival occurred in all US and most Canadian populations. Average annual Atlantic salmon returns to US rivers from 1991 to present is 1,878 only 86% of the time series average. There has been a downward trend in production of salmon on both side of the Atlantic (particularly populations dominated by 2SW fish) that have affected US populations. In addition, recovery from historical impacts was never sufficient so US populations were at low absolute abundance when the phase shift began.

Returns to US waters in 2009 were 2,336 fish ranking 17th out of the 43 year time series which is nearly 1,000 fish above complete series median and only the second year above median since 1998. Relative to the average during the current marine survival regime (1991-present), returns were 6th highest out of the 19 years. Given consistency in stocking levels and natural smolt production measurements, increased marine survival is thought to be the primary cause of this increase.

To assess the relative status of the stocks, it is informative to examine conservation spawning escapement (CSE) goals. CSE targets also called conservation limits, are widely used (e.g., ICES 2005) to describe the status of Atlantic salmon populations. Because juvenile rearing habitat can be measured or estimated efficiently, these data are used to calculate target spawning requirements to fully seed all juvenile rearing habitats. US CSE goals are set for accessible rivers using a method developed by Elson (1975). This method assumes a target egg deposition of 2.4 eggs/m² is needed to fully seed a river (Elson 1975); female fecundity averages 7,200 eggs/female (Baum and Meister 1971, Baum 1997); Healthy salmon populations often greatly exceed CSE and as such this metric should be thought of as a minimal conservation target not fishery target. In self-sustaining populations, the number of returns would frequently exceed CSE by 50 to 100 percent allowing for sustainable harvests and buffers against losses between returns and spawning.

For the 3 stock complexes managed primarily by the United States, the CSE goal has been calculated at 29,199 spawners (Table 1). The average percent of the CSE achieved for the modern time series averaged 7.4% and 2009 returns were 8.0% of CSE. In the last decade, total returns have accounted for less than 2 percent of goals for Long Island Sound and Central New England stock complexes. However, salmon returns to the Gulf of Maine DPS have been as high as 20% of CSE during this period, largely due to hatchery smolt returns to the Penobscot River. In smaller rivers of the Gulf of Maine stock complex CSE ranged from 3-15%.

As such, the status of US Atlantic salmon populations is critically low for all stock complexes, with the remnant populations of the Gulf of Maine stock complex listed as endangered. Since the modern era of Atlantic salmon restoration and recovery (circa 1967) began, instances of spawners meeting a replacement rate of 1 let alone

exhibiting population growth are infrequent at a population level and non-existent at a stock complex level. Because recovery and restoration plans have goals of self-sustaining (i.e., exclusively naturally reproducing) populations, the failure to meet replacement with extensive hatchery stocking support indicates that these populations are well below natural replacement rates. This indicates all US stock complexes are under the minimum stock size threshold (MSST) under current natural mortality regimes because they are below the minimum stock size at which rebuilding to BMSY will occur within 10 years.

Fisheries and Current Management Structures

The ocean migrations of US Atlantic salmon take them to distant-water feeding grounds in the sub-Arctic where they overlap with Canadian and European stock complexes. Historically, fisheries for Atlantic salmon have been in coastal environments – both close to natal rivers and in distant coastal waters outside the US Exclusive Economic Zone (EEZ). US stocks were primarily harvested gill nets, weirs, and pound nets in the coastal waters of the United States, West Greenland, Newfoundland, and Labrador. Some catch also occurred in the Bay of Fundy, Scotian Shelf and East Greenland. As such, US Atlantic salmon become part of a mixed-stock marine resource that ventures into the fisheries zones of other countries where they may be exploited. Recognizing this reality, the United States joined the North Atlantic Salmon Conservation Organization (NASCO). NASCO is an international organization, established by an inter-governmental Convention in 1984 with the objective to conserve, restore, enhance and rationally manage Atlantic salmon through international cooperation taking account of the best available scientific information. To date, regulatory and other measures established by NASCO and its Parties have greatly reduced harvests of wild Atlantic salmon all around the North Atlantic.

The last commercial fishery for Atlantic salmon in the US was closed after the 1948 season in the Penobscot River, Maine (Fay et. al. 2006). Although never commonly targeted in offshore waters, regulations have been in place since 1987 restricting ocean harvest. According to the Atlantic salmon fishery management plan (FMP) - "The management unit for the Atlantic salmon FMP is intended to encompass the entire range of the species of U.S. origin while recognizing the jurisdictional authority of the signatory nations to NASCO" (NEFMC 1987). Accordingly, the management unit for this FMP is: "All anadromous Atlantic salmon of U.S. origin in the North Atlantic area through their migratory ranges except while they are found within any foreign nation's territorial sea or fishery conservation zone (or the equivalent), to the extent that such sea or zone is recognized by the United States." Presently there is a prohibition on the possession of Atlantic salmon in the EEZ. Effectively this protects all US populations in marine waters and is complementary to management practiced by the states in riverine and coastal waters.

Distant water fisheries must be considered to effectively conserve and restore US salmon populations. Commercial fisheries for Atlantic salmon in Canada and

Greenland are managed under the auspices of the North Atlantic Salmon Conservation Organization (NASCO), of which the United States is a member. The mixed-stock fisheries in Canada were managed by time-area closures and quotas, however all commercial fisheries for Atlantic salmon in Canada have been closed since 2000. The Greenland fishery has been managed by a quota system since 1972. In 1993, a modified quota system was agreed to, which provided a framework for quotas based on a forecast model of salmon abundance. From 1993-1994, quotas were bought out through a private initiative, but the fishery resumed in 1995 under forecast modeling-based quotas. In 2002, salmon conservationists and the Organization of Fishermen and Hunters in Greenland signed a five-year, annually renewable agreement, which suspended all commercial salmon fishing within Greenland territorial waters while allowing for an annual internal use only fishery. In 2007, a similar agreement was signed and will be in effect through 2013.

Within US waters directed fishing for other species does have the potential to intercept salmon as by-catch. Beland (1984) reported that fewer than 100 salmon per year were caught incidental to other commercial fisheries in the coastal waters of Maine. In most modern fisheries, the surface oriented pelagic distribution of Atlantic salmon would minimize encounters. Using resources from the Standardized Bycatch Reporting Methodology confirms only rare occurrences of Atlantic salmon in observer databases. For the period of record 1989-2009, Atlantic salmon were observed in 5 of 12 years. A total of 44.8 pounds of Atlantic salmon were reported about 50% was discarded. A typical catch was documented in NEFSC (2010) – 9 pounds retained from statistical area 514, coastal Massachusetts from a large mesh sink gillnet in the July 2008 – June 2009 time period. Most observations appear to be single fish. Recent investigations also confirm the potential for Atlantic salmon bycatch to be occurring in pelagic and midwater fisheries due to an overlap in space and time of the fish and gear. As noted above there are few observations of it actually happening. The by-catch of Atlantic salmon in herring fisheries is not considered a significant mortality source for U.S. stocks (ICES 2004). This analysis was conducted in 2004 and with increased observer coverage in the midwater trawl Atlantic herring fleet the ability to assess this should be increased as this time-series grows.

Biological Reference Points

Biological reference points for US Atlantic salmon are different from most other assessed species because they are managed in numbers not biomass at the population level. Also, triggers are different because 1) they are a protected species and 2) none of the historic domestic fisheries that targeted salmon are still active. Conventional fisheries targets (MSY, BMSY, FMSY, etc.) have not been developed because current populations are so low relative even to sustainable conservation levels. A proxy for MSST for US Atlantic salmon is CSE since this provides the minimum population number needed to fully utilize available freshwater nursery habitat. As noted above, since 1967 CSE has not been achieved and the highest abundances noted are less than 25% of these conservation minima even with hatchery support.

Table 1 Two-sea winter (2SW) conservation spawning escapement requirements for US River populations and 2SW returns (with % of CSE) in 2009

| <u>DPS or Other Composite</u> | <u>CSE</u> | <u>Returns 2009 (%)</u> |
|-------------------------------|------------|-------------------------|
| Long Island Sound | 10,094 | 75 (0.7%) |
| Central New England | 3,435 | 92 (2.7%) |
| Gulf of Maine DPS | 15,670 | 2,169 (13.8%) |
| Total | 29,199 | 2,336(8%) |




Figure 1. Map of New England Atlantic Salmon management stock complexes by region from north to south- Outer Bay of Fundy (OBF), Gulf of Maine DPS (GoM), Central New England (CNE), and Long Island Sound (LIS) Regions




Figure 2. Estimated total returns to New England 1967-2009 from USASAC databases for Outer Bay of Fundy (OBF), Central New England (CNE), and Long Island Sound (LIS) Regions and the Gulf of Maine (GoM) Distinct Population Segment.

Literature Cited

Beland, K. F. 1984. Strategic plan for management of Atlantic salmon in the state of Maine. Atlantic Sea Run Salmon Commission. Bangor, ME. 92 pp.

Elson, P. F. 1975. Atlantic salmon rivers, smolt production and optimal spawning: an overview of natural productions. In New England Atlantic salmon restoration conference 1975. J.R. Bohhe and L. Sochasky, eds. International Atlantic Salmon Foundation Special Publication Series 6: 96-119.

Fay, C., Bartron, M., Craig, S., Hecht, A., Pruden, J., Saunders, R., Sheehan, T., and Trial, J. 2006. Status Review for Anadromous Atlantic Salmon (*Salmo salar*) in the United States. Report to the National Marine Fisheries Service and Wildlife Service. <http://www.nmfs.noaa.gov/pr/pdfs/statusreviews/atlanticsalmon.pdf>

ICES (International Council for the Exploration of the Sea). 2004. Report of the Working Group on North Atlantic Salmon . 29 March - 8 April 2004 Halifax, Canada. ICES CM 2004/ACFM:20. ICES. Copenhagen, Denmark. 293 pp.

ICES (International Council for the Exploration of the Sea). 2005. Report of the Working Group on North Atlantic Salmon . 5-14 April 2005, Nuuk, Greenland. ICES CM 2005/ACFM:17. ICES. Copenhagen, Denmark. 291 pp.

NMFS (National Marine Fisheries Service) and FWS (Fish and Wildlife Service). 2005. Final Recovery Plan for the Gulf of Maine Distinct Population Segment of Atlantic Salmon (*Salmo salar*). National Marine Fisheries Service, Silver Spring, MD. <http://www.nero.noaa.gov/nero/hotnews/salmon/index.html>

NEFMC (New England Fishery Management Council). 1987. Fishery management plan for Atlantic salmon: incorporating an environmental assessment and regulatory impact review/initial regulatory flexibility analysis . New England Fishery Management Council. Saugus, MA. 64 pp. http://nefmc.org/salmon/fmp/salmon_original_fmp_oct1987.pdf

NEFSC (Northeast Fisheries Science Center). 2010. Standardized Bycatch Reporting Methodology Annual Discard Report 2010. Internal document. 1531 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA. 02543-1026 <http://www.nefsc.noaa.gov/femad/fsb/SBRM%20Annual%20Discard%20Report/SBRM%20Annual%20Discard%20Reports.htm>

USASAC (United States Atlantic Salmon Assessment Committee). 2010. Annual Report of the U.S. Atlantic Salmon Assessment Committee, Report No. 22 - 2009 Activities. Portland , Maine USA <http://www.nefsc.noaa.gov/USASAC/2010%20USASAC%20Report/USASAC2010-Report%2322-2009-Activities.pdf>