



New England Fishery Management Council

50 WATER STREET | NEWBURYPORT, MASSACHUSETTS 01950 | PHONE 978 465 0492 | FAX 978 465 3116  
John Pappalardo, *Chairman* | Paul J. Howard, *Executive Director*

**MEMORANDUM**

**DATE:** August 7, 2009  
**TO:** Science and Statistical Committee  
**FROM:** Groundfish Plan Development Team  
**SUBJECT:** **Groundfish ABCs/OFLs**

1. The Groundfish PDT provided recommended groundfish ABCs to the Committee in a memorandum dated July 13, 2009. Due to an oversight the PDT neglected to calculate the overfishing levels (OFLs) for these same stocks and include them in the memorandum. Enclosure (1) corrects this oversight and compares the OFLs to the ABC recommendations. In general, the OFLs were calculated using the same analytic techniques and assumptions used to calculate ABCs. There are three comments worth noting:
  - a. For index-based stock without a projection, the OFL is constant since it is based on the most recent estimate of stock size.
  - b. Two values are shown for GB yellowtail flounder, reflecting two assessment model runs. The value for Pollock was calculated using the SSC assumption for the 2009 survey index.
  - c. An OFL is not shown for white hake. The white hake projection model is not run by the NEFSC; the PDT requests these model runs from a scientist attached to a foreign university. The PDT chair decided to wait until after the SSC meeting to request this projection in case additional projections are needed as a result of SSC discussions.
2. After distributing the July 13 memorandum, the PDT determined the Atlantic wolffish ABCs (Table 4 in the July 13 memorandum) were calculated without projecting future stock size. Enclosure (2) provides the corrected ABCs, which increase slightly from those provided earlier.
3. Atlantic halibut ABCs are also provided in this memo.

**Groundfish Plan Development Team**

**OFLs and Recommended ABCs for 2010 – 2012 (corrected)**

| Species                            | Stock  | Actual<br>2008 catch | 2010   | 2011   | 2012   | msy    | 2010<br>OFL | 2011<br>OFL | 2012<br>OFL |
|------------------------------------|--------|----------------------|--------|--------|--------|--------|-------------|-------------|-------------|
|                                    |        |                      | ABC    | ABC    | ABC    |        |             |             |             |
| Cod <sup>1</sup>                   | GB     | 5,134                | 4,812  | 5,616  | 6,214  | 31,159 | 6,272       | 7,311       | 8,090       |
| Cod                                | GOM    | 8,499                | 8,530  | 9,012  | 9,018  | 10,014 | 11,089      | 11,715      | 11,742      |
| Haddock <sup>1</sup>               | GB     | 20,901               | 62,515 | 46,784 | 39,846 | 32,746 | 80,007      | 59,948      | 51,150      |
| Haddock                            | GOM    | 1,197                | 1,265  | 1,206  | 1,013  | 1,360  | 1,617       | 1,536       | 1,296       |
| Yellowtail Flounder <sup>1,2</sup> | GB     | 1,276                | 2,100  | 2,379  | 2,600  | 9,400  | 7,255       | 8,080       | 8,715       |
| Yellowtail Flounder                | SNE/MA | 504                  | 493    | 687    | 1,003  | 6,100  | 1,553       | 2,174       | 3,166       |
| Yellowtail Flounder                | CC/GOM | 727                  | 863    | 1,041  | 1,159  | 1,720  | 1,124       | 1,355       | 1,508       |
| American Plaice                    | GB/GOM | 1,348                | 3,156  | 3,444  | 3,632  | 4,011  | 4,110       | 4,483       | 4,727       |
| Witch Flounder                     | GB     | 1,063                | 944    | 1,369  | 1,639  | 2,352  | 1,239       | 1,792       | 2,141       |
| Winter Flounder                    | GB     | 963                  | 2,052  | 2,224  | 2,543  | 3,500  | 2,660       | 2,886       | 3,297       |
| Winter Flounder                    | GOM    | 402                  | 238    | 238    | 238    | 917    | 441         | 570         | 685         |
| Winter Flounder                    | SNE/MA | 1,432                | 644    | 897    | 1,198  | 9,742  | 1,568       | 2,117       | 2,830       |
| Redfish                            | GB/GOM | 1,364                | 7,586  | 8,356  | 9,224  | 10,139 | 9,899       | 10,903      | 12,036      |
| White Hake                         | GB/GOM | 1,876                | 2,832  | 3,295  | 3,638  | 5,800  |             |             |             |
| Pollock <sup>1,3</sup>             | GB/GOM | 11,370               | 2,097  | 2,527  | 3,043  | 11,320 | 2,252       | 3,074       | 3,702       |
| Windowpane                         | GOM/GB | 350                  | 169    | 169    | 169    | 700    | 225         | 225         | 225         |
| Windowpane                         | SNE/MA | 363                  | 237    | 237    | 237    | 500    | 317         | 317         | 317         |
| Ocean Pout                         |        | 125                  | 271    | 271    | 271    | 3,754  | 361         | 361         | 361         |
| Atlantic Halibut <sup>1</sup>      |        | 84                   | 71     | 78     | 85     | 3,500  | 119         | 130         | 143         |
| Atlantic Wolffish <sup>4</sup>     |        |                      |        |        |        |        | 92-         | 92-         | 92-         |
|                                    |        |                      |        |        |        |        | 278 - 311   | 192         | 192         |

Notes:

1. All ABCs are for entire stock as assessed by U.S. These values include U.S. and Canadian catch.

2. PDT recommends 2011 and 2012 ABC's be revisited when TRAC results available in future years.

3. Based on index-projection. See PDT memo for comments.

4. No specific PDT recommendation. See PDT July 13, 2009 memo and enclosure (2) for range of possible catches.

Table 4 (corrected)– Atlantic wolffish catches based on different assessment runs (corrected version)

| 2007                             |                    |               |                     |       |         |                            |
|----------------------------------|--------------------|---------------|---------------------|-------|---------|----------------------------|
|                                  | Length at maturity | Total Biomass | Exploitable Biomass | Fmsy  | 75%Fmsy | Fmsy * Exploitable Biomass |
| <b>Run 1<br/>(slope=0.15)</b>    |                    |               |                     |       |         |                            |
| 40 cm                            | 1118.9             | 533.8         | 0.319               | 0.239 | 192     | 144                        |
| 65cm                             | 1118.9             | 533.8         | 0.233               | 0.175 | 140     | 105                        |
| 75 cm                            | 1118.9             | 533.8         | 0.185               | 0.139 | 111     | 83                         |
| <b>Run 2 (L<sub>50</sub>=90)</b> |                    |               |                     |       |         |                            |
| 40 cm                            | 1008.3             | 215.3         | 0.686               | 0.515 | 168     | 126                        |
| 65cm                             | 1008.3             | 215.3         | 0.486               | 0.365 | 119     | 89                         |
| 75 cm                            | 1008.3             | 215.3         | 0.374               | 0.281 | 92      | 69                         |